
GUYANA



	Females		0						
		6 000 LIVES CAN BE S	AVE	D BY 2025 BY IMPLEMENTING ALL OF THE WI	HO "BEST BUYS	3"			
LOD:		NATIONAL TARGET SET			DATA YEAR	MALES	FEMALES	TOTA	
OR	Premature mortality		Т	otal NCD deaths	2016	2 300	2 200	4 400	
	from NCDs	✓	R	tisk of premature death between 30-70 years (%)	2016	32	29	31	
4	Suicide mortality	-	S	uicide mortality rate (per 100 000 population)	2016	-	-	29	
ISK	FACTORS								
	Harmful use of alcohol	√		otal alcohol per capita consumption, adults aged 15+ itres of pure alcohol)	2016	11	2	6	
*	Physical inactivity	✓	P	Physical inactivity, adults aged 18+ (%)	2016				
)	Salt/Sodium intake	✓	٨	Mean population salt intake, adults aged 20+ (g/day)	2010	7	6	6	
9	Tobacco use	✓	C	Current tobacco smoking, adults aged 15+ (%)	2016				
8	Raised blood pressure	X	R	laised blood pressure, adults aged 18+ (%)	2015	21	18	20	
Å	Diabetes	X	R	laised blood glucose, adults aged 18+ (%)	2014	7	11	9	
B	Obseiter	V	C	Dbesity, adults aged 18+ (%)	2016	12	26	19	
W.	Obesity	X		Obesity, adolescents aged 10-19 (%)	2016	8	9	9	
V	Ambient air pollution	•		exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2	
•	Household air pollution	-		Population with primary reliance on polluting fuels and echnologies (%)	2016	-	-	26	
LEC	CTED ADULT RISK FACTOR	R TRENDS							
	CURRENT TOBACCO SMOKING			OBESITY		RAISED BLOOD PRESSURE			
	NO DATA AVAILABI	FI % of the population	40 30 20 10	0	30 at the population 20 2000	2005 2010	2015 2020	······································	
A1	ONAL EVETTING DESPONS		es nales	Past trends Projected linear trends Global targets0					
ATT(ONAL SYSTEMS RESPONS)-	D	Proportion of population at high risk for CVD or with					
			е	roportion of population at high risk for CVD or with xisting CVD (%) Proportion of high risk persons receiving any drug therapy	-				
)	Drug therapy to prevent heart attacks and	✓	а	nd counselling to prevent heart attacks and strokes (%)	-				
	strokes		C	Proportion of primary health care centres reported as iffering CVD risk stratification	2017		25% to 50%		
				Reported having CVD guidelines that are utilized in at east 50% of health facilities	2017		Yes		
1	Essential NCD medicines and basic technologies	X	н	lumber of essential NCD medicines reported as generally available"	2017		9 out of 10		
	to treat major NCDs			lumber of essential NCD technologies reported as generally available"	2017		6 out of 6		