


URUGUAY




50 800 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"							
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
4OR	TALITY						
	Premature mortality from NCDs	✓	Total NCD deaths	2016	13 600	14 200	27 800
			Risk of premature death between 30-70 years (%)	2016	21	13	17
Ţ.	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	18
RISK	FACTORS						
	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	18	4	11
K	Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016	19	28	24
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	7	7	7
⊗	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	19	13	16
	Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	27	22	24
À	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	10	12	11
i i	Obesity	✓	Obesity, adults aged 18+ (%)	2016	25	32	29
			Obesity, adolescents aged 10-19 (%)	2016	14	10	12
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)°	2016	-	-	-
h	Household air pollution	-	Population with primary reliance on polluting fuels and technologies $(\!\%\!)$	2016	-	-	<5









NATIONAL SYSTEMS RESPONSE



to treat major NCDs



х

