


## **SWITZERLAND**




## > 31% 24% Cardiovascular Other NCDs diseases **4**% 27% Communicable, Cancers maternal, perinatal NCDs are 5% and nutritional estimated to account for 90% conditions Chronic of all deaths. respiratory 6% diseases Injuries 2% Diahetes

	Males Females									
34 700 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"										
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL			
MOR'	TALITY									
	Premature mortality from NCDs	X	Total NCD deaths	2016	29 100	31 600	60 700			
			Risk of premature death between 30-70 years (%)	2016	11	7	9			
E	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	17			
RISK FACTORS										
	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	18	5	11			
K	Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016	23	28	26			
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	10	9	9			
<b>©</b>	Tobacco use	X	Current tobacco smoking, adults aged 15+ (%)	2016	27	20	24			
<b>9</b> 8	Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	27	20	24			
i	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	7	4	6			
	Obesity	x	Obesity, adults aged 18+ (%)	2016	24	19	21			
<b>~</b> /\			Obesity, adolescents aged 10-19 (%)	2016	6	4	5			
<b>%</b>	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	1			
à	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	<5			
SELECTED ADULT RISK FACTOR TRENDS										
<b>CURRENT TOBACCO SMOKING</b>			OBESITY	F	RAISED BLOOD PRESSURE					
40				40						







NATIONAL SYSTEMS RESPONSE									
			Proportion of population at high risk for CVD or with existing CVD (%)	-					
	Drug therapy to prevent heart attacks and strokes	V	Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%)	-					
		X	Proportion of primary health care centres reported as offering CVD risk stratification	2017	More than 50%				
			Reported having CVD guidelines that are utilized in at least 50% of health facilities	2017	Yes				
	Essential NCD medicines and basic technologies to treat major NCDs	X	Number of essential NCD medicines reported as "generally available"	2017	10 out of 10				
			Number of essential NCD technologies reported as "generally available"	2017	6 out of 6				