

2016 TOTAL POPULATION: 261 100 000 **2016 TOTAL DEATHS:** 1 863 000


50 Probability (%) of premature death 40 30 ·····• 20 10 0 2000 2005 2010 2015 2020 2025 Past trends Projected linear trends Global targets Males Females

35% 15% Cardiovascular Other NCDs diseases 21% 12% Communicable, Cancers NCDs are maternal, perinatal 6% and nutritional estimated to account for 73% conditions Chronic of all deaths. respiratory 6% diseases Injuries 6% Diabetes

	remates							
193 900 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"								
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL	
MORTALITY*								
	Premature mortality	/	Total NCD deaths	2016	722 400	642 300	1 365 000	
	from NCDs	V	Risk of premature death between 30-70 years (%)	2016	30	23	26	
P.	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	3	
RISK FACTORS								
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	1	0	1	
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	22	21	22	
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	9	8	9	
⊗	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	76	3	39	
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	23	21	22	
i.	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	7	7	7	
Ä	Obesity	✓	Obesity, adults aged 18+ (%)	2016	5	9	7	
			Obesity, adolescents aged 10-19 (%)	2016	6	4	5	
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2	
4	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	42	
SELECTED ADULT RISK FACTOR TRENDS								
CURRENT TORACCO SMOVING			ORESITY	DAISED BLOOD DDESSUDE			DE	











Proportion of population at high risk for CVD or with existing CVD (%)	
Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%)	
Proportion of primary health care centres reported as offering CVD risk stratification	

Reported having CVD guidelines that are utilized in at least 50% of health facilities Number of essential NCD medicines reported as "generally available" Number of essential NCD technologies reported as

2017 Less than 25% 2017 No 2017 10 out of 10 2017 5 out of 6