

**2016 TOTAL POPULATION:** 8 735 000 **2016 TOTAL DEATHS:** 46 000

## 42% 12% 50 Cardiovascular Other NCDs Probability (%) of premature death 40 diseases > 23% 10% 30 Communicable, Cancers 20 NCDs are maternal, perinatal 4% and nutritional estimated to 10 account for 69% conditions Chronic of all deaths. respiratory 0 8% diseases 2000 2005 2010 2015 2020 2025 Injuries 2% Past trends Projected linear trends Global targets Males ....0 Diabetes .....ō Females

9 300 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"							
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY*							
	Premature mortality	<b>√</b>	Total NCD deaths	2016	17 000	14 900	31 900
	from NCDs	v	Risk of premature death between 30-70 years (%)	2016	29	22	25
<u>F</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	2
RISK	FACTORS						
	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	6	1	3
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	19	37	28
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	14	13	14
$\odot$	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016			
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	21	20	21
À	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	8	8	8
À	Obesity	✓	Obesity, adults aged 18+ (%)	2016	10	15	13
			Obesity, adolescents aged 10-19 (%)	2016	3	2	2
<b>%</b>	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	4
h	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	20
SELEC	TED ADULT RISK FACTO	OR TRENDS					








Males Females **OBESITY** 








## NATIONAL SYSTEMS RESPONS



**Essential NCD medicines** and basic technologies to treat major NCDs



offering CVD risk stratification

Number of essential NCD medicines reported as "generally available" Number of essential NCD technologies reported as

and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as Reported having CVD guidelines that are utilized in at least 50% of health facilities

% of the population

54 Less than 25% Yes 9 out of 10

4 out of 6

10

2017

<sup>=</sup> no data available