
NCDs are

estimated to

account for 72%

of all deaths.

ECUADOR



	Males Females		••••					
		113 800 LIVES CAN	BE S	AVED BY 2025 BY IMPLEMENTING ALL OF THE V	VHO "BEST BUY	S"		
ORT	TALITY TALITY	NATIONAL TARGET S	SET		DATA YEAR	MALES	FEMALES	ТОТА
_	Premature mortality	V		Total NCD deaths	2016	31 500	29 300	60 700
	from NCDs	X		Risk of premature death between 30-70 years (%)	2016	14	12	13
Į	Suicide mortality	-		Suicide mortality rate (per 100 000 population)	2016	-	-	7
SK	FACTORS							
	Harmful use of alcohol	X		Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	7	1	4
	Physical inactivity	X		Physical inactivity, adults aged 18+ (%)	2016	24	29	27
b .	Salt/Sodium intake	X		Mean population salt intake, adults aged 20+ (g/day)	2010	8	7	8
3	Tobacco use	X		Current tobacco smoking, adults aged 15+ (%)	2016	12	2	7
3	Raised blood pressure	X		Raised blood pressure, adults aged 18+ (%)	2015	19	15	17
İ	Diabetes	X		Raised blood glucose, adults aged 18+ (%)	2014	7	8	7
ń	Obesity	X		Obesity, adults aged 18+ (%)	2016	14	24	19
A)	Obesity	A		Obesity, adolescents aged 10-19 (%)	2016	9	7	8
>	Ambient air pollution	-		Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	1
	Household air pollution	-		Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	<5
	CURRENT TOBACCO SMOKING			OBESITY	ı	RAISED BLOOD PRESSURE		
40			ion	40	40			
30			of the population	30	of the population			
20			bob	20	20			
		******	the		the		***************************************	O
10		0	% of	10	5 10 8			О
0		······································	•	0	0			
	2000 2005 2010 20	015 2020 2025		2000 2005 2010 2015 2020 2025		2005 2010	2015 2020	2025
			Male: Fema					
\TI	ONAL SYSTEMS RESPONS	SE						_
				Proportion of population at high risk for CVD or with existing CVD (%)	-			
7	Drug therapy to prevent	v		Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%)	-			
4	heart attacks and strokes	X		Proportion of primary health care centres reported as offering CVD risk stratification	2017		Less than 25%	
				Reported having CVD guidelines that are utilized in at	2247		NI=	
				least 50% of health facilities	2017		No	
	Essential NCD medicines	Y			2017		9 out of 10	
]	Essential NCD medicines and basic technologies to treat major NCDs	х		least 50% of health facilities Number of essential NCD medicines reported as				