GERMANY

2016 TOTAL POPULATION: 81 915 000
2016 TOTAL DEATHS: 921 000

RISK OF PREMATURE DEATH DUE TO NCDS (%)

Males — Past trends — Projected linear trends — Global targets
Females

PROPORTIONAL MORTALITY

▶ **37%** Cardiovascular diseases
▶ **26%** Cancers
▶ **6%** Chronic respiratory diseases
▶ **3%** Diabetes
▶ **19%** Other NCDs
▶ **5%** Communicable, maternal, perinatal and nutritional conditions
▶ **4%** Injuries

NCDs are estimated to account for **91%** of all deaths.

---

340 800 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"

|	NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY						
Premature mortality from NCDs	X	Total NCD deaths	2016	406 900	432 700	839 500
		Risk of premature death between 30-70 years (%)	2016	15	9	12
Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	14
RISK FACTORS						
Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	21	6	13
Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016	44	48	46
Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	9	9	9
Tobacco use	X	Current tobacco smoking, adults aged 15+ (%)	2016	30	23	27
Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	31	25	28
Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	8	6	7
Obesity	✓	Obesity, adults aged 18+ (%)	2016	27	25	26
		Obesity, adolescents aged 10-19 (%)	2016	10	6	8
Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	1
Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	<5

SELECTED ADULT RISK FACTOR TRENDS

CURRENT TOBACCO SMOKING

OBESITY

RAISED BLOOD PRESSURE

Males — Past trends — Projected linear trends — Global targets
Females

NATIONAL SYSTEMS RESPONSE

	NATIONAL TARGET SET		DATA YEAR		
Drug therapy to prevent heart attacks and strokes	X	Proportion of population at high risk for CVD or with existing CVD (%)	-	...	
		Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%)	-	...	
		Proportion of primary health care centres reported as offering CVD risk stratification	2017	More than 50%	
		Reported having CVD guidelines that are utilized in at least 50% of health facilities	2017	Yes	
Essential NCD medicines and basic technologies to treat major NCDs	X	Number of essential NCD medicines reported as "generally available"	2017	10 out of 10	
		Number of essential NCD technologies reported as "generally available"	2017	6 out of 6	

... = no data available