


COSTA RICA




> 29% 20% Cardiovascular Other NCDs diseases 23% 6% Communicable, Cancers NCDs are maternal, perinatal 7% and nutritional estimated to account for 83% conditions Chronic of all deaths. respiratory 10% diseases Injuries 4% Diabetes

	Females		· O							
16 000 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"										
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL			
MORT	TALITY									
R	Premature mortality	/	Total NCD deaths	2016	11 000	9 500	20 500			
	from NCDs	*	Risk of premature death between 30-70 years (%)	2016	14	9	12			
P.	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	8			
RISK FACTORS										
	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	8	1	5			
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	38	55	46			
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	8	8	8			
8	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	17	6	12			
	Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	20	16	18			
À	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	8	9	9			
	a.		Obesity, adults aged 18+ (%)	2016	21	30	26			
	Obesity	√	Obesity, adolescents aged 10-19 (%)	2016	11	11	11			
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2			
4	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	7			
SELECTED ADULT RISK FACTOR TRENDS										
CURRENT TOBACCO SMOKING			OBESITY	RAISED BLOOD PRESSURE			RE			
40										







NATIONAL SYSTEMS RESPONSE									
(**)	Drug therapy to prevent heart attacks and strokes		Proportion of population at high risk for CVD or with existing CVD (%)	-					
			Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%)	-					
		V	Proportion of primary health care centres reported as offering CVD risk stratification	2017	More than 50%				
			Reported having CVD guidelines that are utilized in at least 50% of health facilities	2017	Yes				
	Essential NCD medicines and basic technologies to treat major NCDs	✓	Number of essential NCD medicines reported as "generally available"	2017	10 out of 10				
			Number of essential NCD technologies reported as "generally available"	2017	6 out of 6				