

ANDORRA


NO DATA AVAILABLE

NO DATA AVAILABLE


		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY [†]							
	Premature mortality from NCDs	X	Total NCD deaths	2016			
			Risk of premature death between 30-70 years (%)	2016			
E.	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	
RISK FACTORS							
	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	18	5	11
K	Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016	41	44	43
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	10	9	10
©	Tobacco use	X	Current tobacco smoking, adults aged 15+ (%)	2016	36	24	30
	Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	28	21	25
į.	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	10	8	9
Å	Obesity	x	Obesity, adults aged 18+ (%)	2016	28	28	28
			Obesity, adolescents aged 10-19 (%)	2016	13	9	11
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)°	2016	-	-	-
各	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	<5
SELEC	TED ADULT RISK FACTO	OR TRENDS					








OBESITY





NATIONAL SYSTEMS RESPONSE



Drug therapy to prevent heart attacks and strokes

х

Proportion of population at high risk for CVD or with existing CVD (%)

Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as

Reported having CVD guidelines that are utilized in at least 50% of health facilities

offering CVD risk stratification

Number of essential NCD medicines reported as Number of essential NCD technologies reported as "generally available"

2017

2017

Don't know 2017 No 10 out of 10 2017

6 out of 6

Essential NCD medicines and basic technologies to treat major NCDs