DOMINICA

NO DATA AVAILABLE

NO DATA AVAILABLE


		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY [†]							
Ä	Premature mortality	✓	Total NCD deaths	2016	***	***	***
	from NCDs	v	Risk of premature death between 30-70 years (%)	2016			
<u>F</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	
RISK FACTORS							
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	14	3	8
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	16	34	25
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	7	7	7
\odot	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016			
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	26	21	23
i.	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	9	14	11
i i	Obesity	✓	Obesity, adults aged 18+ (%)	2016	20	36	28
			Obesity, adolescents aged 10-19 (%)	2016	13	14	13
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2
h	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	9

SELECTED ADULT RISK FACTOR TRENDS


CURRENT TOBACCO SMOKING

NO DATA AVAILABLE





RAISED BLOOD PRESSURE



NATIONAL SYSTEMS RESPONSE



heart attacks and strokes





Proportion of population at high risk for CVD or with existing CVD (%)

Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as offering CVD risk stratification

Reported having CVD guidelines that are utilized in at least 50% of health facilities Number of essential NCD medicines reported as Number of essential NCD technologies reported as

"generally available"

2017 2017

None Yes

9 out of 10 2017 2017 6 out of 6