
GUATEMALA



> 20% 17% Cardiovascular Other NCDs diseases 10% > 25% Communicable, Cancers NCDs are maternal, perinatal 3% and nutritional estimated to account for **59%** conditions Chronic of all deaths. respiratory 16% diseases Injuries 8%

MORT/			E SAVED BY 2025 E					
MORT				BY IMPLEMENTING ALL OF THE W	HO "BEST BUYS"	"		
MORIA	ALITV	NATIONAL TARGET S	ET		DATA YEAR	MALES	FEMALES	TOTAL
	Premature mortality from NCDs	✓	Total NCD deat	hs ure death between 30-70 years (%)	2016 2016	23 900 15	23 500 15	47 500 15
F	Suicide mortality	-	Suicide mortali	ty rate (per 100 000 population)	2016	-	-	3
RISK F	ACTORS							
	Harmful use of alcohol	✓		Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)		4	1	2
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)		2016	35	36	35
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)		2010	8	7	7
8	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)		2016			
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)		2015	18	16	17
À	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)		2014	7	8	8
	Obesity	✓	Obesity, adults	Obesity, adults aged 18+ (%)		13	24	19
	Obesity	V	Obesity, adoles	cents aged 10-19 (%)	2016	8	9	8
%	Ambient air pollution	-		Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)		-	-	2
4	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)		2016	-	-	55
SELECT	TED ADULT RISK FACTO	R TRENDS						
CURRENT TOBACCO SMOKING				OBESITY		RAISED BLOOD PRESSURE		RE
% of the population		30	0	% of the population 20 20 10 10			8	

