


NORWAY




4 000 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"							
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY							
	Premature mortality from NCDs	✓	Total NCD deaths	2016	17 300	18 300	35 600
			Risk of premature death between 30-70 years (%)	2016	11	8	9
<u>F</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	12
RISK FACTORS							
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	12	3	7
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	31	36	34
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	10	9	10
8	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	21	19	20
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	29	21	25
-i	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	8	6	7
i	Obesity	✓	Obesity, adults aged 18+ (%)	2016	25	25	25
			Obesity, adolescents aged 10-19 (%)	2016	9	7	8
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)°	2016	-	-	-
h	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	<5









NATIONAL SYSTEMS RESPONSI





х



Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as offering CVD risk stratification

Reported having CVD guidelines that are utilized in at least 50% of health facilities

Number of essential NCD medicines reported as "generally available"

Number of essential NCD technologies reported as "generally available"







^{... =} no data available onot exceeding