



NIGERIA



	134 500 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"						
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY*							
	Premature mortality	/	Total NCD deaths	2016	293 700	323 600	617 300
	from NCDs	•	Risk of premature death between 30-70 years (%)	2016	21	24	22
<u>r</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	10
RISK	FACTORS						
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	22	5	13
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	22	27	25
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	8	7	7
\odot	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	11	1	6
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	18	19	18
i.	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	4	4	4
À	Obesity	✓	Obesity, adults aged 18+ (%)	2016	4	12	8
			Obesity, adolescents aged 10-19 (%)	2016	1	2	2
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	5
h	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	>95
SELEC	TED ADULT RISK FACTO	OR TRENDS					













and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as offering CVD risk stratification

Reported having CVD guidelines that are utilized in at least 50% of health facilities Number of essential NCD medicines reported as

"generally available" Number of essential NCD technologies reported as

