PAKISTAN



Current tobacco smoking, adults aged 15+ (%)

Exceedance of WHO guidelines level for annual PM2.5

Population with primary reliance on polluting fuels and

OBESITY

Raised blood pressure, adults aged 18+ (%)

Raised blood glucose, adults aged 18+ (%)

Obesity, adolescents aged 10-19 (%)

concentration (by a multiple of)

technologies (%)

Obesity, adults aged 18+ (%)

SELECTED ADULT RISK FACTOR TRENDS

X

X

8

48

Tobacco use


Diabetes


Obesity


Raised blood pressure

Ambient air pollution


Household air pollution











NATIONAL SYSTEMS RESPO



Essential NCD medicines and basic technologies to treat major NCDs

Proportion of population at high risk for CVD or with

Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as

offering CVD risk stratification Reported having CVD guidelines that are utilized in at least 50% of health facilities

Number of essential NCD medicines reported as generally available Number of essential NCD technologies reported as 2017

2016

2015

2014

2016

2016

2016

2016

of the population

35

27

10

6

3

3

24

10

10

2

19

25

10

8

57

None Yes

2017 2017 4 out of 10 2017 4 out of 6

â

⁼ no data available