

## ALBANIA




	Premature mortality from NCDs	✓	Total NCD deaths	2016	13 300	10 900	24 300
		V	Risk of premature death between 30-70 years (%)	2016	21	13	17
E	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	6
RISK FACTORS							
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	13	3	7
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	***		
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	10	9	9
<b>(S)</b>	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	51	8	29
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	35	29	32
-A	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	8	8	8
À	Obesity	,	Obesity, adults aged 18+ (%)	2016	22	23	22
		<b>√</b>	Obesity, adolescents aged 10-19 (%)	2016	8	5	6
<b>%</b>	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2
各	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	23

## SELECTED ADULT RISK FACTOR TRENDS









**RAISED BLOOD PRESSURE** 















5 out of 6

â

2017

<sup>=</sup> no data available