


HONDURAS


2016 TOTAL POPULATION: 9 113 000 **2016 TOTAL DEATHS:** 39 000



3 500 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"							
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORT	ALITY*						
\square	Premature mortality from NCDs	✓	Total NCD deaths	2016	13 000	13 200	26 200
	TIONI NODS		Risk of premature death between 30-70 years (%)	2016	16	12	14
	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	3
RISKI	ACTORS						
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	7	1	4
K	Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016			
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	8	7	7
\odot	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016		2	
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	19	16	17
À	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	7	8	7
À	Obesity	✓	Obesity, adults aged 18+ (%)	2016	14	25	19
			Obesity, adolescents aged 10-19 (%)	2016	8	9	8
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2
备	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	47
SELECTED ADULT RISK FACTOR TRENDS							








NATIONAL SYSTEMS RESPONSI










Number of essential NCD medicines reported as "generally available"

Number of essential NCD technologies reported as



4 out of 6

2017

% of the population

^{... =} no data available