

PARAGUAY


2016 TOTAL POPULATION: 6 725 000 **2016 TOTAL DEATHS:** 36 000




		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY*							
	Premature mortality	✓	Total NCD deaths	2016	14 000	13 100	27 000
	from NCDs	V	Risk of premature death between 30-70 years (%)	2016	19	16	18
E	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	9
RISK FACTORS							
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	12	2	7
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	37	36	36
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	11	10	11
8	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	21	5	13
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	25	18	22
į.	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	7	7	7
À	Obesity	/	Obesity, adults aged 18+ (%)	2016	16	22	19
			Obesity, adolescents aged 10-19 (%)	2016	10	8	9
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	1
b	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	34









NATIONAL SYSTEMS RESPO





Proportion of population at high risk for CVD or with existing CVD (%)

....00

Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as offering CVD risk stratification

Reported having CVD guidelines that are utilized in at least 50% of health facilities Number of essential NCD medicines reported as

"generally available" Number of essential NCD technologies reported as



Less than 25%

2017 Yes

2017 8 out of 10 2017 3 out of 6

â

⁼ no data available