

The Fluid Dynamics of the Casimir Vacuum: Planck's Viscosity, Wave Expansion, and the Geon Collapse Threshold

Daniel Gezin
Independent Researcher
[ORCID: 0009-0003-7309-7050](https://orcid.org/0009-0003-7309-7050)

February 18, 2026

Abstract

Standard quantum mechanics treats the vacuum as a passive void and Planck's constant (h) as an abstract quantization rule. This paper reconceptualizes the Casimir vacuum as a hyper-elastic, zero-point tensor fluid, deriving Planck's constant mechanically as the fundamental kinematic action viscosity of empty space. By applying classical fluid principles to electromagnetic propagation, we derive a novel metric—the Topological Reynolds Number of the Vacuum (Re_{vac}). This metric strictly explains why low-energy light disperses spherically (the inverse-square law), why high-energy gamma radiation focuses linearly like a projectile, and calculates the exact 1.022 MeV threshold where wave expansion is overpowered by vacuum crush. Furthermore, we define the asymmetrical topological shear (the "Nuclear Anvil") required to torque these extreme linear waves, strictly deriving the experimental mechanics of Pair Production.

1 The Hyper-Elastic Casimir Vacuum

The empirical existence of the Casimir effect [1] proves that the vacuum is not a void; it is an active, pressurized field of zero-point fluctuation. To support continuous wave propagation, this field must possess the mechanical characteristics of a hyper-elastic tensor fluid.

When a free electromagnetic wave propagates, it does not move through "nothing." It must physically displace the structural tension of this vacuum fluid. Like any physical medium, the vacuum naturally resists displacement. The observable energy of an electromagnetic wave is simply the kinetic effort required to continuously deform this elastic medium at velocity c .

2 The Mechanical Origin of Planck's Constant (h)

In 1901, Max Planck introduced h strictly as a mathematical proportionality constant ($E = hf$) to quantize radiation [2]. However, dimensionally, Planck's constant possesses the units of Action (Joules $\times$ seconds), which strictly equates to:

$$h = 6.62607 \times 10^{-34} \text{ kg} \cdot \text{m}^2/\text{s} \quad (1)$$

In classical fluid dynamics, kinematic viscosity (ν) dictates the energy required to shear a fluid over time. By incorporating mass-energy equivalence, h perfectly aligns with the dimensions of a mass-weighted viscous medium. In this framework, h is strictly redefined as the **Kinematic Action Viscosity of the Casimir Vacuum**.

The foundational equation $E = hf$ is mechanically decoded as a fluid-dynamic power equation. A propagating photon must violently push against the baseline viscosity of space. The

total energy (E) required to exist is directly proportional to how many times per second (f) the wave-front must cycle against the fundamental sheer resistance (h) of the vacuum medium.

3 Wave Expansion and the Inverse Square Law

Why does light from a star expand in a spherical shell, classically obeying the inverse square law ($I \propto 1/r^2$)?

In our fluidic vacuum, the transverse electric ($\vec{E}$) vectors of a propagating wave represent regions of intense local elastic stress. Like a highly compressed gas expanding into a lower-pressure atmosphere, the localized electromagnetic stress naturally seeks to minimize its energy state. The electric vectors repel one another laterally along the wave-front, causing the spatial boundary of the wave to continuously dilate.

The macroscopic expansion pressure (P_{exp}) of the wave-front pushing outward spherically is proportional to its kinetic energy distributed over the expanding surface area:

$$P_{exp} \propto \frac{E}{4\pi r^2 c \tau} \quad (2)$$

This continuous geometric dilation smoothly distributes the energy over an ever-widening area, mechanically deriving the classical phenomenon of spherical diffraction.

4 The "Bullet in Water" Effect: High-Energy Focusing

Experimental physics demonstrates that high-energy photons (X-rays, gamma rays) do not diffract spherically like radio waves; they travel in tight, highly focused linear beams with minimal lateral spread.

We establish this via the **"Bullet in Water" Analogy**. When a low-velocity projectile enters water, the fluid has time to react, creating broad, rippling wakes (spherical dispersion). When an extreme-velocity projectile enters the water, it punches through the medium faster than the fluid's bulk modulus can respond laterally, creating a narrow, highly focused cavitation tunnel.

4.1 The Topological Reynolds Number (Re_{vac})

To calculate this focusing effect, we introduce a novel parameter balancing the wave's inertial penetration against the vacuum's lateral crushing force: the **Topological Reynolds Number of the Vacuum** (Re_{vac}).

$$Re_{vac} = \frac{\text{Wave Inertial Penetration}}{\text{Vacuum Viscous Relaxation}} = \frac{E_\gamma}{h \cdot f_{vac}} \quad (3)$$

Where f_{vac} represents the baseline relaxation frequency of the local zero-point field.

- **Low Energy (Radio/Visible, Low Re_{vac}):** The wave displaces the vacuum slowly. P_{exp} dominates. The stress distributes laterally, creating an expanding spherical "gas."
- **High Energy (Gamma, High Re_{vac}):** The wave cycles so violently that it punches a strict linear tunnel through the vacuum. The inward lateral crush of the surrounding Casimir field occurs too slowly to dissipate the energy, forcing the wave into an ultra-tight, focused cylinder.




Figure 1: Fluid Dynamics of Light: Low energy waves expand laterally like a gas. High energy waves possess high Re_{vac} , piercing the vacuum linearly while being strictly constrained by lateral Casimir pressure.

5 The Geon Threshold: Mathematical Overpowering

As the wavelength (λ) shrinks into the picometer scale, the spatial volume of the wave packet decreases cubically. The external inward Casimir vacuum pressure (P_{vac}) attempting to crush this highly concentrated displacement scales inversely with the cube of the wavelength:

$$P_{vac} \propto \frac{hc}{2\lambda^3} \quad (4)$$

Because λ^3 shrinks geometrically faster than the lateral r^2 expansion, there exists a strict, calculable mathematical singularity where the inward crushing force of the vacuum absolutely overpowers the internal repulsive expansion of the wave:

$$P_{vac} > P_{exp} \quad (5)$$

5.1 Deriving the Critical Limit (1.022 MeV and 1.213 pm)

To physically collapse a linear wave into stable, circulating leptonic loops (an electron and positron), the wave must possess enough internal energy to satisfy the topological rest-mass boundary conditions of both resulting geometries ($E \geq 2m_e c^2$).

We calculate the exact critical energy threshold (E_{crit}):

$$E_{crit} = 2(0.510998 \text{ MeV}) = 1.02199 \text{ MeV} \approx 1.637 \times 10^{-13} \text{ J} \quad (6)$$

At this specific energy, the wavelength reaches its critical compression limit (λ_{crit}):

$$\lambda_{crit} = \frac{hc}{E_{crit}} = \frac{1.986 \times 10^{-25} \text{ J} \cdot \text{m}}{1.637 \times 10^{-13} \text{ J}} \approx 1.213 \times 10^{-12} \text{ m} \quad (7)$$

At exactly $\lambda \leq 1.213 \text{ pm}$, the volumetric Casimir crush vastly exceeds the linear wave's ability to expand. However, if the wave collapses, why do higher-energy gamma rays (e.g., 10 TeV from astrophysical sources) survive in deep space?

6 Catalyzed Topological Collapse: The Nuclear Anvil

In a pure, homogeneous vacuum, the lateral Casimir pressure crushing inward on the high-energy wave is perfectly symmetric. Because the wave possesses immense forward linear momentum, it acts as a balanced hypersonic projectile. The symmetric inward crush cannot overcome the extreme forward inertia without a structural perturbation; therefore, the wave penetrates the vacuum indefinitely.

To buckle the linear wave, there must be an **asymmetrical topological shear**.

6.1 Nuclear Pair Production

This requirement is historically validated by **Nuclear Pair Production**. In 1933, Blackett and Occhialini [?] proved that a gamma-ray ≥ 1.022 MeV must pass near a heavy atomic nucleus to transform into an electron and positron.

In the Geon framework, the heavy nucleus (a massive multi-trefoil topology) projects an immense, highly concentrated spatial gradient of both electrostatic stress and Casimir shadowing. When the high-energy gamma "bullet" grazes this steep gradient, the side of the wave closest to the nucleus experiences massive drag compared to the far side.

This extreme asymmetrical shear acts as a topological anvil. The forward linear momentum of the wave is abruptly arrested and torqued. Because the intense Casimir pressure ($P_{vac} > P_{exp}$) prevents the torqued wave from dispersing laterally, the wave violently buckles. To conserve angular momentum, it shears into two counter-rotating vortices.




Figure 2: Catalyzed Topological Collapse: The extreme forward inertia of a high-energy gamma ray prevents spontaneous collapse in a pure vacuum. When it grazes a heavy nucleus, the asymmetrical spatial gradient applies severe shear. The wave buckles under the Casimir crush, locking its phase into two counter-rotating, circularly polarized loops (Topological Geons).

To survive annihilation, the sheared wave ends lock their phases, curling into two mirrored 4π loops. The linear kinetic energy is permanently converted into the continuous circularly polarized rotation of the Topological Geon, strictly deriving the genesis of structural mass ($m = E/c^2$).

6.2 The Breit-Wheeler Process: Self-Shearing in a Vacuum

In 1934, Breit and Wheeler [?] theorized that two high-energy photons colliding in an empty vacuum would provide enough concentrated energy density to trigger this collapse without a nucleus. This was definitively proven by the STAR collaboration in 2021 [?].

Within this framework, the Breit-Wheeler process is perfectly consistent. When two high-energy beams perfectly cross in a pure vacuum, their intersecting momentum vectors act as the extreme topological shear gradient *for each other*. They serve as mutual topological anvils, providing the exact asymmetrical torque required to buckle their linear tracks under the surrounding Casimir pressure, resulting in spontaneous topological pair production in empty space.

7 Conclusion

By redefining the quantum vacuum as an active, hyper-elastic fluid, we achieve total mechanical unity for electromagnetic propagation. Planck's constant (h) is explicitly defined as the kinematic action viscosity of empty space. The inverse-square law is derived as the lateral relaxation of this fluid, while high-energy focusing is governed by the Topological Reynolds Number (Re_{vac}). Finally, the explicit calculation of the 1.213 pm boundary perfectly derives the 1.022 MeV threshold where vacuum pressure overpowers wave expansion, requiring only topological shear to catalyze the genesis of fundamental particles.

References

- [1] Hendrik B. G. Casimir. On the attraction between two perfectly conducting plates. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen*, 51:793–795, 1948.
- [2] Max Planck. Ueber das gesetz der energieverteilung im normalspectrum. *Annalen der Physik*, 309(3):553–563, 1901.

Original Preprint – D. Gezin