

# 01160611 — SELECT TOPICS IN LIVING SYSTEMS

## LECTURES

DR. RAM ADAR\*

*Daniel Haim Breger<sup>†</sup>*

*Physics 300, Technion*

*29<sup>th</sup> October 2025 – 0<sup>th</sup> February 2026*

*These notes were typed during the 2026 winter semester in the Technion.  
The lectures were given in Hebrew and were live-translated by me to English.  
The document is provided as is and likely contains many errors.*

---

## CONTENTS

1	Phase transitions in equilibrium	2
1.1	Motivation . . . . .	2
1.2	Physical description . . . . .	2
1.3	Landau Theory of Phase Transitions . . . . .	3
1.3.1	Where a comes from . . . . .	5
1.3.2	A dynamical description of the phase transition . . . . .	5
1.4	Spatial Dependence: The Ginsburg-Landau Theory . . . . .	6

---

\*radar@technion.ac.il (speaker)

<sup>†</sup>danielbreger@campus.technion.ac.il (scribe)

Lecture 1 (2 hours)  
29<sup>th</sup> October 2025

## 1 PHASE TRANSITIONS IN EQUILIBRIUM

### 1.1 Motivation

Phase is a distinct state of matter which can be described by an "order parameter". We'll focus on isotropic polar phase transition. In this case the order parameter is the polarization  $\vec{p} = \langle \hat{n} \rangle$  where  $\hat{n}$  is the unit vector of the particles' direction.

Phase transition: in certain areas of physics we see systems which aren't ordered at high temperatures but become ordered at low temperatures. Why? A competition between energy and entropy (Free energy being  $F = U - TS$ ). The temperature at which order emerges is called the critical temperature, marked  $T_C$  and the change in the system is called a phase transition.




Figure 1: Polar and Isotropic ordering of matter.

### 1.2 Physical description

We'll assume free energy of the following form:

$$F(T, N, \psi) = F_0(T, N) + F_\psi(T, N, \psi) \quad (1)$$

where  $\psi$  is the order parameter.  $\psi$  is not a conserved quantity but rather a degree of freedom of the system (in contrast to the number of particles, for example). Therefore under equilibrium we find

$$F_{eq}(T, N, \psi) = F_0(T, N) + F_\psi(T, N, \psi_{eq}) \quad (2)$$

and the equilibrium value is determined by

$$\left. \frac{\partial F_\psi}{\partial \psi} \right|_{\psi=\psi_{eq}} = 0 \quad (3)$$

and the following is true:

$$\psi_{eq}(T > T_C) = 0 \quad (4)$$

$$\psi_{eq}(T < T_C) \neq 0 \quad (5)$$

We'll focus on phase transition of 2nd order, for which the order parameter changes continuously, thus  $\psi_{eq}(T = T_C) = 0$ . Otherwise the phase transition is of first order.

### 1.3 Landau Theory of Phase Transitions

The theory was developed by Lev Landau, for which he earned the Nobel Prize in Physics in 1962.

Close to the critical temperature  $T_C$  the physics of phase transitions is universal. For a phase transition of 2nd order, we will write  $F_\psi$  in a phenomenological form as a Taylor series expansion:

$$F_\psi = \frac{a}{2}\psi^2 + \frac{b}{4}\psi^4 \quad (6)$$

The terms being of even powers because we don't care about directions. The quadratic term's coefficient is of the form

$$a = a_0 \cdot \frac{T - T_C}{T_C} \quad (7)$$

Where  $a_0 > 0$ . This quantifies the competition between energy (which wins at  $T < T_C$ ) and entropy (which wins at  $T > T_C$ ). The parameter  $b > 0$  ensures stability:  $\psi_{eq}$  receives finite values and does not approach infinity.

Because of  $a$ 's sign, we find  $\psi_{eq} \neq 0$  for  $T < T_C$ . Explicitly:

$$0 = \left. \frac{\partial F}{\partial \psi} \right|_{\psi=\psi_{eq}} = a\psi_{eq} + b\psi_{eq}^3 = b\psi_{eq} \left( \psi_{eq}^2 + \frac{a}{b} \right) \quad (8)$$




Figure 2: The graphs for the functions  $2x^2$  (blue),  $0.1x^4$  (green) and  $0.1x^4 - 2x^2$  (red).

therefore

$$\psi_{eq} = 0, \pm \sqrt{\frac{a_o}{b} \cdot \frac{T - T_C}{T_C}} \quad (9)$$




Figure 3: The possible solutions from equation 9.

There is no distinction between the positive and negative sign. Either can be chosen arbitrarily (spontaneous symmetry break). There will be a preference given an external field which will add an element of the form  $-\hbar\psi$  to the free energy.

### 1.3.1 Where $a$ comes from

We'll assume a simple model of polarisation with energy

$$U = - \sum_{i,j \text{ neighbors}} u n_{\alpha}^i n_{\alpha}^j, \quad u > 0 \quad (10)$$

The energy is lower when  $i,j$  point in the same direction. Using mean field approximation the internal energy is

$$U = \frac{u}{2} \sum_{i,j} \langle n_{\alpha}^i \rangle \cdot \langle n_{\alpha}^j \rangle = -\frac{u}{2} N Z p_{\alpha}^2 \quad (11)$$

where  $p_{\alpha} = \langle n_{\alpha} \rangle$ ,  $N$  is the number of particles, and  $Z$  the number of close neighbors. As for the entropy, we can find from a Taylor expansion

$$S = S(p=0) - \alpha N p^2 \quad (12)$$

and summing it up we find

$$F = U - TS = N \left( \alpha T - \frac{u}{2} Z \right) p^2 = a_0 \cdot \frac{T - T_C}{T_C} p^2 \quad (13)$$

where

$$T_C = \frac{uZ}{2\alpha} \quad a_0 = \alpha N T_C \quad (14)$$

If instead of the vector  $n_{\alpha}$  there is spin ( $\pm$ ) then we find a similar result in the mean field framework. This model is called the Ising model, one of the most basic and useful models in statistical mechanics.

### 1.3.2 A dynamical description of the phase transition

We described the phase transition from the free energy but we can also describe it from dynamics. For a non-conserved order parameter we can write the dynamics as follows:

$$\dot{\psi} = -\Gamma_{\psi} \frac{\partial F}{\partial \psi}, \quad \Gamma_{\psi} > 0 \quad (15)$$

the idea being that the dynamics guarantee that the free energy only gets smaller (the total entropy grows). We can see this using the chain rule:

$$\dot{F} = \frac{\partial F}{\partial \psi} \dot{\psi} = -\Gamma_{\psi} \left( \frac{\partial F}{\partial \psi} \right)^2 < 0 \quad (16)$$

this dynamics is called Model A (and often appears with a "noise" parameter). We'll insert F and find:

$$\dot{p} = -\Gamma_p(ap + bp^3) \quad (17)$$

the fixed points are  $\frac{\partial F}{\partial p} = 0$ , the extremums of the free energy. The stability of the solution is determined in phase space.




Figure 4: The graphs of  $f(x) = x - 2x^3$  (blue) and  $g(x) = -x - 2x^3$  (red). The positive and negative points marked on the  $x$  axis are stable points of equilibrium for  $f(x)$ . The point marked at  $(0,0)$  is a stable point for  $g(x)$  but an unstable one for  $f(x)$ .

in the language of dynamical systems, a phase transition is called a bifurcation. This approach is useful for out of equilibrium systems, for which we can't write down a free energy. Later on in the course, we'll come across dynamical equations of a similar form.

In general dynamical systems, when a function F whose derivative is strictly non-positive with a finite minimum exists (like free energy), then it's called a Lyapunov function.

#### 1.4 Spatial Dependence: The Ginsburg-Landau Theory

We saw that the direction of  $\vec{p}$  is arbitrary, in practice different areas of the system will initially develop order in different directions. For simplicity, we'll assume  $\vec{p} = p\hat{y}$  where  $p$  can be either positive or negative. We'll look at the case where  $\vec{p}$  points down at  $x \rightarrow -\infty$  and up at  $x \rightarrow \infty$ .




Figure 5: A smooth phase transition with Domain Wall  $\xi$ .




Figure 6: A smooth phase transition with Domain Wall  $\xi$ .

We'll find a finite range around 0 of width  $\xi$  in which  $|p| \neq p_{eq}$ . This region is called the Domain Wall.

To quantify the cost to change  $p$  in space we'll write the density of free energy per length unit:

$$F[p] = \int dx f = \int dx \left( \frac{a}{2} p^2 + \frac{b}{4} p^4 + \frac{k}{2} p'^2 \right) \quad (18)$$

where  $k > 0$ . The last element quantifies the cost to change  $p$ . In higher dimensions we'll have  $(\vec{\nabla} p)^2$ . Why is it like this? The cost comes from the gradient, and given there is no preferred direction, a positive and negative change should cost the same.

\* \* \*