

Quantum 2 HW3

Daniel Haim Breger, 316136944

Technion

(Date: December 5, 2025)

1 Simple Model for Quantization of the Electromagnetic Field

1. Use Maxwell's equations to write time development equations for $\mathcal{E}$ and $\mathcal{B}$.
2. Express $U(t)$ using $V, \mathcal{E}(t), \mathcal{B}(t), \mu_0, \varepsilon_0$. Use

$$\int_V d\mathbf{r} \sin^2 kz = \int_V d\mathbf{r} \cos^2 kz = \frac{V}{2}$$

3. We'll define $\omega = ck$ and

$$\chi(t) = \sqrt{\frac{\varepsilon_0 V}{2\hbar\omega}} \mathcal{E}(t) \qquad \Pi(t) = \sqrt{\frac{V}{2\mu_0\hbar\omega}} \mathcal{B}(t)$$

show that the equations of motion for $\chi(t)$ and $\Pi(t)$ are identical in their form to those of X and P in subsection 1.

1.1 Time Development

The relevant Maxwell's equations for time development are

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \qquad \nabla \times \mathbf{B} = \mu_0 \left(\mathbf{J} + \varepsilon_0 \frac{\partial \mathbf{E}}{\partial t} \right) \quad (1)$$

and since no current or charges at all other than out waves were specified, we assume the current $\mathbf{J} = 0$. Plugging the given fields into the equations we find

$$k\mathcal{E} = \frac{\partial \mathcal{B}}{\partial t} \qquad k\mathcal{B} = -\mu_0\varepsilon_0 \frac{\partial \mathcal{E}}{\partial t} \quad (2)$$

we can take the derivative of the equations and plug them into the other ones to find 2nd order differential equations

$$\frac{\partial^2 \mathcal{E}}{\partial t^2} + \frac{k^2}{\mu_0\varepsilon_0} \mathcal{E} = 0 \qquad \frac{\partial^2 \mathcal{B}}{\partial t^2} + \frac{k^2}{\mu_0\varepsilon_0} \mathcal{B} = 0 \quad (3)$$

1.2 Energy in Time

$$U(t) = \int_V d\mathbf{r} \left(\frac{\varepsilon_0}{2} \mathbf{E}^2 + \frac{1}{2\mu_0} \mathbf{B}^2 \right) \quad (4)$$

$$= \int_V d\mathbf{r} \left(\frac{\varepsilon_0}{2} \mathcal{E}^2 \sin^2(kz) + \frac{1}{2\mu_0} \mathcal{B}^2 \cos^2(kz) \right) \quad (5)$$

$$= \frac{V}{2} \left(\frac{\varepsilon_0}{2} \mathcal{E}^2 + \frac{1}{2\mu_0} \mathcal{B}^2 \right) \quad (6)$$

1.3 Same Form

The equations of motion for $\mathcal{E}$ and $\mathcal{B}$ are 2nd order linear ODEs, which are invariant to multiplication by scalars. The transformation of parameters from $\mathcal{E}$ and $\mathcal{B}$ to χ and Π simply multiplies by a scalar, therefore the equations of motion will remain the same.

2 Quantization of the Field

1. Express $\hat{H}_F$ in terms of $\hat{a}$ and $\hat{a}^\dagger$ and the number operator $\hat{N} = \hat{a}^\dagger \hat{a}$.
2. Calculate the expectation values $\langle \hat{E}(\mathbf{r}) \rangle_n$, $\langle \hat{B}(\mathbf{r}) \rangle_n$, $\langle \hat{H}_F \rangle$ in state $|n\rangle$.

2.1 Field Hamiltonian

From the definitions for the photon creation and annihilation operators we can find expressions for $\hat{\chi}$ and $\hat{\Pi}$ using the operators:

$$\hat{\chi} = \frac{1}{\sqrt{2}} (\hat{a} + \hat{a}^\dagger) \quad \hat{\Pi} = \frac{1}{\sqrt{2}i} (\hat{a} - \hat{a}^\dagger) \quad (7)$$

therefore

$$\hat{H}_F = \frac{\hbar\omega}{2} (\hat{\chi}^2 + \hat{\Pi}^2) \quad (8)$$

$$= \frac{\hbar\omega}{4} ((\hat{a} + \hat{a}^\dagger)^2 - (\hat{a} - \hat{a}^\dagger)^2) \quad (9)$$

$$= \frac{\hbar\omega}{4} (\hat{a}^2 + \hat{a}\hat{a}^\dagger + \hat{a}^\dagger\hat{a} + \hat{a}^{\dagger 2} - \hat{a}^2 + \hat{a}\hat{a}^\dagger + \hat{a}^\dagger\hat{a} - \hat{a}^{\dagger 2}) \quad (10)$$

$$= \frac{\hbar\omega}{2} (\hat{a}\hat{a}^\dagger + \hat{N}) \quad (11)$$

$$= \frac{\hbar\omega}{2} (\hat{a}^\dagger\hat{a} - \hat{a}^\dagger\hat{a} + \hat{a}\hat{a}^\dagger + \hat{N}) \quad (12)$$

$$= \frac{\hbar\omega}{2} (2\hat{N} + [\hat{a}, \hat{a}^\dagger]) \quad (13)$$

$$= \frac{\hbar\omega}{2} (2\hat{N} + 1) \quad (14)$$

$$= \hbar\omega \left(\hat{N} + \frac{1}{2} \right) \quad (15)$$

2.2 Expectation Values

First we notice

$$\langle \hat{E}(\mathbf{r}) \rangle_n = \langle n | \hat{E} | n \rangle \quad (16)$$

which if we look at the definition of the creation and annihilation operators which define $\hat{E}$, the result of the above equation is always zero, therefore

$$\langle \hat{E}(\mathbf{r}) \rangle_n = 0 \quad (17)$$

and the same holds for $\langle \hat{B}(\mathbf{r}) \rangle_n$. For $\hat{H}_F$:

$$\langle n | \hat{H}_F | n \rangle = \hbar\omega \langle n | \left(\hat{N} + \frac{1}{2} \right) | n \rangle = \hbar\omega \left(n + \frac{1}{2} \right) \quad (18)$$

3 Coherent States of the Electromagnetic Field

The coherent states of the Electromagnetic field are defined as

$$|\alpha\rangle = e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle, \quad \alpha \in \mathbb{C} \quad (19)$$

1. Show that the state $|\alpha\rangle$ is a normalized eigenstate of the annihilation operator a . Find the corresponding eigenvalue. Calculate the expectation value $\langle N \rangle_\alpha$ in this state.
2. Given $|\psi(t=0)\rangle = |\alpha\rangle$, show that

$$|\psi(t)\rangle = \exp\left(-i\frac{\omega t}{2}\right) |\alpha e^{-i\omega t}\rangle$$

3. Calculate the expectation values $\langle \mathbf{E}(\mathbf{r}) \rangle_t$ and $\langle \mathbf{B}(\mathbf{r}) \rangle_t$ in the state $|\psi(t)\rangle$. Assume α is real.
4. Show that $\langle \mathbf{E}(\mathbf{r}) \rangle_t$ and $\langle \mathbf{B}(\mathbf{r}) \rangle_t$ satisfy Maxwell's equations.
5. Calculate the classical energy of the field for $\mathbf{E}_{cl} = \langle \mathbf{E}(\mathbf{r}) \rangle_t$ and $\mathbf{B}_{cl} = \langle \mathbf{B}(\mathbf{r}) \rangle_t$ and compare with $\langle H_F \rangle_t$.
6. Assume $|\alpha| \gg 1$ and justify the statement that the states $|\alpha\rangle$ have a "classical" character.

3.1 Normalized Eigenstate

To show $|\alpha\rangle$ is an eigenstate of the annihilation operator we'll activate it on the state:

$$a |\alpha\rangle = e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} \cdot a |n\rangle \quad (20)$$

$$= e^{-\frac{|\alpha|^2}{2}} \sum_{n=1}^{\infty} \frac{\alpha^n}{\sqrt{n!}} \sqrt{n} |n-1\rangle \quad (21)$$

$$= e^{-\frac{|\alpha|^2}{2}} \sum_{n=1}^{\infty} \frac{\alpha^n}{\sqrt{(n-1)!}} |n-1\rangle \quad (22)$$

$$= e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha^{n+1}}{\sqrt{n!}} |n\rangle \quad (23)$$

$$= \alpha e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} \cdot a |n\rangle \quad (24)$$

$$= \alpha |\alpha\rangle \quad (25)$$

therefore $|\alpha\rangle$ is an eigenstate of the annihilation operator with eigenvalue α . In line 21 the sum begins from 1 because the ground state annihilates into a zero, so it can be removed from the sum. Next we check that

the state is normalized:

$$\langle \alpha | \alpha \rangle = e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} \cdot e^{-\frac{|\alpha^*|^2}{2}} \sum_{n=0}^{\infty} \frac{(\alpha^*)^n}{\sqrt{n!}} \langle n | n \rangle \quad (26)$$

$$= e^{-|\alpha|^2} \sum_{n=0}^{\infty} \frac{\alpha^{*n} \cdot \alpha^n}{n!} \quad (27)$$

$$= e^{-|\alpha|^2} \sum_{n=0}^{\infty} \frac{|\alpha|^{2n}}{n!} \quad (28)$$

$$= e^{-|\alpha|^2} \cdot e^{|\alpha|^2} \quad (29)$$

$$= 1 \quad (30)$$

noting that in the general case it should be expanded as $\langle n | m \rangle$ but since it obviously becomes δ_{nm} I just skipped there directly. Next the expectation value of N is:

$$\langle N \rangle_{\alpha} = \langle \alpha | N | \alpha \rangle = \langle \alpha | a^{\dagger} a | \alpha \rangle = \alpha^* \alpha \quad (31)$$

3.2 Showing That

We know that $|\alpha\rangle$ is an eigenstate of the annihilation operator with eigenvalue α . As we know from question 2, the hamiltonian (which is time independent) of the EM field can be expressed using the creation and annihilation operators. This means the state $|\alpha\rangle$ is also an eigenstate of the EM field, which means we can use standard time evolution of eigenstates inserting the energy from question 2:

$$|\psi(t)\rangle = e^{-i\omega t(N+\frac{1}{2})} |\alpha\rangle \quad (32)$$

$$= e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} e^{-i\omega t(n+\frac{1}{2})} \frac{\alpha^n}{\sqrt{n!}} |n\rangle \quad (33)$$

$$= e^{-\frac{|\alpha|^2}{2} - i\omega t} \sum_{n=0}^{\infty} \frac{(\alpha \cdot e^{-i\omega t})^n}{\sqrt{n!}} |n\rangle \quad (34)$$

$$= [\beta \equiv \alpha \cdot e^{-i\omega t}] = e^{-\frac{|\beta|^2}{2} - i\omega t} \sum_{n=0}^{\infty} \frac{\beta^n}{\sqrt{n!}} |n\rangle \quad (35)$$

so

$$|\psi(t)\rangle = e^{-\frac{i\omega t}{2}} |\beta\rangle = \exp\left(-i\frac{\omega t}{2}\right) |\alpha e^{-i\omega t}\rangle \quad (36)$$

3.3 Expectation Values

$$\langle \mathbf{E}(\mathbf{r}) \rangle_t = \langle \psi(t) | \mathbf{E} | \psi(t) \rangle \quad (37)$$

$$= \langle \alpha e^{i\omega t} | \mathbf{E} | \alpha e^{-i\omega t} \rangle \quad (38)$$

$$= \varepsilon_x \sqrt{\frac{\hbar\omega}{\varepsilon_0 V}} \sin(kz) \langle \alpha e^{i\omega t} | (\alpha^{\dagger} + \alpha) | \alpha e^{-i\omega t} \rangle \quad (39)$$

$$= \varepsilon_x \sqrt{\frac{\hbar\omega}{\varepsilon_0 V}} \sin(kz) (\alpha e^{-i\omega t} + \alpha e^{i\omega t}) \quad (40)$$

$$= 2\alpha\varepsilon_x \sqrt{\frac{\hbar\omega}{\varepsilon_0 V}} \sin(kz) \cos(\omega t) \quad (41)$$

and

$$\langle \mathbf{B}(\mathbf{r}) \rangle_t = \langle \alpha e^{i\omega t} | \mathbf{B} | \alpha e^{-i\omega t} \rangle \quad (42)$$

$$= i\varepsilon_y \sqrt{\frac{\mu_0 \hbar \omega}{V}} \cos(kz) \langle \alpha e^{i\omega t} | (a^\dagger - a) | \alpha e^{-i\omega t} \rangle \quad (43)$$

$$= i\varepsilon_y \sqrt{\frac{\mu_0 \hbar \omega}{V}} \cos(kz) (\alpha e^{i\omega t} - \alpha e^{-i\omega t}) \quad (44)$$

$$= -2\alpha\varepsilon_y \sqrt{\frac{\mu_0 \hbar \omega}{V}} \cos(kz) \sin(\omega t) \quad (45)$$

3.4 Satisfying Maxwell's Equations

For the first equation, on one hand

$$\nabla \times \langle E \rangle = \nabla \times \left(2\alpha\varepsilon_x \sqrt{\frac{\hbar\omega}{\varepsilon_0 V}} \sin(kz) \cos(\omega t) \right) \quad (46)$$

$$= 2\alpha k \varepsilon_y \sqrt{\frac{\hbar\omega}{\varepsilon_0 V}} \cos(kz) \cos(\omega t) \quad (47)$$

$$= [\omega = ck, c = \frac{1}{\sqrt{\mu_0 \varepsilon_0}}] = 2\alpha ck \sqrt{\frac{\mu_0 \hbar ck}{V}} \cos(kz) \cos(ckt) \varepsilon_y \quad (48)$$

on the other hand

$$\frac{\partial \langle B \rangle}{\partial t} = -2\alpha\omega\varepsilon_y \sqrt{\frac{\mu_0 \hbar \omega}{V}} \cos(kz) \sin \cos \omega t \quad (49)$$

$$= -2\alpha ck \sqrt{\frac{\mu_0 \hbar ck}{V}} \cos(kz) \cos(ckt) \varepsilon_y \quad (50)$$

next

$$\nabla \times \langle B \rangle = \nabla \times \left(-2\alpha\varepsilon_y \sqrt{\frac{\mu_0 \hbar \omega}{V}} \cos(kz) \sin(\omega t) \right) \quad (51)$$

$$= -2\alpha k \varepsilon_x \sqrt{\frac{\mu_0 \hbar \omega}{V}} \sin(kz) \sin(\omega t) \quad (52)$$

$$= -2\alpha k \sqrt{\frac{\mu_0 \hbar ck}{V}} \sin(kz) \sin(ckt) \varepsilon_x \quad (53)$$

and

$$\mu_0 \varepsilon_0 \frac{\partial \langle E \rangle}{\partial t} = -2\alpha\omega\varepsilon_x \mu_0 \sqrt{\frac{\varepsilon_0 \hbar \omega}{V}} \sin(kz) \sin(\omega t) \quad (54)$$

$$= -2\alpha k \sqrt{\frac{\mu_0 \hbar ck}{V}} \sin(kz) \sin(ckt) \varepsilon_x \quad (55)$$

3.5 Classical Energy

The classical energy of the EM field is

$$U = \int_V d\mathbf{r} \left(\frac{\varepsilon_0}{2} \langle E \rangle^2 + \frac{1}{2\mu_0} \langle B \rangle^2 \right) \quad (56)$$

$$= \int_V d\mathbf{r} \left(\frac{\varepsilon_0}{2} \left(2\alpha\varepsilon_x \sqrt{\frac{\hbar\omega}{\varepsilon_0 V}} \sin(kz) \cos(\omega t) \right)^2 + \frac{1}{2\mu_0} \left(-2\alpha\varepsilon_y \sqrt{\frac{\mu_0 \hbar\omega}{V}} \cos(kz) \sin(\omega t) \right)^2 \right) \quad (57)$$

$$= 2\alpha^2 \frac{\hbar\omega}{V} \int_V d\mathbf{r} (\sin^2(kz) \cos^2(\omega t) + \cos^2(kz) \sin^2(\omega t)) \quad (58)$$

$$= \alpha^2 \hbar\omega \quad (59)$$

In comparison, the energy from H_F is

$$\langle H_F \rangle = \hbar\omega \left(\langle \psi | N | \psi \rangle + \frac{1}{2} \right) = \hbar\omega \left(\alpha^2 + \frac{1}{2} \right) \quad (60)$$

So the classical energy is lower than the quantized energy.

3.6 Justification

If $|\alpha| \gg 1$ then the classical energy and the quantized energy are effectively the same, the $1/2$ term becoming too small to notice. In that case, the quantized behavior of the energy levels is effectively the same as the classical behavior.

4 Question 4

1. Show that the set of states A is an eigenbasis of $H_0 = H_A + H_F$. Find the eigenvalues.
2. Draw qualitatively on the energy axis the location of the five first eigenenergies of H_0 . Show that aside from the ground state, H_0 's eigenstates are bunched in pairs of degenerate states.
3. The coupling between the field and the atom is a dipole coupling and is given by

$$W = \gamma (a\sigma_+ + a^\dagger\sigma_-)$$

where $\gamma = -d \sin(kz_0) \sqrt{\hbar\omega/\varepsilon_0 V}$ where d is the dipole moment, determined experimentally.

- (a) Write the action of $\hat{W}$ on the states $|g, n\rangle$ and $|e, n\rangle$
- (b) Which physical processes to the operators $a\sigma_+$ and $a^\dagger\sigma_-$ describe?

4.1 Eigenbasis

I'll assume the meaning of the question is to prove the states in set A are eigenstates of the hamiltonian, not to prove they span the energy space.

$$H_0 |g, n\rangle = \left(\frac{\hbar\omega_A}{2} \sigma_z + \frac{\hbar\omega}{2} (\chi^2 + \Pi^2) \right) |g, n\rangle \quad (61)$$

$$= -\frac{\hbar\omega_A}{2} + \hbar\omega \left(n + \frac{1}{2} \right) |g, n\rangle \quad (62)$$

and

$$H_0 |e, n\rangle = \left(\frac{\hbar\omega_A}{2} \sigma_z + \frac{\hbar\omega}{2} (\chi^2 + \Pi^2) \right) |e, n\rangle \quad (63)$$


$$= \frac{\hbar\omega_A}{2} + \hbar\omega \left(n + \frac{1}{2} \right) |e, n\rangle \quad (64)$$

4.2 Drawing

Since $\omega_A = \omega$ the energies are now

$$H_0 |g, n\rangle = \hbar\omega n |g, n\rangle \quad H_0 |e, n\rangle = \hbar\omega (n+1) |e, n\rangle \quad (65)$$

we can see that the ground state energy is 0, then every energy level has is degenerate with 2 options for achieving that energy.



4.3 Coupling

4.3.1 Action of W

$$W |g, n\rangle = \gamma (a\sigma_+ + a^\dagger\sigma_-) |g, n\rangle \quad (66)$$

$$= \gamma a\sigma_+ |g, n\rangle \quad (67)$$

$$= \gamma\sqrt{n} |e, n-1\rangle \quad (68)$$

$$W |e, n\rangle = \gamma (a\sigma_+ + a^\dagger\sigma_-) |e, n\rangle \quad (69)$$

$$= \gamma a^\dagger\sigma_- |e, n\rangle \quad (70)$$

$$= \gamma\sqrt{n+1} |g, n+1\rangle \quad (71)$$

and of course the $W |g, 0\rangle = 0$.

4.3.2 Physical Process

$a\sigma_+$ describes a photon being absorbed by the atom raising it to the excited state. $a^\dagger\sigma_-$ describes the opposite.

5 Rabi Oscillations

1. Find the state of the system $|\psi(t)\rangle$ at a time $t > 0$.
2. What is the probability $P_g(T)$ to find the atom in the ground state $|g\rangle$ at time T , when the atom leaves the resonator? Show that $P_g(T)$ is a periodic function.
3. For Rubidium atoms $d = 1.1 \times 10^{-26}$ C m and $\frac{\omega}{2\pi} = 5.0 \times 10^{10}$ Hz and the experiment is done in a resonator of volume $V = 1.87 \times 10^{-6}$ m³. The graph of $P_g(T)$ and of the real part of the fourier transform $J(\nu) = \int_0^\infty dT P_g(T) \cos(2\pi\nu T)$ are presented. Compare theory to experiment.

5.1 State at Time

First we need to check if the hamiltonian $H = H_0 + W$ can even be diagonalized and for that we check if H_0 and W share an eigenbasis. To check that we look at their commutator

$$[H_0, W] = \left[\left(\frac{\hbar\omega}{2} \sigma_z + \hbar\omega \left(N + \frac{1}{2} \right) \right), \gamma (a\sigma_+ + a^\dagger\sigma_-) \right] \quad (72)$$

$$= \frac{\hbar\omega\gamma}{2} ([\sigma_z, a\sigma_+ + a^\dagger\sigma_-] + 2[N, a\sigma_+ + a^\dagger\sigma_-]) \quad (73)$$

$$= \hbar\omega\gamma (a\sigma_+ - a^\dagger\sigma_- - a\sigma_+ + a^\dagger\sigma_-) \quad (74)$$

$$= 0 \quad (75)$$

So H can be diagonalized by diagonalizing H_0 and W together. We'll try to express the eigenstates of W using the eigenstates of H_0 . Since W moves excited states to ground states and vice versa while removing or adding a photon respectively I'll guess normalized vectors of the form

$$|\psi_{\pm, n}\rangle = \frac{1}{\sqrt{2}} (|e, n\rangle \pm |g, n+1\rangle) \quad (76)$$

since they represent superpositions of states with an excited atom and n photons and ground state atoms with $n+1$ photons. We'll check

$$W |\psi_{+, n}\rangle = W \frac{1}{\sqrt{2}} (|e, n\rangle + |g, n+1\rangle) \quad (77)$$

$$= \gamma (a\sigma_+ + a^\dagger\sigma_-) \frac{1}{\sqrt{2}} (|e, n\rangle + |g, n+1\rangle) \quad (78)$$

$$= \frac{\gamma}{\sqrt{2}} (a^\dagger |g, n\rangle + a |e, n+1\rangle) \quad (79)$$

$$= \frac{\gamma}{\sqrt{2}} \sqrt{n+1} (|g, n+1\rangle + |e, n\rangle) \quad (80)$$

$$= \gamma \sqrt{n+1} |\psi_{+, n}\rangle \quad (81)$$

and

$$W |\psi_{-,n}\rangle = W \frac{1}{\sqrt{2}} (|e, n\rangle - |g, n+1\rangle) \quad (82)$$

$$= \gamma (a\sigma_+ + a^\dagger\sigma_-) \frac{1}{\sqrt{2}} (|e, n\rangle - |g, n+1\rangle) \quad (83)$$

$$= \frac{\gamma}{\sqrt{2}} (a^\dagger |g, n\rangle - a |e, n+1\rangle) \quad (84)$$

$$= \frac{\gamma}{\sqrt{2}} \sqrt{n+1} (|g, n+1\rangle - |e, n\rangle) \quad (85)$$

$$= -\gamma \sqrt{n+1} |\psi_{-,n}\rangle \quad (86)$$

So those are indeed eigenstates of W too, therefore these are also eigenstates of the full hamiltonian. The energies of these eigenstates are

$$E_{\pm,n} = \hbar\omega (n+1) \pm \gamma \quad (87)$$

We can now express the initial state as a combination of these

$$|\psi(0)\rangle = |e, 0\rangle = \frac{1}{\sqrt{2}} (|\psi_{+,0}\rangle + |\psi_{-,0}\rangle) \quad (88)$$

and use the time independent propagator

$$|\psi(t)\rangle = \frac{1}{\sqrt{2}} U(t) (|\psi_{+,0}\rangle + |\psi_{-,0}\rangle) \quad (89)$$

$$= \frac{1}{\sqrt{2}} e^{-i(\omega + \frac{\gamma}{\hbar})t} |\psi_{+,0}\rangle + \frac{1}{\sqrt{2}} e^{-i(\omega - \frac{\gamma}{\hbar})t} |\psi_{-,0}\rangle \quad (90)$$

$$= \frac{1}{\sqrt{2}} e^{-i\omega t} \left(e^{-\frac{i\gamma}{\hbar}t} |\psi_{+,0}\rangle + e^{\frac{i\gamma}{\hbar}t} |\psi_{-,0}\rangle \right) \quad (91)$$

$$= \frac{1}{2} e^{-i\omega t} \left(e^{-\frac{i\gamma}{\hbar}t} (|e, 0\rangle + |g, 1\rangle) + e^{\frac{i\gamma}{\hbar}t} (|e, 0\rangle - |g, 1\rangle) \right) \quad (92)$$

$$= \frac{1}{2} e^{-i\omega t} \left(\left(e^{-\frac{i\gamma}{\hbar}t} + e^{\frac{i\gamma}{\hbar}t} \right) |e, 0\rangle + \left(e^{-\frac{i\gamma}{\hbar}t} - e^{\frac{i\gamma}{\hbar}t} \right) |g, 1\rangle \right) \quad (93)$$

$$= e^{-i\omega t} \left(\cos\left(\frac{\gamma t}{\hbar}\right) |e, 0\rangle - i \sin\left(\frac{\gamma t}{\hbar}\right) |g, 1\rangle \right) \quad (94)$$

5.2 Probability of Ground State

$$P_g(T) = |\langle g, 1 | \psi(T) \rangle|^2 = \sin^2\left(\frac{\gamma T}{\hbar}\right) \quad (95)$$

which is periodic.

5.3 Rubidium

We found the probability to be periodic and since it's a sine square the period is twice the expression in the sine So

$$\nu = f/2\pi = \frac{\gamma}{\pi\hbar} \quad (96)$$

$$= \frac{d \sin(kz_0) \sqrt{\hbar\omega/\varepsilon_0 V}}{\pi\hbar} \quad (97)$$

$$= 1.1 \times 10^{-26} \text{ C m} \cdot \sqrt{\frac{2 \cdot 5.0 \times 10^{10} \text{ Hz}}{\pi \cdot 1.05 \times 10^{-34} \text{ J s} \cdot 8.85 \times 10^{-12} \text{ F m}^{-1} \cdot 1.87 \times 10^{-6} \text{ m}^3}} \quad (98)$$

$$= 47 \text{ kHz} \quad (99)$$

which fits the peak in the frequency graph nicely.