

Notes on Template Games and Differential Logic

Daniel Rogozin

Contents

1 General Remarks

1

1 General Remarks

The results from [FGHW08] introduce the bicategorical interpretation of linear logic based on distributors and generalised species. The idea is that every formula A of linear logic is interpreted as a small category $\llbracket A \rrbracket$ and every derivation tree

$$\frac{\begin{array}{c} \pi \\ \dots \end{array}}{A_1, \dots, A_n \vdash A}$$

is interpreted as a distributor (or a profunctor)

$$\llbracket \pi \rrbracket : \llbracket A_1 \rrbracket \otimes \dots \otimes \llbracket A_n \rrbracket \not\rightarrow \llbracket A \rrbracket.$$

Recall that a distributor $M : A \not\rightarrow B$ is defined as a functor $M : A \times B^{op} \rightarrow \mathbf{Set}$. The category **Dist** of small categories and distributors is symmetric monoidal with the tensor product defined by the direct product of categories in **Cat**, the category of small categories and functors, and the unit defined by the terminal (a single-element) category. **Dist** is also *-autonomous (and, moreover, compact closed) with linear negation defined as $A \mapsto A^{op}$. We thus have the following isomorphism

$$\mathbf{Dist}(A \otimes B, C) \cong \mathbf{Dist}(B, A^{op} \otimes C)$$

natural A , B and C , providing the currfication in **Dist**.

The exponential modality $A \mapsto !A$ is interpreted using the 2-monad **Sym** : **Cat** $\rightarrow$ **Cat** sending each small category $\mathcal{C}$ to the free symmetric monoidal category generated by $\mathcal{C}$. **Sym**($\mathcal{C}$) has objects defined as finite sequences $A_1 \dots A_n$ for objects from $\mathcal{C}$ for where $A_i \in \mathcal{C}$ for $1 \leq i \leq n$, and morphisms

$$(\sigma, f_1 \dots f_n) : A_1 \dots A_n \rightarrow B_1 \dots B_n$$

defined as pairs $(\sigma, f_1 \dots f_n)$ consists of a permutation $\sigma \in \mathcal{S}_n$ on the set of n -elements along with a sequence of morphisms from $\mathcal{C}$ for $1 \leq k \leq n$

$$f_k : A_k \rightarrow B_{\sigma(k)}.$$

The category $\mathbf{Sym}(\mathcal{C})$, as strict monoidal, comes with a pair of functors $\otimes : \mathbf{Sym}(\mathcal{C}) \times \mathbf{Sym}(\mathcal{C}) \rightarrow \mathbf{Sym}(\mathcal{C})$ and $I_A : \mathbf{1} \rightarrow \mathbf{Sym}(\mathcal{C})$ defined as the concatenation and the empty sequence respectively. $\mathbf{Sym}(\mathcal{C})$ with those two functors also forms a monoid in $\mathbf{Cat}$, and such a monoid is commutative up to isomorphism γ (the symmetry of the monoidal category):

$$\begin{array}{ccc} \mathbf{Sym}(\mathcal{C}) \times \mathbf{Sym}(\mathcal{C}) & \xrightarrow{(21)} & \mathbf{Sym}(\mathcal{C}) \times \mathbf{Sym}(\mathcal{C}) \\ & \searrow \otimes \quad \xRightarrow{\alpha} \quad \swarrow \otimes & \\ & \mathbf{Sym}(\mathcal{C}) & \end{array}$$

where (21) is the symmetry of the cartesian category $\mathbf{Sym}(\mathcal{C}) \times \mathbf{Sym}(\mathcal{C})$. The exponential modality $A \mapsto !A$ is, thus, interpreted in $\mathbf{Dist}$ by turning $\mathbf{Sym}$ into a 2-comonad

$$\mathbf{Sym} : \mathbf{Dist} \rightarrow \mathbf{Dist}$$

using the distributivity law between $\mathbf{Sym}$ and the presheaf construction in $\mathbf{Cat}$. Further, every functor $F : \mathcal{C} \rightarrow \mathcal{D}$ induces an adjoint pair $L_F \dashv R_F$ of distributors $L_F : A \not\rightarrow B$ and $R_F : B \not\rightarrow A$ in the bicategory $\mathbf{Dist}$, where the distributors L_F and R_F are given as the following presheaves

$$\begin{aligned} L_F(B, A) &= \mathcal{D}(FB, A) : B^{op} \times A \rightarrow \mathbf{Set} \\ R_F(A, B) &= \mathcal{D}(A, FB) : A^{op} \times B \rightarrow \mathbf{Set} \end{aligned}$$

The comonoid structure $(\mathbf{Sym}(\mathcal{C}), d_{\mathcal{C}}, e_{\mathcal{C}})$ of the exponential modality $\mathbf{Sym}$ is given by the following distributors

$$\begin{aligned} d_{\mathcal{C}} &= R_{\otimes_{\mathcal{C}}} : \mathbf{Sym}(\mathcal{C}) \not\rightarrow \mathbf{Sym}(\mathcal{C}) \otimes \mathbf{Sym}(\mathcal{C}) \\ e_{\mathcal{C}} &= R_{I_{\mathcal{C}}} : \mathbf{Sym}(\mathcal{C}) \not\rightarrow \mathbf{1} \end{aligned}$$

right adjoint to $\otimes_{\mathcal{C}}$ and $I_{\mathcal{C}}$ respectively. Note that $(\mathbf{Sym}(\mathcal{C}), d_{\mathcal{C}}, e_{\mathcal{C}})$ is not commutative, but only symmetric, so it is commutative only up to isomorphism:

$$\begin{array}{ccc} & \mathbf{Sym}(\mathcal{C}) & \\ d_{\mathcal{C}} \swarrow & \xRightarrow{\alpha} & \searrow d_{\mathcal{C}} \\ \mathbf{Sym}(\mathcal{C}) \otimes \mathbf{Sym}(\mathcal{C}) & \xrightarrow{(21)} & \mathbf{Sym}(\mathcal{C}) \otimes \mathbf{Sym}(\mathcal{C}) \end{array}$$

where γ is the symmetry of $\mathbf{Sym}(\mathcal{C}) \otimes \mathbf{Sym}(\mathcal{C})$, which is a distributor in $\mathbf{Dist}$ as well.

References

- [FGHW08] Marcelo Fiore, Nicola Gambino, Martin Hyland, and Glynn Winskel. The cartesian closed bicategory of generalised species of structures. *Journal of the London Mathematical Society*, 77(1):203–220, 2008.