

Notes on Ross Street's Quantum Groups

Daniel Rogozin

1 Duality between geometry and algebra

Let $\mathbf{k}$ be a field, a $\mathbf{k}$ -algebra is a ring A equipped with a ring morphism $\eta : \mathbf{k} \rightarrow A$. Note that either A is trivial or η is injective. *Scalar multiplication* $\mathbf{k} \times A \rightarrow A$ is defined by $(\kappa, a) \mapsto \eta(\kappa)a$. Let A and B be $\mathbf{k}$ -algebras, a $\mathbf{k}$ -algebra morphism is a ring morphism such that the following triangle commutes:

$$\begin{array}{ccc} & \mathbf{k} & \\ \eta \swarrow & & \searrow \eta \\ A & \xrightarrow{f} & B \end{array}$$

$\text{Alg}_{\mathbf{k}}(A, B)$ is the set of all $\mathbf{k}$ -algebra morphisms from A to B .

Let X be a compact Hausdorff space, $C(X)$ is the algebra of continuous complex-valued functions $a : X \rightarrow \mathbb{C}$ with the operations defined pointwisely.

A continuous function $f : X \rightarrow Y$ induces an algebra morphism $C(f) : C(Y) \rightarrow C(X)$ with $C(f)(b) = a$ for $a(x) = b(f(x))$. The unique $X \rightarrow 1$ gives the algebra morphism $\eta : \mathbb{C} = C(1) \rightarrow C(X)$, whilst each point $x : 1 \rightarrow X$ of the space gives an algebra morphism $C(X) \rightarrow \mathbb{C}$.

$C(X)$ is a *commutative C^* -algebra*, and, moreover, each commutative C^* -algebra is isomorphic to $C(X)$ for some compact Hausdorff space X ; and each C^* -algebra morphism $C(Y) \rightarrow C(X)$ has the form $C(f)$ for a unique continuous map $f : X \rightarrow Y$. This fact is known as *Gelfand duality*.

A variety is a space of solutions to polynomials in several variables. Consider the variety given by a polynomial $x^2 + 2y^3 = z^4$ over $\mathbf{k}$. We pass to the $\mathbf{k}$ -algebra:

$$A = \mathbf{k}[x, y, z]/(x^2 + 2y^3 = z^4)$$

where $\mathbf{k}[x, y, z]/(x^2 + 2y^3 = z^4)$ is the quotient algebra obtained from $\mathbf{k}[x, y, z]$ by identifying those polynomials that can be transformed to $x^2 + 2y^3 = z^4$ by using the algebra axioms.

Let B be a $\mathbf{k}$ -algebra, a $\mathbf{k}$ -algebra morphism $f : \mathbf{k}[x, y, z] \rightarrow B$ is uniquely determined by its values on x, y, z . So we have a bijection:

$$\text{Alg}_{\mathbf{k}}(\mathbf{k}[x, y, z], B) \cong B^3$$

and, similarly:

$$\mathrm{Alg}_{\mathbf{k}}(A, B) \cong \{(u, v, w) \in B^3 \mid u^2 + 2v^3 = w^4\}$$

a $\mathbf{k}$ -algebra morphism $A \rightarrow B$ corresponds to a map of varieties in the reverse direction.

A topological group is a group in the category of topological spaces and continuous maps. A Lie group is a group in the category of smooth manifolds and smooth maps.

We will be studying groups in the category $(\mathbf{CommAlg}_{\mathbf{k}})^{op}$ of commutative $\mathbf{k}$ -algebras and reversed morphisms, *affine groups* over $\mathbf{k}$. Algebraically, they are called *commutative Hopf algebras* over $\mathbf{k}$. A commutative Hopf algebra H over $\mathbf{k}$ is concised of the following choice of morphisms: $\varepsilon : H \rightarrow \mathbf{k}$ (counit), $\delta : H \rightarrow H \otimes_{\mathbf{k}} H$ (comultiplication) and $\nu : H \rightarrow H$ (antipode). Let A be a commutative $\mathbf{k}$ -algebra, $\mathrm{Alg}_{\mathbf{k}}(H, A)$ is the group of A -points in H .

Affine monoids (or commutative bialgebras) are concised of counits and comultiplications without antipods.

Example 1.1. Let $M(2)$ be $\mathbf{k}[a, b, c, d]$ as a commutative $\mathbf{k}$ -algebra. Define a counit $\varepsilon : M(2) \rightarrow \mathbf{k}$ is given by $\varepsilon(a) = \varepsilon(d) = 1$ and $\varepsilon(b) = \varepsilon(c) = 0$. One has

$$\mathbf{k}[a, b, c, d] \otimes_{\mathbf{k}} \mathbf{k}[a, b, c, d] \cong \mathbf{k}[a, b, c, d, a', b', c', d']$$

with the coprojections

$$\mathbf{k}[a, b, c, d] \longrightarrow \mathbf{k}[a, b, c, d, a', b', c', d'] \longleftarrow \mathbf{k}[a, b, c, d]$$

defined as $a, b, c, d \mapsto a, b, c, d$ and $a, b, c, d \mapsto a', b', c', d'$. The comultiplication $\delta : M(2) \rightarrow M(2) \otimes_{\mathbf{k}} M(2)$ is given by

$$a, b, c, d \mapsto aa' + bc', ab' + bd', ca' + dc', cb' + dd'.$$

So $M(2)$ is a commutative bialgebra. We also have a monoid isomorphism

$$\mathrm{Alg}_{\mathbf{k}}(M(2), A) \cong \mathrm{Mat}(2, A)$$

where $\mathrm{Mat}(2, A)$ is the multiplicative monoid of 2×2 matrices with entries in A . $M(2)$ is the coordinate $\mathbf{k}$ -algebra of the variety of 2×2 matrices.

To obtain the coordinate $\mathbf{k}$ -algebra of the general linear group, one can take:

$$GL(2) = \mathbf{k}[a, b, c, d, x]/(xad - xbc = 1).$$

There is an epimorphic $\mathbf{k}$ -algebra morphism $M(2) \rightarrow GL(2)$ which induces a bialgebra structure on $GL(2)$ from $M(2)$. The antipode $\nu : M(2) \rightarrow M(2)$ with $a, b, c, d, x \mapsto xd, -xb, -xc, xa, ad - bc$, so $GL(2)$ is a commutative Hopf algebra.

2 The quantum general linear group

Quantisation deforms commutative algebras to non-commutative since $xy = e^{\hbar}yx$. Quantum spaces correspond to more general $\mathbf{k}$ -algebras that do not have to be commutative.

Fix a field $\mathbf{k}$ and $q \in \mathbf{k}$ such that $q \neq 0$. Let $\mathbf{k}\langle x_1, \dots, x_n \rangle$ denote the $\mathbf{k}$ -algebra of polynomials in non-commuting indeterminates $x_1, \dots, x_n$. As a vector space $\mathbf{k}$, a basis is given by

$$x_{\xi(1)}^{m_1}, \dots, x_{\xi(r)}^{m_r}$$

for $r < \mathbb{N}$, $m_1, \dots, m_r \in \mathbb{N}^+$, $\xi : \{1, \dots, r\} \rightarrow \{1, \dots, n\}$. Note that $\mathbf{k}[x, y] = \mathbf{k}\langle x, y \rangle / (xy = yx)$.

The coordinate algebra of quantum 2×2 matrices is defined by

$$M_q(2) = \mathbf{k}\langle a, b, c, d \rangle / R$$

where R is the system of equations

$$\begin{aligned} ab &= q^{-1}ba, \quad ac = q^{-1}ca, \quad cd = q^{-1}dc, \quad bd = q^{-1}db, \\ bc &= cb, \quad ad - da = (q^{-1} - q)bc. \end{aligned}$$

The monomials $a^{m_1}b^{m_2}c^{m_3}d^{m_4}$ form a basis for the algebra as the vector space.

$$\text{Alg}_{\mathbf{k}}(M_1(2), A) \cong \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Mat}(2, A) \mid R \text{ holds} \right\}$$

Theorem 2.1. Let $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $\begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$ be two A -points of $M_q(2)$ such that each entry of the first matrix commute with each entry of the second, then:

1. The matrix product $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$ is an A -point of $M_q(2)$.
2. The q -determinant

$$\det_q \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - q^{-1}bc$$

commutes with each of a, b, c, d and the following holds

$$\det_q \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} \right) = \det_q \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) \times \det_q \left(\begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} \right).$$

3. If $\det_q \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right)$ is invertible in A , then the following is an A -point of $M_{q^{-1}}(2)$:

$$\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right)^{-1} = \left(\det_q \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) \right)^{-1} \begin{pmatrix} d & -qb \\ -q^{-1}c & a \end{pmatrix}.$$

The *quantum plane* is defined as the following $\mathbf{k}$ -algebra

$$\mathbb{A}_q^{2|0} = \mathbf{k}\langle x, y \rangle / (xy = q^{-1}yx).$$

The monomials $x^m y^n$ for $m, n \in \mathbb{N}$ form a basis for $\mathbb{A}_q^{2|0}$ as a vector space. We also need the quantised version of a Grassmannian algebra in two variables also called the *quantum superplane*:

$$\mathbb{A}_q^{0|2} = \mathbf{k}\langle \xi, \eta \rangle / (\xi^2 = \eta^2 = 0 = \xi\eta + q\eta\xi).$$

Monomials $\xi^m \eta^n$ for $m, n < 2$ form a basis for this algebra.

An A -point of B is called *generic* when the algebra morphism $B \rightarrow A$ is injective.

Theorem 2.2. Let (x, y) be a generic A -point of $\mathbb{A}_q^{2|0}$ and let (ξ, η) be a generic A -point of $\mathbb{A}_q^{0|2}$. Assume $a, b, c, d \in A$ commute with x, y, ξ, η . Put:

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}, \begin{pmatrix} x'' \\ y'' \end{pmatrix} = \begin{pmatrix} a & c \\ b & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}, \begin{pmatrix} \xi' \\ \eta' \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix}$$

If $q^2 = -1$, then the following are equivalent:

1. (x', y') and (x'', y'') are points of $\mathbb{A}_q^{2|0}$;
2. (x', y') is a point of $\mathbb{A}_q^{2|0}$ and (ξ', η') is a point of $\mathbb{A}_q^{0|2}$;
3. $\begin{pmatrix} a & c \\ b & d \end{pmatrix}$ is a point of $M_q(2)$.

If $q^2 = -1$, then one only has $(3) \Rightarrow (1) \& (2)$.

Proof.

- $(1) \Leftrightarrow (3)$.

Let (x', y') be a point of $\mathbb{A}_q^{2|0}$ iff $x'y' = q^{-1}y'x'$, that is, iff $(ax + by)(cx + dy) = q^{-1}(cx + dy)(ax + by)$. One can multiply out the products using the fact that a, b, c, d each commute with x and y . (x, y) is generic, so one can equate coefficients of x^2, y^2, xy . The single equation is equivalent to the following set of equations:

$$ac = q^{-1}ca, bd = q^{-1}db, ad - da = q^{-1}cb - qbc. \quad (1)$$

If we interchange b and c , then we can see that (x'', y'') is a point of $\mathbb{A}_q^{2|0}$ iff

$$ab = q^{-1}ba, cd = q^{-1}dc, ad - da = q^{-1}bc - qcb. \quad (2)$$

Therefore we have $q^{-1}cb - qbc = q^{-1}bc - qcb$, thus, $(q + q^{-1})(bc - cb) = 0$ and, thus, $bc = cb$ provided $q^2 \neq -1$.

- (2) $\Leftrightarrow$ (3) Let (ξ', η') be a point of $\mathbb{A}_q^{0|2}$ iff $0 = (a\xi + b\eta)^2 = (c\xi + d\eta)^2 = (a\xi + b\eta)(c\xi + d\eta)$. Using $\xi^2 = \eta^2 = 0$, we have $ab\xi\eta + ba\eta\xi = 0$ and $cd\xi\eta + dc\eta\xi = 0$ and $ab\xi\eta + bc\eta\xi + q(cb\xi\eta + da\eta\xi) = 0$. Now we use $\xi\eta = -q\eta\xi$ and the fact that η and ξ in A , so we have $-qab + ba = 0$ and $-qcd + dc = 0$ and also $-q(ad + qcb) + bc + qda = 0$.

□

Proof of Theorem 2.1.

1. Let B be the free $\mathbf{k}$ -algebra containing the indeterminates $a, b, c, d, a', b', c', d', x, y$ satisfying the relations from Theorem 2.1 and Theorem 2.2. Then (x, y) is generic and $\begin{pmatrix} a & c \\ b & d \end{pmatrix}$ and $\begin{pmatrix} a' & c' \\ b' & d' \end{pmatrix}$ are B -points of $M_q(2)$. By Theorem 2.2, $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ and $\begin{pmatrix} a' & c' \\ b' & d' \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ are B -points of $\mathbb{A}_q^{2|0}$. Each coordinate in the first of these commutes with all of a', b', c', d' whilst coordinates in the second commute with a, b, c, d . Also $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ is also generic since when composed with $B \rightarrow \mathbb{A}_q^{2|0}$ for which $(a, b, c, d, x, y) \mapsto (1, 0, 0, 1, x, y)$ one gets (x, y) , which is generic. For the same reason, $\begin{pmatrix} a' & c' \\ b' & d' \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ is generic. By Theorem 2.2, $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \begin{pmatrix} a' & c' \\ b' & d' \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ and $\begin{pmatrix} a' & c' \\ b' & d' \end{pmatrix} \begin{pmatrix} a & c \\ b & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ are B -points of $\mathbb{A}_q^{2|0}$. By Theorem 2.2, $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \begin{pmatrix} a' & c' \\ b' & d' \end{pmatrix}$ is a B -point of $\mathbb{A}_q^{2|0}$. To obtain the result for A , one should apply $B \rightarrow A$ for which $(a, b, \dots, d', x, y) \mapsto (a, b, \dots, d', 0, 0)$.

2. (2) and (3) are straightforward.

□

The *quantum general linear group* is defined from 2×2 matrices by inverting the determinant:

$$\mathrm{GL}_q(2) = M_q(2)[t]/(ta = at, tb = bt, tc = ct, dt = td, t \det_q = 1).$$

The *quantum special linear group* is defined by requiring that the determinant is equal to 1:

$$\mathrm{SL}_q(2) = M_q(2)(\det_q = 1).$$

3 Modules and Tensor Products

Let R be a ring (not necessarily commutative). R^{op} stands for the ring with opposite multiplication:

$$R \times R \xrightarrow{\sigma} R \times R \xrightarrow{\mu} R$$

R is commutative if $R = R^{op}$.

A *left R -module* is an Abelian group M along with a function $R \times M \rightarrow M$ with $(r, m) \mapsto rm$ called *scalar multiplication* such that:

- $1m = m$,
- $(r + r')m = rm + r'm$,
- $(rs)m = r(sm)$,
- $r(m + m') = rm + rm'$.

A *right R -module* is defined similarly $M \times R \rightarrow M$.

A left R^{op} -module structure on an Abelian group M “is the same” as a right R -module structure. If $\mu : R \times M \rightarrow M$ is a scalar multiplication for a left R^{op} -module iff $M \times R \xrightarrow{\sigma} R \times M \xrightarrow{\mu} M$ is one for a right R -module. If R is a field, then an R -module is a vector space.

A subset X of an R -module M generates M (or spans M), when, for each $m \in M$, there exist $r_1, \dots, r_n \in R$ and $x_1, \dots, x_n \in X$ such that

$$m = r_1x_1 + \dots r_nx_n$$

M is finitely generated if X is finite. A subset $X \subseteq M$ is *linearly independent* if $x_1, \dots, x_n \in X$ and x_i and x_j are distinct for each $i, j \leq n$ such that $i \neq j$, then $r_1x_1 + \dots + r_nx_n = 0$ for $r_1, \dots, r_n \in R$ implies that $r_1 = \dots = r_n = 0$.

An R -module F is *free* if it is generated by some linearly independent subset. Each set X determines an R -module

$$\mathcal{F}_R(X) = \{r_1x_1 + \dots + r_nx_n \mid r_i \in R, x_i \in X, n < \omega\}$$

$\mathcal{F}_R(X)$ is free and generated by X .

Let M, N be R -modules, a function $f : M \rightarrow N$ is (left) R -linear (or an *R -module morphism*) when $f(m + m') = fm + fm'$ and $f(rm) = rf(m)$. $\text{Hom}_R(M, N)$ is the Abelian group of R -linear functions with addition given by $(f + g)(m) = f(m) + g(m)$. If R is commutative, then $\text{Hom}_R(M, N)$ is an R -module. Clearly, one has the following isomorphism of Abelian groups:

$$\text{Hom}_R(\mathcal{F}_R(X), M) \cong M^X$$

where the addition on M^X is pointwise.

A subset H is a *submodule* of an R -module M if it is closed under addition and scalar multiplication. So we have an equivalence relation $\equiv_H$ on M :

$$m \equiv_H m' \Leftrightarrow m - m' \in H$$

The equivalence class containing $m \in M$ is $m + H = \{m + h \mid h \in H\}$ is called the *H -coset containing m* . The set M/H of H -cosets is an R -module itself via

- $(m + H) + (n + H) = (m + n) + H$,

- $r(m + H) = rm + H$.

There is a surjective R -linear function $\rho : M \rightarrow M/H$ for which $\rho : m \mapsto m + H$. For each R -linear $g : M \rightarrow N$ such that $g(m) = 0$ for each $m \in H$, there is a unique R -linear $\hat{g} : M/H \rightarrow N$ such that $\hat{g} \circ \rho = g$. The kernel $\ker(f) = \{m \in M \mid f(m) = 0\}$ for any R -linear $f : M \rightarrow N$ is a submodule of M . Therefore, the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \rho \downarrow & & \uparrow \\ M/\ker(f) & \xrightarrow{\cong} & \text{im}(f) \end{array}$$

The submodule $\langle X \rangle$ generated by a subset X of an R -module M is the smallest submodule of M containing X .

Let M be a right R -module and let N be a left R -module. Let A be an Abelian group, then a function $f : M \times N \rightarrow A$ is R -bilinear if it satisfies the following:

- $f(m, n + n') = f(m, n) + f(m, n')$,
- $f(m + m', n) = f(m, n) + f(m', n)$,
- $f(mr, n) = f(m, rn)$.

$\mathbf{Bil}_R(M, N; A)$ is the Abelian group, a subgroup of $A^{M \times N}$ of R -bilinear functions $f : M \times N \rightarrow A$. Let us construct $\lambda : M \times N \rightarrow M \otimes_R N$.

Let B be the subset of the Abelian group $\mathbf{F}_{\mathbb{Z}}(M \times N)$ consisting of the elements of the following form for $m, m' \in M$, $n, n' \in N$ and $r \in R$:

- $(m + m', n) - (m, n) - (m', n)$,
- $(m, n + n') - (m, n) - (m, n')$,
- $(mr, n) - (m, rn)$

Put

$$M \otimes_R N := \mathbf{F}_{\mathbb{Z}}(M \times N)/(B)$$

Then one has the following isomorphisms:

$$\begin{aligned} \text{Hom}_{\mathbb{Z}}(M \otimes_R N, A) &= \text{Hom}_{\mathbb{Z}}(\mathbf{F}_{\mathbb{Z}}(M \times N)/(B), A) \cong \\ &\{g \in \text{Hom}_{\mathbb{Z}}(\mathbf{F}_{\mathbb{Z}}(M \times N), A) \mid \forall (m, n) \in B \ g(m, n) = 0\} \cong \\ &\{f \in A^{M \times N} \mid f \text{ is } R\text{-bilinear}\} = \mathbf{Bil}_R(M, N; A) \end{aligned}$$


Take $(m, n) \in M \times N$, put $m \otimes n := \lambda(m, n)$. Elements of $M \otimes_R N$ has the following form for $m_1, \dots, m_k \in M$ and $n_1, \dots, n_k \in N$

$$\sum_{i=1}^k m_i \otimes n_i$$

Such elements satisfy the following equalities:

- $(m + m') \otimes n = m \otimes n + m' \otimes n$,
- $m \otimes (n + n') = m \otimes n + m \otimes n'$,
- $mr \otimes n = m \otimes rn$.

Let R and S be rings, a *module M from R to S* , written as $M : R \nrightarrow S$, is an Abelian group M endowed with a left R -module structure and a right S -module structure related by $(rm)s = r(ms)$ for $r \in R$, $m \in M$ and $s \in S$. So one can view tensor product as the following triangle:



So $M \otimes_S N$ is a module from R to T by putting:

$$r(m \otimes n)t = (rm) \otimes (nt)$$

Let R and S be rings and let X be a set, a free module from R to S generated by X , denoted by $\mathcal{F}_R^S(X)$, consists of the elements of the form:

$$\sum_{i=1}^i r_i x_i s_i$$


for $i < \omega$, $r_i \in R$, $s_i \in S$ and $x_i \in X$.

With every module $M : R \nrightarrow S$ we have:

$$\text{Hom}_R^S(\mathcal{F}_R^S(X), M) \cong M^X$$

where $\text{Hom}_R^S(N, M)$ is the Abelian group of left R /right S -module morphisms from $N \rightarrow M$.

Given rings and modules:



A function $f : M_1 \times \dots \times M_n \rightarrow L$ is *multilinear* if it satisfies the following for $r_i \in R_i$, $m_i, m'_i \in M_i$:

- $f(m_1, \dots, m_i + m'_i, \dots, m_n) = f(m_1, \dots, m_i, \dots, m_n) + f(m_1, \dots, m'_i, \dots, m_n)$,

- $r_0 f(m_1, \dots, m_n) = f(r_0 m_1, \dots, m_n)$,
- $f(m_1, \dots, m_i r_i, m_{i+1}, \dots, m_n) = f(m_1, \dots, m_i, r_i m_{i+1}, \dots, m_n)$,
- $f(m_1, \dots, m_n) r_n = f(m_1, \dots, m_n r_n)$.

$\mathbf{Mult}(M_1, \dots, M_n; L)$ is the Abelian group of such functions f . So we can construct a module

$$M_1 \otimes_{R_1} M_2 \otimes_{R_2} \dots \otimes_{R_{n-1}} M_n : R_0 \not\rightarrow R_n$$

along with the multilinear function

$$\lambda : M_1 \times \dots \times M_n \rightarrow M_1 \otimes_{R_1} M_2 \otimes_{R_2} \dots \otimes_{R_{n-1}} M_n$$

such that the following universal property holds:

$$\begin{array}{ccc} M_1 \times \dots \times M_n & \xrightarrow{\lambda} & M_1 \otimes_{R_1} M_2 \otimes_{R_2} \dots \otimes_{R_{n-1}} M_n \\ & \searrow \exists! g & \downarrow g \\ & & L \end{array}$$

Thus we have the following isomorphism of Abelian groups:

$$\mathbf{Mult}(M_1, \dots, M_n; L) \cong \text{Hom}_{R_0}^{R_n}(M_1 \otimes_{R_1} M_2 \otimes_{R_2} \dots \otimes_{R_{n-1}} M_n, L)$$

Note that the diagonal map $M \rightarrow M \otimes M$ with $m \mapsto m \otimes m$ does not preserve addition. The empty tensor product $M_1 \otimes \dots \otimes M_n$ for $n = 0$, which is R_0 as a module $R_0 \not\rightarrow R_0$ with multiplication in R as scalar multiplication. We have the following canonical isomorphisms:

$$R \otimes_R M \cong M \cong M \otimes_S S.$$

Given $M, M' : R \not\rightarrow S$, one writes $f : M \Rightarrow M' : R \not\rightarrow S$ or

$$\begin{array}{ccc} & M & \\ R & \begin{array}{c} \curvearrowright \\ \Downarrow f \\ \curvearrowleft \end{array} & S \\ & M' & \end{array}$$

to emphasise that $f : M \rightarrow M'$ is a left R - and right S -module morphism. Now assume we are given the following data:

$$\begin{array}{ccccccc} & M_1 & & M_2 & & \dots & & M_n \\ R_0 & \begin{array}{c} \curvearrowright \\ \Downarrow f_1 \\ \curvearrowleft \end{array} & R_1 & \begin{array}{c} \curvearrowright \\ \Downarrow f_2 \\ \curvearrowleft \end{array} & R_2 & \begin{array}{c} \curvearrowright \\ \dots \\ \curvearrowleft \end{array} & R_{n-1} & \begin{array}{c} \curvearrowright \\ \Downarrow f_n \\ \curvearrowleft \end{array} & R_n \\ & M'_1 & & M'_2 & & & & M'_n \end{array}$$

we obtain

$$f_1 \otimes \dots \otimes f_n : M_1 \otimes_{R_1} \dots \otimes_{R_{n-1}} M_n \Rightarrow M'_1 \otimes_{R_1} \dots \otimes_{R_{n-1}} M'_n : R_0 \not\rightarrow R_n$$

given by $(f_1 \otimes \dots \otimes f_n) \circ \lambda = \lambda \circ (f_1 \times \dots \times f_n)$.

Given a triangle of modules:

$$\begin{array}{ccc} R & \xrightarrow{M} & S \\ & \searrow L & \swarrow N \\ & T & \end{array}$$

we enrich the Abelian group $\text{Hom}_R(M, L)$ (or $\text{Hom}^T(N, L)$) of left R - (right T -) module morphisms with the following structure:

$$\begin{aligned} \text{Hom}_R(M, L) &: S \not\rightarrow T \\ \text{Hom}^T(N, L) &: R \not\rightarrow S \end{aligned}$$

with the following scalar multiplications:

$$\begin{aligned} (sft)(m) &= f(ms)t \\ (rgs)(n) &= rg(sn) \end{aligned}$$

Thus there are the following Abelian group isomorphisms:

$$\text{Hom}_S^T(N, \text{Hom}_R(M, L)) \cong \mathbf{Mult}(M, N; L) \cong \text{Hom}_R^S(M, \text{Hom}^T(N, L))$$

induced by the following canonical bijections:

$$(L^M)^N \cong L^{M \times N} \cong (L^N)^M$$

Thus we have:

$$\text{Hom}_S^T(N, \text{Hom}_R(M, L)) \cong \text{Hom}_R^T(M \otimes_S N, L) \cong \text{Hom}_R^S(M, \text{Hom}^T(N, L))$$

the isomorphisms are determined by the evaluation morphisms:

$$\begin{aligned} \text{ev}_M &: M \otimes_S \text{Hom}_R(M, L) \rightarrow L, m \otimes f \mapsto f(m), \\ \text{ev}^N &: \text{Hom}^T(N, L) \otimes_S N \rightarrow L, g \otimes n \mapsto g(n). \end{aligned}$$

3.1 Exercises

Exercise 3.1. Describe $\mathbb{Z}/(2) \otimes_{\mathbb{Z}} \mathbb{Z}/(5)$.

Proof. Take $a \in \mathbb{Z}/(2)$ and $b \in \mathbb{Z}/(5)$, then $a \otimes b$ ranges over the elements $\mathbb{Z}/(2) \otimes_{\mathbb{Z}} \mathbb{Z}/(5)$. Then:

$$\begin{aligned} 5(a \otimes b) &= a \otimes 5b = a \otimes 0 = a \otimes (0 \cdot 0) = 0 \otimes 0 = 0 \\ 2(a \otimes b) &= 2a \otimes b = 0 \otimes b = 0 \end{aligned}$$

So $a \otimes b = (5 - 2 \cdot 2)a \otimes b = 0$. So $\mathbb{Z}/(2) \otimes_{\mathbb{Z}} \mathbb{Z}/(5)$ is trivial. $\square$

Exercise 3.2. Let R, S be rings, describe the canonical ring structure $R \otimes_{\mathbb{Z}} S$. Is a function $\varphi : R \rightarrow R \otimes_{\mathbb{Z}} S$ with $r \mapsto r \otimes 1$ a ring morphism? Show that $R \otimes_{\mathbb{Z}} S$ is the coproduct in the category of commutative rings.

Proof. The multiplication and unit on $R \otimes_{\mathbb{Z}} S$ are given by:

$$(R \otimes S) \otimes (R \otimes S) \xrightarrow{1 \otimes \sigma_{R,S} \otimes 1} (R \otimes R) \otimes (S \otimes S) \xrightarrow{\mu \otimes \mu} R \otimes S$$

$$\mathbb{Z} \cong \mathbb{Z} \otimes \mathbb{Z} \xrightarrow{\eta \otimes \eta} R \otimes S$$

the multiplication is realised as

$$(r \otimes s)(r' \otimes s') = (rr') \otimes (ss'),$$

whereas the unit is realised as $1 = 1 \otimes 1$.

The function $\varphi : R \rightarrow R \otimes_{\mathbb{Z}} S$ with $\varphi : r \mapsto r \otimes 1$ is a morphism of Abelian groups, but it also preserves multiplication and unit:

$$\begin{aligned} \varphi(rr') &= rr' \otimes 1 = (r \otimes 1)(r' \otimes 1) = \varphi(r)\varphi(r') \\ \varphi(1) &= 1 \otimes 1 = 1 \end{aligned}$$

We have canonical injections:

$$R \xrightarrow{\varphi} R \otimes_{\mathbb{Z}} S \xleftarrow{\psi} S$$

realised as $\varphi(r) = r \otimes 1$ and $\psi(s) = 1 \otimes s$. Now take ring morphisms $f, g : R, S \rightarrow T$, we must show that there is a unique morphism $h : R \otimes S \rightarrow T$ such that the following diagram commutes:

$$\begin{array}{ccccc} & & R \otimes S & & \\ & \nearrow \varphi & \downarrow \exists! h & \nwarrow \psi & \\ R & \xrightarrow{f} & T & \xleftarrow{g} & S \end{array}$$

so we define:

$$h(r \otimes s) = h((r \otimes 1)(1 \otimes s)) = h(r \otimes 1)h(1 \otimes s) = f(r)g(s)$$

and h is bilinear and is an Abelian group morphism. h also preserves multiplication:

$$\begin{aligned} h((r \otimes s)(r' \otimes s')) &= h(rr' \otimes ss') = f(rr')g(ss') = \\ &= f(r)f(r')g(s)g(s') = f(r)g(s)f(r')g(s') = h(r \otimes s)h(r' \otimes s') \end{aligned}$$

□

Exercise 3.3. Show that there is a bijection between modules M from R to S and left $R \otimes_{\mathbb{Z}} S^{op}$ -module.

Proof. A module $M : R \nrightarrow S$ is an abelian group $\mu : R \otimes M \otimes S \rightarrow M$ such that $\mu(r \otimes m \otimes s) = rms$ and:

- $r'(rms)s' = (r'r)m(ss')$,
- $m = 1m1$.

A left $R \otimes S^{op}$ -module is an Abelian group M with an Abelian $\bar{\mu} : (R \otimes S) \otimes M \rightarrow M$ such that $\bar{\mu}(r \otimes m \otimes s) = (r \otimes s)m$ and:

- $(r'r \otimes ss')m = (r' \otimes s')((r \otimes s)m)$,
- $(1 \otimes 1)m = m$.

so the bijection is obtained via the following diagram:

$$\begin{array}{ccc}
 R \otimes M \otimes S & & \\
 \downarrow \cong & \searrow \mu & \\
 & & M \\
 R \otimes S \otimes M & \nearrow \bar{\mu} &
 \end{array}$$

□

Exercise 3.4. Describe the construction of $M \otimes_S N \otimes_T L$ explicitly.

Proof. Assume we are given modules $M : R \nrightarrow S$, $N : S \nrightarrow T$ and $L : T \nrightarrow U$. We have an Abelian group:

$$B := M \otimes_S N \otimes_T L$$

where $B \subseteq \mathcal{F}_{\mathbb{Z}}(M \times N \times L)$ consisting of the following elements:

- $(m + m', n, l) - (m, n, l) - (m', n, l)$,
- $(m, n + n', l) - (m, n, l) - (m, n', l)$,
- $(m, n, l + l') - (m, n, l) - (m, n, l')$,
- $(ms, n, l) - (m, sn, l)$,
- $(m, nt, l) - (m, n, tl)$.

$m \otimes n \otimes l$ is the equivalence class of (m, n, l) . We define $r(m \otimes n \otimes l)\mu = (rm) \otimes n \otimes (l\mu)$, so we have $M \otimes_S N \otimes_T L : R \nrightarrow U$. Thus:

$$\text{Hom}_R^U(M \otimes_S N \otimes_T L, K) \cong \mathbf{Mult}(M, N, L; K).$$

□

4 Cauchy modules

A module $M : R \nrightarrow S$ induces a module $M* = \text{Hom}_R(M, R) : S \nrightarrow R$ called the *left dual* of M . There is a canonical module morphism:

$$\rho_L^M : M^* \otimes_R L \rightarrow \text{Hom}_R(M, R)$$

given by $\rho_L^M : (u \otimes l)(m) = u(m)l$ for each left R -module L .

A module $M : R \nrightarrow S$ is *Cauchy* whenever ρ_L^M is iso for each L .

A module P is *projective* when, for all surjective module morphisms $e : L \rightarrow L'$ and all module morphisms $f : P \rightarrow L'$, there is a module morphism $g : P \rightarrow L$ such that:

$$\begin{array}{ccc} P & \xrightarrow{f} & L' \\ & \searrow g & \uparrow e \\ & & L \end{array}$$

A morphism $r : M \rightarrow N$ is said to be a *retraction* when there is a morphism $i : N \rightarrow M$ such that $r \circ i = 1_N$. In this case we say that N a retract of M .

Proposition 4.1. A module is projective iff P is a retract of some free module F .

Proof. 1. A retract Q of a projective P is projective.

Assume we are given the following retraction:

$$\begin{array}{ccccc} & & 1_Q & & \\ & \nearrow & & \searrow & \\ Q & \xrightarrow{i} & P & \xrightarrow{r} & Q \end{array}$$

Let $e : L \rightarrow L'$ be a surjective morphism and let $f : Q \rightarrow L'$, then $f \circ r : P \rightarrow L'$, and P is projective, there is a morphism $h : P \rightarrow L$ such that $e \circ h = f \circ r$. But:

$$e \circ (h \circ i) = (e \circ h) \circ i = f \circ r \circ i = f \circ 1_Q = f$$

Therefore $g = h \circ i$, and thus $e \circ g = f$.

2. Free modules are projective.

Let $e : L \rightarrow L'$ be surjective and take any $f : F(X) \rightarrow L'$. Then we can choose by the axiom of choice $g(x) \in L$ for each $x \in X$ such that $e(g(x)) = f(x)$. $F(X)$ is free, so we can extend g uniquely to a morphism $g : F(X) \rightarrow L$ and thus $e \circ g = f$.

3. For each module M there is a free module F and a surjective morphism $e : F \rightarrow M$. Let F be the free module $F(\underline{M})$, where $\underline{M}$ is the carrier of M . We take $e(m) = m$ and e is clearly onto.

4. Moreover, if $e : F \rightarrow P$ is onto and P is projective, then e is a retraction. $\square$

Theorem 4.1 (The fundamental theorem of Morita theory). Let $M : R \nrightarrow S$ be a module, then the following are equivalent:

1. M is Cauchy,
2. There is a morphism $d : S \Rightarrow M^* \otimes M : S \nrightarrow S$ such that the following composites are identities:

$$\begin{aligned} M &\cong M \otimes_S S \xrightarrow{1_M \otimes d} M \otimes_S M^* \otimes_S M \xrightarrow{\text{ev}_M \otimes 1_M} R \otimes_R M \cong M \\ M^* &\cong S \otimes_S M^* \xrightarrow{d \otimes 1_{M^*}} M^* \otimes_R M \otimes_S M^* \xrightarrow{1_{M^*} \otimes \text{ev}_M} M^* \otimes_R R \cong M^* \end{aligned}$$

3. There is a module $N : S \nrightarrow R$ and morphisms $e : M \otimes_S N \rightarrow R$ and $d : S \rightarrow N \otimes_R M$ such that the following composite is identity:

$$M \cong M \otimes_S S \xrightarrow{1_M \otimes d} M \otimes_S N \otimes_R M \xrightarrow{e \otimes 1_M} R \otimes_R M \cong M.$$

4. M is a finitely generated projective left R -module.

Proof.

1. (i) $\Rightarrow$ (ii).

Since $\rho_M^M : M^* \otimes_R M \rightarrow \text{Hom}_R(M, M)$ is an isomorphism, so there is an element of $M^* \otimes_R M$ that ρ_M^M maps to 1_M . This element determines a unique morphism $d : S \rightarrow M^* \otimes_R M$, whose value at $1 \in S$ is the element. Let us write $d(1) = \sum_i u_i \otimes m_i$. Thus the condition $\rho_M^M(d(1))(m) = m$ becomes $\sum_i u_i(m)m_i = m$ for each $m \in M$. Thus the first composite for the second item takes m to m . The second composite takes $u \in M^*$ to itself since:

$$u(m) = u(\sum_i u_i(m)m_i) = \sum_i u_i(m)u(m_i).$$

2. (ii) $\Rightarrow$ (iii). Take $N := M^*$, $e = \text{ev}_M$ and a given d .
3. (iii) $\Rightarrow$ (iv). Put

$$d(1) = \sum_{i=1}^k n_i \otimes m_i \in N \otimes_R M.$$

As far as the composite from (iii) is the identity, one has

$$\forall m \in M \quad \sum_i e(m \otimes n_i)m_i = m.$$

Thus M is generated by $m_1, \dots, m_k$. To see that M , take a surjective $s : L \rightarrow L'$ and $f : M \rightarrow L'$. Thus we can choose $l_1, \dots, l_k \in L$ with $s(l_i) = f(m_i)$. Define $g : M \rightarrow L$ by:

$$g(m) = \sum_i e(m \otimes n_i)l_i$$

Therefore we get:

$$s(g(m)) = \sum_i e(m \otimes n_i) s(l_i) = \sum_i e(m \otimes n_i) f(m_i) = f\left(\sum_i e(m \otimes n_i) m_i\right) = f(m)$$

4. (iv) $\Rightarrow$ (i).

First, a retract of a Cauchy module is Cauchy. Let us show that $M = F_R(X)$ is Cauchy for some finite $X = \{x_1, \dots, x_k\}$. One has $M^* = \text{Hom}_R(F(X), R) \cong R^k$ and $\text{Hom}_R(M, L) = \text{Hom}_R(F(X), L) \cong L^k$. Under these isomorphisms ρ_L^M carries across to the morphism $R^k \otimes_R L \rightarrow L^k$ with $(r_1, \dots, r_k) \otimes l \mapsto (r_1 l, \dots, r_k l)$ which has inverse $(l_1, \dots, l_k) \mapsto \sum_{i=1}^k u_i \otimes l_i$, in which $u_i \in R^k$ projects to 0 in all components except the i -th, where it projects to 1. Thus ρ_L^M is iso.

□

Let R and S be rings and let $f : S \rightarrow R$ be a ring morphism, we obtain two modules ${}_f R : S \nrightarrow R$ and $R_f : R \nrightarrow S$, and both of them have R as the underlying Abelian group. They have scalar multiplications:

- $S \times {}_f R \rightarrow {}_f R$ with $(s, r) \mapsto f(s)r$,
- $R_f \times S \rightarrow R_f$ with $(r, s) \mapsto rf(s)$,
- ${}_f R \times R \rightarrow {}_f R$ with $(r, r') \mapsto rr'$,
- $R \times R_f \rightarrow R_f$ with $(r', r) \mapsto r'r$.

For any module $L : R \nrightarrow T$ we have canonical isomorphism

$${}_f R \otimes_R L \cong L \cong \text{Hom}_R(R_f, L)$$

with $r \otimes l \mapsto l$ for ${}_f R \otimes_R L \cong L$ and $u \mapsto u(1)$ for $\text{Hom}_R(R_f, L) \cong L$.

Therefore we have:

$$(R_f)^* n \cong {}_f R$$

and, thus, R_f is Cauchy.

A module $M : R \nrightarrow S$ is *convergent* if there exists a ring morphism $f : S \rightarrow R$ and a module isomorphism $M \cong R_f$.

The *product* $\prod_{i \in I} M_i : R \nrightarrow S$ of a family of modules $\{M_i : R \nrightarrow S \mid i \in I\}$ consists of elements $\mathbf{m} = (m_i)_{i \in I}$ for $m_i \in M_i$. Addition and scalar multiplication are given by:

- $\mathbf{m} + \mathbf{m}' = (m_i + m'_i)_{i \in I}$,
- $r\mathbf{m} = (rm_i)_{i \in I}$

There are projections for each $j \in I$:

$$\pi_j : \prod_{i \in I} M_i \rightarrow M_j$$

given by $\pi_j(\mathbf{m}) = m_j$. There are injective module morphisms for each $j \in J$:

$$\iota_j : M_j \rightarrow \prod_{i \in I} M_i$$

given by $\iota_j(m) = \mathbf{m}$ where $m_j = m$ and $m_i = 0$ for $i \neq j$.

The *direct sum* $\sum_{i \in I} M_i : R \nrightarrow S$ is the submodule of $\prod_{i \in I} M_i$ consisting of those $\mathbf{m}$ such that $m_i = 0$ for finitely many $i \in I$. This is the submodule generated by the union $\cup_{i \in I} M_i$, so we write $\sum_{i \in I} m_i$ instead of $\mathbf{m} \in \sum_{i \in I} M_i$.

Proposition 4.2. There are module isomorphisms:

1. $\text{Hom}_R(\sum_{i \in I} M_i, L) \cong \prod_{i \in I} \text{Hom}_R(M_i, L)$ with $f \mapsto (f \circ \iota_i)_{i \in I}$.
2. $(\sum_{i \in I} M_i) \otimes_S N \cong \sum_{i \in I} M_i \otimes_S N$ with $(\sum_i m_i) \otimes n \mapsto \sum_i (m_i \otimes n)$.

Proof.

1. If $f \circ \iota_i = 0$ for each $i \in I$, then f is zero on each M_i . But if we have $(f_i : M_i \rightarrow L)_{i \in I}$, define $f : \sum M_i \rightarrow L$ by $f(\sum m_i) = \sum f_i(m_i)$.
2. Take any L :

$$\begin{aligned} \text{Hom}_R((\sum_{i \in I} M_i) \otimes_S N, L) &\cong \text{Hom}^S(\sum_{i \in I} M_i, \text{Hom}^R(N, L)) \cong \\ &\prod_{i \in I} \text{Hom}^S(M_i, \text{Hom}^R(N, L)) \cong \\ &\prod_{i \in I} \text{Hom}_R(M_i \otimes_S N, L) \cong \\ &\text{Hom}_R(\sum_{i \in I} M_i \otimes_S N, L) \end{aligned}$$

□

4.1 Exercises

Exercise 4.1. Let M be a finitely generated projective module over a commutative ring R . Then M^* is a finitely generated projective module and the canonical morphism $M \rightarrow M^{**}$ is bijective.

Proof. Let M be finitely generated and projective, then by the Fundamental Theorem of Morita theory, we have the morphisms $d : R \rightarrow M^* \otimes M$, $e : M \otimes M^* \rightarrow R$ such that:

$$\begin{array}{ccc} M^* & \xrightarrow{d \otimes 1} & M^* \otimes_R M \otimes_R M^* \\ & \searrow & \downarrow 1 \otimes e \\ & & M^* \end{array} \quad \begin{array}{ccc} M & \xrightarrow{1 \otimes d} & M \otimes M^* \otimes M \\ & \searrow & \downarrow e \otimes 1 \\ & & M \end{array}$$

Then we put $d' = R \xrightarrow{d} M^* \otimes_R M \xrightarrow{\sigma} M \otimes_R M^*$ and $e' = M^* \otimes_R M \xrightarrow{\sigma} M \otimes_R M^* \xrightarrow{e} R$. $\square$

Exercise 4.2. Prove directly from the definition of a Cauchy module that a retract of a Cauchy module is Cauchy.

Proof. Note that $\rho_L^M : M^* \otimes_R L \rightarrow \text{Hom}_R M, L$ is natural in M , so any module morphism $f : M \rightarrow N : R \nrightarrow S$, so the following “ f -square” commutes:

$$\begin{array}{ccc} N^* \otimes_R L & \xrightarrow{\rho_L^N} & \text{Hom}_R(N, L) \\ f^* \otimes 1 \downarrow & & \downarrow _ \circ f \\ M^* \otimes_R L & \xrightarrow{\rho_L^M} & \text{Hom}_R(M, L) \end{array}$$

Suppose that M is a retract of a Cauchy module N , thus one has $i : M \rightarrow N$, $r : N \rightarrow M$, $r \circ i = 1_M$ and ρ_L^N is invertible. The claim is that the following composite:

$$\text{Hom}(M, L) \xrightarrow{\circ r} \text{Hom}(n, L) \xrightarrow{(\rho_L^N)^{-1}} N^* \otimes L \xrightarrow{i^* \otimes 1} M^* \otimes L$$

is an inverse from ρ_L^M , which is shown the following way:

$$\begin{aligned} \rho_L^M \circ i^* \otimes 1 \circ (\rho_L^N)^{-1} \circ (_ \circ r) &= (_ \circ i) \circ \rho_L^N \circ (\rho_L^N)^{-1} \circ (_ \circ r) \\ &= (_ \circ i) \circ 1_{\text{Hom}(N, L)} \circ (_ \circ r) \\ &= (_ \circ i) \circ (_ \circ r) \\ &= _ \circ ri \\ &= _ \circ 1_M = \\ &= 1_{\text{Hom}(M, L)} \end{aligned}$$

and

$$\begin{aligned} i^* \otimes 1 \circ (\rho_L^N)^{-1} \circ (_ \circ r) \circ \rho_L^M &= (i^* \otimes 1) \circ (\rho_L^N)^{-1} \circ \rho_L^N \circ (r^* \otimes 1) \\ &= (i^* \otimes 1) \circ (r^* \otimes 1) = (ri)^* \otimes 1 = 1_{M^*} \otimes 1_L = 1_{M^* \otimes L}. \end{aligned}$$

$\square$

5 Algebras

Let R be a ring, an *algebra over* R is a module $A : R \nrightarrow R$ along with module morphisms $\mu : A \otimes_R A \rightarrow A$ and $\eta : R \rightarrow A$ such that the following principles

are satisfied (associativity and identity):

$$\begin{array}{ccccc}
A \otimes_R A \otimes_R A & \xrightarrow{\mu \otimes 1_A} & A \otimes_R A & \xrightarrow{\mu} & A \\
& \xrightarrow{1_A \otimes \mu} & & & \\
A & \xrightarrow{\eta \otimes 1_A} & A \otimes_R A & \xrightarrow{\mu} & A \\
& \xrightarrow{1_A \otimes \eta} & & & \\
& & \searrow 1_A & & \nearrow
\end{array}$$

A is also a ring with $ab = \mu(a \otimes b)$ and $1 = \eta(1)$.

Let $A, B : R \nrightarrow R$ be R -algebras, an *algebra morphism* $f : A \rightarrow B$ is a module morphism:

$$\begin{array}{ccc}
A \otimes_R A & \xrightarrow{f \otimes f} & B \otimes_R B \\
\mu \downarrow & & \downarrow \mu \\
A & \xrightarrow{f} & B
\end{array}
\quad
\begin{array}{ccc}
& R & \\
\eta \swarrow & & \searrow \eta \\
A & \xrightarrow{f} & B
\end{array}$$

$\mathbf{Alg}_R(A, B)$ is the set of all algebra morphisms from A to B .

Example 5.1. Let $M : R \nrightarrow S$, the *endomorphism algebras* over S and R are given by:

$$\begin{aligned}
\mathbf{End}_R(M) &= \text{Hom}_R(M, M) : S \nrightarrow S \\
\mathbf{End}^S(M) &= \text{Hom}^S(M, M) : R \nrightarrow R
\end{aligned}$$

where the multiplication is given by composition. A module morphism

$$\hat{\mu} : A \Rightarrow \mathbf{End}^S(M) : R \nrightarrow R$$

corresponds to a module morphism:

$$\mu : A \otimes_R M \Rightarrow M : R \nrightarrow S.$$

Example 5.2. Let $M : R \nrightarrow R$ be a module, then $M^{\otimes n} = M \otimes_R \dots \otimes_R M$ (n times). The *tensor algebra* on M is defined by the geometric series:

$$T(M) = \sum_{n < \omega} M^{\otimes n}$$

with multiplication $\mu : T(M) \otimes T(M) \rightarrow T(M)$ induced by canonical isomorphisms for each $p, q < \omega$:

$$M^{\otimes p} \otimes M^{\otimes q} \xrightarrow{\cong} M^{\otimes(p+q)}$$

and unit $\eta : R \rightarrow T(M)$ defined by the injection:

$$\iota_0 : R \rightarrow \sum_{n < \omega} M^{\otimes n}$$

Composition with the injection $\iota_1 : M \rightarrow T(M)$ induces a bijection:

$$\mathbf{Alg}_R(T(M), A) \cong \text{Hom}_R^R(M, A)$$

for any A . The inverse takes $f : M \rightarrow A$ to $g : T(M) \rightarrow A$ given by $g(m_1 \otimes \dots \otimes m_r) = f(m_1) \dots f(m_r)$. For $M = A$ and $f = 1_A$ we have $\mu : T(A) \rightarrow A$ with $\mu(a_1 \otimes \dots \otimes a_r) = a_1 \dots a_r$.

Example 5.3. Let G be a monoid, an R -algebra $R(G)$ is the free module $\mathcal{F}_R^R(G)$ on the underlying set of G along with the multiplication μ extending G as follows:

$$\begin{array}{ccc} G \times G & \hookrightarrow & \mathcal{F}_R^R(G \times G) \cong \mathcal{F}_R^R(G) \otimes \mathcal{F}_R^R(G) \\ \downarrow \mu & & \downarrow \mu \\ G & \hookrightarrow & \mathcal{F}_R^R(G) \end{array}$$

$R(G)$ is the monoid R -algebra of G , the group R -algebra of G if G is a group. Each monoid morphism $G \rightarrow A$ into the multiplicative monoid of A uniquely extends to an R -algebra morphism $R(G) \rightarrow A$.

A *representation* of G on M is an $R(G)$ -module with scalar multiplication $R(G) \otimes_R M \rightarrow M$ that can be viewed as a monoid morphism $G \rightarrow \mathbf{End}_R(M)$. The subset of M given by $\{gm - m \mid g \in G, m \in M\}$ generates a submodule $(gm - m \mid g \in G, m \in M)$. M/G is the quotient module $M/(gm - m \mid g \in G, m \in M)$.

An ideal in an algebra A is a submodule I such that $axb \in I$ for all $x \in I$ and $a, b \in A$. There is a canonical algebra morphism $\rho : A \rightarrow A/I$. The kernel of any algebra morphism $f : A \rightarrow B$ is an ideal in A .

If X is a subset of A , (X) is the smallest ideal containing X , which is the submodule (AXA) generated by the subset $AXA = \{axb \mid a, b \in A, x \in X\} \subseteq A$. Let $g : A \rightarrow B$ be an algebra morphism $g(x) = 0$ for all $x \in X$, then an algebra morphism $f : A/(X) \rightarrow B$ such that $f \circ \rho = g$.

If R is commutative, then left R -modules and right R -modules are the same thing. Each left R -module can be regarded as a module $M : R \not\rightarrow R$ by defining $rms = (rs)m$ for all $r, s \in R$ and $m \in M$. For R -modules $M_1, \dots, M_n$, we have an R -module $M_1 \otimes_R \dots \otimes_R M_n$.

Every permutation $\xi : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ induces a canonical module isomorphism:

$$\sigma_\xi : M_1 \otimes_R \dots \otimes_R M_n \xrightarrow{\cong} M_{\xi(1)} \otimes_R \dots \otimes_R M_{\xi(n)}$$

An algebra A is commutative if $\mu \circ \sigma = 1_A$.

Example 5.4. Let X be a set and let R be a commutative ring, then the set of all functions R^X is a commutative R -algebra with the operations defined pointwise. The unit $\eta : R \rightarrow R^X$ is given by $\lambda r. \lambda x. r$.

Example 5.5. Let M be a module of the commutative ring R . There is a representation of the symmetric group $\mathcal{S}_n$ on $M^{\otimes n}$ given by $\sigma() : \mathcal{S}_n \rightarrow \mathbf{End}_R(M^{\otimes n})$ such that:

$$\sigma : \xi \mapsto (m_1 \otimes \dots \otimes m_n \mapsto m_{\xi(1)} \otimes \dots \otimes m_{\xi(n)}).$$

The *symmetric R -algebra* on M is induced by *exponential series* of the following form:

$$\mathcal{S}(M) = \sum_{n < \omega} M^{\otimes n} / \mathcal{S}_n.$$

Alternatively, one can construct the symmetric R -algebra the following way. Take any R -algebra A , then we form a commutative R -algebra by factorising A through the ideal $(ab - ba \mid a, b \in A)$. Then we apply this construction to the tensor algebra $T(M)$.

Let A be a commutative R -algebra, then

$$\mathbf{Alg}_R(\mathcal{S}(M), A) \cong \mathbf{Hom}_R(M, A)$$

The skew commutativity principle of the form $ab + ba = 0$ is a too strong requirement for each $a, b \in A$. For example, if $b = 1$, then we have $(1 + 1)a = 0$, then we have $a = 0$ for any field of characteristic $\neq 2$.

An R -algebra is *skew commutative* if for each $a \in A$ we have either $a^2 = 0$ or $a \in \eta(R)$. If $a, b, a + b \notin \mathbf{Im}(R)$, then:

$$ab + ba = (a + b)^2 - a^2 - b^2 = 0.$$

Example 5.6. Let A be an R -algebra, one can form the quotient modulo $(a^2 \mid a \notin \eta(R))$ to obtain a skew commutative algebra. Factorising the tensor algebra $T(M)$ modulo this ideal induces *the exterior algebra* $\Lambda(M)$. Alternatively, let $\Lambda_n(M)$ be the quotient module of $M^{\otimes n}$ by the submodule generated by the elements $m_1 \otimes \dots \otimes m_n$ such that $m_i = m_j$ for some $i \neq j$, then we put

$$\Lambda(M) = \sum_{n < \omega} \Lambda_n(M)$$

$m_1 \wedge \dots \wedge m_n$ stands the image of $m_1 \otimes \dots \otimes m_n$ in $\Lambda(M)$. For each $x, y, z \in M$, one has:

- $x \wedge x = 0$, and, thus $x \wedge y = -y \wedge x$,
- $(rx + sy) \wedge z = r(x \wedge z) + s(y \wedge z)$.

If $M = \mathbf{F}_R(x_1, \dots, x_k)$ is a free module, then $\Lambda_n(M)$ is a free module on a C_n^k , so $\Lambda(M)$ is a free module with 2^n generators. $\Lambda_k(M)$ is free and is generated by $\{x_1 \wedge \dots \wedge x_k\}$. Thus, if

$$y_i = \sum_{j=1}^k r_{ij} x_j$$

then $y_1 \wedge \dots \wedge y_k$ be a unique scalar multiple:

$$y_1 \wedge \dots \wedge y_k = \det(r_{ij}) x_1 \wedge \dots \wedge x_k$$

of $x_1 \wedge \dots \wedge x_k$.

If A is skew commutative, then one has the following bijection:

$$\mathbf{Alg}_R(\Lambda(M), A) \cong \mathbf{Hom}_R(M, A)$$

An *R-Lie algebra* is an R -algebra L along with a module morphism $\beta : L \otimes_R L \rightarrow L$ such that:

- $\beta(x, x) = 0$,
- $\beta(\beta(x, y), z) + \beta(\beta(z, x), y) + \beta(\beta(y, z), x) = 0$ (Jacobi identity).

β is called a *Lie bracket* on the module L .

Example 5.7. Let A be an R -algebra, then the commutator $[a, b] = ab - ba$ defines a Lie bracket on A :

$$\begin{aligned} [[a, b], c] + [[c, a], b] + [[b, c], a] &= \\ [a, b]c - c[a, b] + [c, a]b - b[c, a] + [b, c]a - a[b, c] &= \\ (ab - ba)c - c(ab - ba) + (ca - ac)b - b(ca - ac) + (bc - cb)a - a(bc - cb) &= 0 \end{aligned}$$

A_L is the Lie subalgebra of A , that is, subalgebra of those elements that are closed under commutator.

Example 5.8. Let A be an R -algebra and let $M : A \not\rightarrow A$ be a module. Then a derivation $D : A \rightarrow M$ is an R -module morphism such that:

$$D(ab) = D(a)b + aD(b)$$

$\mathbf{Der}_R(A, M)$ is a submodule of $\mathbf{Hom}_R(A, M)$. $\mathbf{Der}_R(A, M)$ is closed under commutator in $\mathbf{End}_R(A)$ by putting

$$[D_1, D_2] = D_1 \circ D_2 - D_2 \circ D_1$$

Let L and L' be R -Lie algebras and $f : L \rightarrow L'$ be an R -module morphism, then f is a Lie algebra morphism if:

$$f(\beta(x, y)) = \beta(f(x), f(y))$$

Let L be an R -Lie-algebra, the *universal enveloping algebra* of L is an R -algebra $\mathcal{U}(L)$ such that there is a natural isomorphism:

$$\mathbf{Alg}_R(\mathcal{U}(L), A) \cong \mathbf{Lie}(L, A_L)$$

We construct $\mathcal{U}(L)$ by factorising the tensor algebra modulo the appropriate ideal:

$$\mathcal{U}(L) := T(L) / (x \otimes y - y \otimes x - \beta(x, y) \mid x, y \in L).$$

The *direct sum* $L_1 \oplus L_2$ of Lie algebras L_1 and L_2 is their direct sum of modules along with the Lie bracket:

$$\beta((x_1, x_2), (y_1, y_2)) = (\beta(x_1, y_1), \beta(x_2, y_2))$$

Theorem 5.1. There is an algebra isomorphism:

$$\mathcal{U}(L_1 \oplus L_2) \cong \mathcal{U}(L_1) \otimes \mathcal{U}(L_2)$$

whose composite $i : L_1 \oplus L_2 \rightarrow \mathcal{U}(L_1 \oplus L_2)$ takes the pair (x_1, x_2) to $x_1 \otimes 1 + 1 \otimes x_2$.

Proof. First, the following maps are Lie algebra morphisms:

- $L_1 \oplus L_2 \rightarrow (\mathcal{U}(L_1) \otimes_R \mathcal{U}(L_2))_L, (x_1, x_2) \mapsto x_1 \otimes 1 + 1 \otimes x_2$;
- $L_1 \rightarrow (\mathcal{U}(L_1) \otimes_R \mathcal{U}(L_2))_L$ with $x_1 \mapsto (x_1, 0)$;
- $L_2 \rightarrow (\mathcal{U}(L_1) \otimes_R \mathcal{U}(L_2))_L$ with $x_2 \mapsto (0, x_2)$.

We have the following algebra morphisms:

$$\begin{aligned} \phi : \mathcal{U}(L_1 \oplus L_2) &\rightarrow \mathcal{U}(L_1) \otimes_R \mathcal{U}(L_2) \\ \phi_1 : \mathcal{U}(L_1) &\rightarrow \mathcal{U}(L_1 \oplus L_2) \\ \phi_2 : \mathcal{U}(L_2) &\rightarrow \mathcal{U}(L_1 \oplus L_2) \end{aligned}$$

Define $\psi : \mathcal{U}(L_1) \otimes_R \mathcal{U}(L_2) \rightarrow \mathcal{U}(L_1 \oplus L_2)$ by $\psi(a \otimes b) = \psi_1(a)\psi_2(b)$. Then we have:

$$\begin{aligned} \psi(\phi(x_1, x_2)) &= \psi(x_1 \otimes 1 + 1 \otimes x_2) = (x_1, 0) + (0, x_2) = (x_1, x_2) \\ \phi(\psi_1(x_1)) &= \phi(x_1, 0) = x_1 \otimes 1 \\ \phi(\psi_2(x_2)) &= \phi(0, x_2) = 1 \otimes x_2 \end{aligned}$$

□

5.1 Exercises

Exercise 5.1. Let R be a commutative ring and let G be a group. Consider left modules M, N, L over the group algebra $R(G)$, then the following holds:

1. $M \otimes_R N$ is an $R(G)$ -module on defining for $g \in G, m \in M, n \in N$:

$$g(m \otimes n) = (gm) \otimes (gn)$$

2. $\text{Hom}_R(M, L)$ is an $R(G)$ -module on defining for $g \in G, u \in \text{Hom}_R(M, L), m \in M$:

$$(gu)(n) = gu(g^{-1}m)$$

3. The evaluation $ev_M : M \otimes_R \text{Hom}_R(M, L) \rightarrow L$ is an $R(G)$ -module morphism.
4. The following isomorphism holds:

$$\text{Hom}_{R(G)}(N, \text{Hom}_R(M, L)) \cong \text{Hom}_{R(G)}(M \otimes_R N, L).$$

Proof. 1. We have that $G \rightarrow \text{End}_R(M \otimes N)$ with $g \mapsto (m \otimes n \mapsto (gm) \otimes (gn))$ is a monoid morphism since $1 \mapsto 1_{M \otimes N}$ and

$$gh \mapsto (m \otimes n \mapsto (hm) \otimes (hn) \mapsto (ghm) \otimes (ghn)).$$

This extends to a unique R -algebra morphism $R(G) \rightarrow \text{End}_R(M \otimes_R N)$, so $M \otimes_R N$ is an $R(G)$ -module.

2. Let $\hat{\mu}(g) : \text{Hom}_R(M, L) \rightarrow \text{Hom}_R(M, L)$ be the R -module morphism given by $\hat{\mu}(g)(\mu)(m) = gh\mu(h^{-1}g^{-1}m)$. So we have

$$\hat{\mu}(gh)(\mu)(m) = gh\mu(h^{-1}g^{-1}m) = g\hat{\mu}(h)(\mu)(g^{-1}m) = (\hat{\mu}(g) \circ \hat{\mu}(h))(\mu)(m)$$

and $\hat{\mu}(1)(\mu)(m) = \mu(m)$. Thus $\hat{\mu} : G \rightarrow \text{End}_R(\text{Hom}_R(M, L))$ is a monoid morphism, so $\text{Hom}_R(M, L)$ is an $R(G)$ -module.

3. The proof is the following:

$$ev_M(g \circ (m \otimes \mu)) = ev_M(gm \otimes g\mu) = (g\mu)(gm) = g\mu(g^{-1}gm) = g\mu(m) = gev_M(m \otimes \mu).$$

4. We need $d : N \rightarrow \text{Hom}_R(M, M \otimes_R N)$ with $n \mapsto (m \mapsto m \otimes n)$ to be an $R(G)$ -module morphism. We have:

$$\begin{aligned} d(gn)(m) &= m \otimes gn \\ &= g((g^{-1}m) \otimes n) \\ &= gd(n)(g^{-1}m) = (gd(n))(m) \end{aligned}$$

The required isomorphism is the restriction of

- $\text{Hom}_R(N, \text{Hom}_R(M, L)) \cong \text{Hom}_R(M \otimes_R N, L)$,
- $f \mapsto e \circ (1_M \otimes f)$,
- $(g \circ _) \circ \leftarrow g$.

□

6 Coalgebras and bialgebras

Let R be a ring. A *coalgebra* over R is a module $C : R \nrightarrow R$ along with module morphisms $\delta : C \rightarrow C \otimes_R C$ and $\varepsilon : C \rightarrow R$ such that:

$$\begin{array}{ccc} C & \xrightarrow{\delta} & C \otimes_R C \xrightarrow[1 \otimes \delta]{\delta \otimes 1} C \otimes_R C \otimes_R C \\ & & \\ C & \xrightarrow{\delta} & C \otimes_R C \xrightarrow[1 \otimes \varepsilon]{\varepsilon \otimes 1} C \end{array}$$

There is a uniquely determined morphism $\delta : C \rightarrow C \otimes_R C \otimes_R \dots \otimes_R C$ defined as:

$$\delta : c \mapsto \sum_i c_{1i} \otimes \dots \otimes c_{ni}$$

We will be using the following notation:

$$\delta(c) = \sum (c)c_{(1)} \otimes \dots \otimes c_{(n)}$$

Given a multilinear function $f : C \times C \dots \times C \rightarrow A$, we write:

$$f(\delta(c)) = \sum_{(c)} f(c_1, \dots, c_{(n)}).$$

So we can rewrite the axioms the following way:

$$\begin{aligned} \sum_{(c)} \delta(c_{(1)}) \otimes c_{(2)} &= \sum_{(c)} c_{(1)} \otimes c_{(2)} \otimes c_{(3)} = \sum_{(c)} c_{(1)} \otimes \delta(c_{(2)}) \\ c &= \sum_{(c)} \varepsilon(c_{(1)}) \otimes c_{(2)} = \sum_{(c)} c_{(1)} \otimes \varepsilon(c_{(2)}) \end{aligned}$$

A *coalgebra morphism* $f : C \rightarrow D$ is a module morphism such that:

$$\begin{array}{ccc} C & \xrightarrow{f} & D \\ \delta \downarrow & & \downarrow \delta \\ C \otimes_R C & \xrightarrow{f \otimes f} & D \otimes_R D \end{array} \quad \begin{array}{ccc} C & \xrightarrow{f} & D \\ \varepsilon \searrow & & \swarrow \varepsilon \\ & R & \end{array}$$

that is for $c, c_1, c_2 \in C$,

$$\begin{aligned} \sum_{(c)} f(c_{(1)}) \otimes f(c_{(2)}) &= \sum_{f(c)} f(c)_{(1)} \otimes f(c)_{(2)} \\ \varepsilon(f(c)) &= \varepsilon(c). \end{aligned}$$

$\mathbf{Cog}_R(C, D)$ is the set of all coalgebra morphisms from C to D .

Let R be commutative, then a coalgebra C over R is commutative whenever:

$$\begin{array}{ccc} & C & \\ \delta \swarrow & & \searrow \delta \\ C \otimes_R C & \xrightarrow{\sigma} & C \otimes_R C \end{array}$$

Let R be any ring, let C be R -coalgebra and let A be an R -algebra. Then $\text{Hom}_R(C, A)$ is an R -algebra under the *convolution structure*:

$$\begin{array}{ccc} \text{Hom}_R(C, A) \otimes_R \text{Hom}_R(C, A) & & R \\ \downarrow - \otimes - & & \downarrow \cong \\ \text{Hom}_R(C \otimes_R C, A \otimes_R A) & & \text{Hom}_R(R, R) \\ \downarrow \mu \circ \delta & & \downarrow \eta \circ \varepsilon \\ \text{Hom}_R(C, A) & & \text{Hom}_R(C, A) \end{array}$$

In other words, let $f, g : C \rightarrow A$, the convolution product $f * g$ is given by:

$$\begin{array}{ccc}
 C & & C \\
 \delta \downarrow & & \downarrow \varepsilon \\
 C \otimes C & & R \\
 f \otimes g \downarrow & & \downarrow \eta \\
 A \otimes A & & A \\
 \mu \downarrow & & \\
 A & &
 \end{array}$$

In other words, $f * g$ is given by the following formula:

$$(f * g)(c) = \sum_{(c)} f(c_{(1)})g(c_{(2)})$$

If $A = R$, then each R -coalgebra C induces a convolution R -algebra structure on the dual $C^* = \text{Hom}_R(C, R)$. One can also view C^* as an R -algebra with the multiplication $\mu : C^* \otimes_R C^* \rightarrow C^*$:

$$\begin{array}{ccc}
 C^* \otimes_R C^* \otimes_R C & \xrightarrow{\mu \otimes 1} & C^* \otimes_R C \\
 1 \otimes 1 \otimes \delta \downarrow & & \downarrow e \\
 C^* \otimes_R C^* \otimes_R C \otimes_R C & \xrightarrow{1 \otimes e \otimes 1} C^* \otimes_R C \xrightarrow{e} & R
 \end{array}$$

Example 6.1. Let $\mathcal{C}$ be any category with finite products. Let $F : \mathcal{C} \rightarrow \mathbf{Mod}_R^R$ is an oplax monoidal functor into the category of modules from R to R , that is, there is a family of natural morphisms:

$$\phi_{A,B} : F(A \times B) \rightarrow FA \otimes FB$$

compatible with the associator. Then FA for each $A \in \mathcal{C}$ is endowed with the structure of a coalgebra with:

$$\begin{array}{ccc}
 FA & \xrightarrow{F\delta} F(A \times A) & \xrightarrow{\varphi_{A,A}} FA \otimes_R FA \\
 & & \downarrow F\varepsilon \\
 & FX & \xrightarrow{F\varepsilon} F\mathbb{1} \xrightarrow{\varphi_{\mathbb{1}}} R.
 \end{array}$$

Example 6.2. The free R -module construction given a functor $\mathcal{F}_R : \mathbf{Set} \rightarrow \mathbf{Mod}_R$. We have natural isomorphisms:

$$\begin{aligned}
 \phi : \mathcal{F}_R(A \times B) &\xrightarrow{\cong} \mathcal{F}_R A \otimes_R \mathcal{F}_R B \\
 \phi : (a, b) &\mapsto a \otimes b
 \end{aligned}$$

The proof:

$$\begin{aligned}
\mathrm{Hom}_R(\mathcal{F}_R(A \times B), M) &\cong M^{A \times B} \cong (M^A)^B \cong \\
&\mathrm{Hom}_R(\mathcal{F}_R B, M^A) \cong \\
&\mathrm{Hom}_R(\mathcal{F}_R B, \mathrm{Hom}_R(\mathcal{F}_R A, M)) \cong \\
&\mathrm{Hom}_R(\mathcal{F}_R A \otimes \mathcal{F}_R B, M)
\end{aligned}$$

Example 6.3. The universal enveloping algebra provides a functor $\mathcal{U} : \mathbf{Lie}_R \rightarrow \mathbf{Mod}_R$ such that $\mathcal{U}(\{0\}) \cong R$ and

$$\begin{aligned}
\mathcal{U}(L_1 \oplus L_2) &\cong \mathcal{U}L_1 \otimes_R \mathcal{U}L_2 \\
(x, x') &\mapsto x \otimes 1 + 1 \otimes x'.
\end{aligned}$$

Each universal enveloping algebra $\mathcal{U}L$ is an R -coalgebra, the comultiplication here is given by:

$$\begin{aligned}
L &\xrightarrow{i} \mathcal{U}L \xrightarrow{\delta} \mathcal{U}L \otimes_R \mathcal{U}L \\
x &\mapsto x \otimes 1 + 1 \otimes x.
\end{aligned}$$

We summarise the above examples as follows:

- $c \in C$ is *set-like* when $\delta(c) = c \otimes c$ and $\varepsilon(c) = 1$. $\mathcal{D}(C)$ is the set of set-like elements of C .
- $c \in C$ is *primitive* when $\delta(c) = c \otimes 1 + 1 \otimes c$. $\mathcal{P}(C)$ is the submodule of primitive elements of C .

Proposition 6.1. Let R be a field, then the set-like elements of any coalgebra C form a linearly independent subset $\mathcal{D}(C)$.

Proof. Assume $\mathcal{D}(C)$ is linearly dependent. Let $n+1$ be the first natural number for which there is a linearly dependent subset of $\mathcal{D}(C)$ with that number of elements. Then any set of n elements of $\mathcal{D}(C)$ is linearly independent, whilst there exist distinct $g, g_1, \dots, g_n \in \mathcal{D}(C)$. Then

$$g = \lambda_1 g_1 + \dots + \lambda_n g_n$$

where each $\lambda_i \in R$, $1 \leq i \leq n$, is non-zero. Therefore:

$$\sum_{i=1}^n \lambda_i g_i \otimes g_i = \sum_{i=1}^n \lambda_i \delta(g_i) = \delta(g) = g \otimes g = \sum_{i,j=1}^n \lambda_i \lambda_j g_i \otimes g_j.$$

$\{g_1, \dots, g_n\}$ is linearly independent in C , so $\{g_i \otimes g_j\}$ is linearly independent in $C \otimes C$, so we can equate $\lambda_i \lambda_j = 0$ for $i \neq j$ and $\lambda_i = \lambda_i^2$. As far as $\lambda_i \neq 0$, so $n = 1$ and $\lambda_i = 1$. And also $g = g_1$, contradiction. $\square$

Example 6.4. Let R be commutative. Let $C = \mathcal{F}_R \omega$ be the free R -module on the countable set ω . Define

$$\delta(n) = \sum_{p+q=n} p \otimes q$$

and

$$\varepsilon(n) = \begin{cases} 1, & \text{for } n = 0, \\ 0, & \text{for } n > 0. \end{cases}$$

Then (C, δ, ε) is a cocommutative coalgebra on C . The convolution structure on $\text{Hom}_R(C, A)$ induces the following isomorphism:

$$\text{Hom}_R(C, A) \cong A^\omega$$

with the multiplication defined as:

$$ab = \left(\sum_{p+1=n} a_p b_q \right)$$

for sequences $a = (a_n) = (a_0, a_1, \dots)$ and $b = (b_n) = (b_0, b_1, \dots)$ in A . The unit sequence is $(1, 0, 0, 0, \dots)$. An *indeterminate* is taken to mean the following sequence:

$$x = (0, 1, 0, 0, 0, \dots) \in A^\omega$$

Each $u \in A$ is identified with $u1 = (u, 0, 0, 0, \dots) \in A^\omega$, then each $a \in A^\omega$ can be thought as a formal power series of the form

$$a = \sum_{n=0}^{\infty} a_n x^n$$

Let $A[[x]]$ denote A^ω with the above algebra structure, so this is the R -algebra of formal power series in A . If A is commutative, so is $A[[x]]$. If $A = R$, then we obtain R -coalgebra $C^* = R[[x]]$.

Example 6.5. Let $\mathbf{n} = \{1, 2, \dots, n\}$ and let C denote $\mathcal{F}_R(\mathbf{n} \times \mathbf{n})$, then C is an R -coalgebra by defining:

$$\delta(i, j) = \sum_{k=1}^n (i, k) \otimes (k, j)$$

and

$$\varepsilon(i, j) = \begin{cases} 1, & \text{for } i = j, \\ 0, & \text{otherwise.} \end{cases}$$

Let A be an R -coalgebra, then we have the following R -module isomorphism:

$$\text{Hom}_R(C, A) \cong A^{\mathbf{n} \times \mathbf{n}}$$

with the convolution product on $\text{Hom}_R(C, A)$ and the matrix multiplication on $A^{\mathbf{n} \times \mathbf{n}}$:

$$(a_{ij})(b_{ij}) = \left(\sum_{k=1}^n a_{ik} b_{kj} \right).$$

We obtain the R -algebra $\mathbf{Mat}(n, A) \cong \text{End}_R(A^n)$ of $n \times n$ with entries in A .

Let R be a commutative ring, an R -bialgebra is an R -module B along with algebra and coalgebra structures:

- $\mu : B \otimes_R B \rightarrow B$,
- $\delta : B \rightarrow B \otimes_R B$,
- $\eta : R \rightarrow B$,
- $\varepsilon : B \rightarrow R$

such that the following diagram commute:

$$\begin{array}{ccc}
 B \otimes_R B & \xrightarrow{\mu} & B \\
 \delta \otimes \delta \downarrow & & \downarrow \delta \\
 B \otimes_R B \otimes_R B \otimes_R B & \xrightarrow{1 \otimes \sigma \otimes 1} B \otimes_R B \otimes_R B \otimes_R B \xrightarrow{\mu \otimes \mu} B \otimes_R B & \\
 \\
 \begin{array}{ccccc}
 B \otimes B & \xrightarrow{\varepsilon \otimes \varepsilon} R \otimes R & \xrightarrow{\eta \otimes \eta} B \otimes B & & B \\
 \mu \downarrow & & \downarrow \cong & \uparrow \delta & \nearrow \eta \\
 B & \xrightarrow{\varepsilon} R & \xrightarrow{\eta} B & & R \\
 & & & & \nwarrow \varepsilon \\
 & & & & R
 \end{array}
 \end{array}$$

Let R be a commutative ring, the tensor product $A \otimes_R A'$ of R -algebras A and A' is an R -algebra with the following multiplication and the unit:

$$(A \otimes_R A') \otimes_R (A \otimes_R A') \xrightarrow{\cong} (A \otimes_R A) \otimes_R (A' \otimes_R A') \xrightarrow{\mu \times \mu} A \otimes_R A'$$

$$R \xrightarrow{\cong} R \otimes_R R \xrightarrow{\eta \otimes \eta} A \otimes_R A'$$

The tensor product $C \otimes_R C'$ of R -coalgebras C and C' is an R -coalgebra with the following comultiplication and the counit:

$$C \otimes_R C' \xrightarrow{\delta \otimes \delta} (C \otimes_R C) \otimes_R (C' \otimes_R C') \xrightarrow{\cong} (C \otimes_R C') \otimes_R (C \otimes_R C')$$

$$C \otimes_R C' \xrightarrow{\varepsilon \otimes \varepsilon} R \otimes_R R \cong R$$

Proposition 6.2. Suppose (μ, η) and (δ, ε) are algebra and coalgebra structures on the R -module B , then TFAE:

1. B is bialgebra,
2. $\mu : B \otimes_R B \rightarrow B$ and $\eta : R \rightarrow B$ are coalgebra morphisms,
3. $\delta : B \rightarrow B \otimes_R B$ and $\varepsilon : B \rightarrow R$ are algebra morphisms.

$\mathbf{Big}_R(B, B')$ is the set of all bialgebra morphisms from B to B' .

Proposition 6.3. If B is a bialgebra then the set-like elements are closed under multiplication: so $\mathcal{D}(B)$ is a monoid.

Proof. Take $b, b' \in \mathcal{D}(B)$:

$$\delta(bb') = \delta(b)\delta(b') = (b \otimes b)(b' \otimes b') = bb' \otimes bb'.$$

□

Proposition 6.4. Let B be a bialgebra, then the set of primitive elements is closed under commutator, so $\mathcal{P}(B)$ is a Lie algebra. Moreover, $\varepsilon(x) = 0$ for all $x \in \mathcal{P}(B)$.

Proof. Let $x, y \in \mathcal{P}(B)$, then:

$$\begin{aligned} \delta([x, y]) &= \delta(x)\delta(y) - \delta(y)\delta(x) \\ &= (x \otimes 1 + 1 \otimes x)(y \otimes 1 + 1 \otimes y) - (y \otimes 1 + 1 \otimes y)(x \otimes 1 + 1 \otimes x) \\ &= (xy \otimes 1 + x \otimes y + y \otimes x + 1 \otimes xy) - (yx \otimes 1 + y \otimes x + x \otimes y + 1 \otimes yx) \\ &= [x, y] \otimes 1 + 1 \otimes [x, y] \end{aligned}$$

And thus $[x, y] \in \mathcal{P}(B)$. Also

$$x = (1 \otimes \varepsilon)\delta(x) = (1 \otimes \varepsilon)(x \otimes 1 + 1 \otimes x) = x + \varepsilon(x)$$

and therefore $\varepsilon(x) = 0$.

□

Example 6.6. There are two conditions on the functor $F : \mathcal{X} \rightarrow \mathbf{Mod}_R$ rising to bialgebras FX .

1. When the morphisms ϕ are invertible. The multiplication and unit for G are given as follows:

- $FG \otimes_R FG \cong F(G \times G) \xrightarrow{F\mu} FG,$
- $R \cong F1 \xrightarrow{F\eta} FG$

on FG . These are coalgebra morphisms since all arrows in $\mathcal{X}$ “commute with diagonals”. Thus FG is a bialgebra. This is the case for the functor $\mathcal{F}_R : \mathbf{Set} \rightarrow \mathbf{Mod}_R$, so for each monoid G the monoid algebra $R(G)$ is a cocommutative bialgebra.

2. When F lifts to $F : \mathcal{X} \rightarrow \mathbf{Alg}_R$. Each FX is a bialgebra since the comultiplication and counit are algebra morphisms.

Example 6.7. Recall the example with the free module $\mathcal{F}_R\omega$:

$$\delta(n) = \sum_{p+q=n} p \otimes q$$

and

$$\varepsilon(n) = \begin{cases} 0, & \text{if } n > 0, \\ 1, & \text{if } n = 0. \end{cases}$$

Now consider the following structure $\mathcal{F}_R(E)$ for $E = \{e_i \mid i < \omega\}$:

$$e_p e_q = \frac{(p+q)!}{p!q!} e_{p+q}$$

and

$$e_0 = 1$$

Then $\mathcal{F}_R(E)$ is a bialgebra, if $\text{char } R = 0$, we put $x = e_1$, so one has:

$$e_n = \frac{1}{n!} x^n.$$

Thus $\mathcal{F}_R(E)$ is isomorphic to $R[x]$. We think of $\mathcal{F}_R(E)$ as the algebra of Hurwitz polynomials in one indeterminate:

$$\sum_{n=0}^k \frac{a_n x^n}{n!}$$

for each $a_n \in R$.

6.1 Exercises

Exercise 6.1. For any R -coalgebra C , prove the following identities:

1. $\delta(c) = \Sigma_{(c)} \varepsilon(c_{(2)}) \otimes \delta(c_{(1)}) = \Sigma_{(c)} \delta(c_{(2)}) \otimes \varepsilon(c_{(1)}) \Sigma_{(c)} c_{(1)} \otimes \varepsilon(c_{(3)}) \otimes c_{(2)}$
2. $\Sigma_{(c)} \varepsilon(c_{(1)}) \otimes c_{(3)} \otimes c_{(2)} = \Sigma_{(c)} c_{(2)} \otimes c_{(1)}$
3. $\Sigma_{(c)} \varepsilon(c_{(1)}) \otimes c_{(3)} \otimes c_{(2)} = c.$

Proof.

1. $\delta(c) = \Sigma_{(c)} \varepsilon(c_{(2)}) \otimes \delta(c_{(1)}) = \Sigma_{(c)} \delta(c_{(2)}) \otimes \varepsilon(c_{(1)}) \Sigma_{(c)} c_{(1)} \otimes \varepsilon(c_{(3)}) \otimes c_{(2)}$ is proved the following way:

$$\begin{array}{ccccccc}
& & C \otimes C & \xrightarrow{\sigma} & C \otimes C & \xrightarrow{\varepsilon \otimes 1} & R \otimes C \\
& \nearrow \delta & \downarrow 1 \otimes \varepsilon & & \nearrow \sigma & & \downarrow 1 \otimes \delta \\
C & \xrightarrow{\cong} & C \otimes R & \xrightarrow{\delta \otimes 1} & C \otimes C \otimes R & \xrightarrow{\cong} & R \otimes C \otimes C \\
\downarrow \delta & & & & & & \parallel \\
C \otimes C & \xrightarrow{\cong} & & & & & R \otimes C \otimes C
\end{array}$$

2. $\Sigma_{(c)}\varepsilon(c_{(1)}) \otimes c_{(3)} \otimes c_{(2)} = \Sigma_{(c)}c_{(2)} \otimes c_{(1)}$ holds as the following diagram commutes:

$$\begin{array}{ccccccc}
& & C \otimes C \otimes C & \xrightarrow{1 \otimes \delta} & C \otimes C \otimes C & & \\
& \nearrow \delta & \uparrow \delta \otimes 1 & \searrow \varepsilon \otimes 1 \otimes 1 & \searrow \varepsilon \otimes 1 \otimes 1 & & \\
C & \xrightarrow{\delta} & C \otimes C & \xrightarrow{\cong} & R \otimes C \otimes C & \xrightarrow{1 \otimes \sigma} & R \otimes C \otimes C \\
& & \searrow \sigma & & \nearrow \cong & & \\
& & C \otimes C & & & &
\end{array}$$

3. $\Sigma_{(c)}\varepsilon(c_{(1)}) \otimes c_{(3)} \otimes c_{(2)} = c$ holds because of:

$$\begin{array}{ccccccc}
& & C \otimes C \otimes C & \xrightarrow{1 \otimes \sigma} & C \otimes C \otimes C & & \\
& \nearrow \delta & \uparrow \delta \otimes 1 & \downarrow \varepsilon \otimes 1 \otimes 1 & \downarrow \varepsilon \otimes 1 \otimes 1 & \searrow \varepsilon \otimes \varepsilon \otimes 1 & \\
C & \xrightarrow{\delta} & C \otimes C & \xrightarrow{\cong} & R \otimes C \otimes C & \xrightarrow{1 \otimes \sigma} & R \otimes C \otimes C \\
& & \searrow 1 \otimes \varepsilon & & \searrow 1 \otimes 1 \otimes \varepsilon & & \searrow 1 \otimes \sigma \\
& & C \otimes R & \xrightarrow{\cong} & R \otimes C \otimes R & &
\end{array}$$

□

7 Dual Algebras of Coalgebras

The dual C^* of a coalgebra C has a natural structure of a coalgebra. However, if we take the dual A^* of an algebra A , then A^* does not have to be a coalgebra since the canonical morphism

$$M^* \otimes_R N^* \rightarrow (M \otimes_R N)^*$$

is not invertible. If M is Cauchy, the morphism is invertible since:

$$(M \otimes_R N)^* = \text{Hom}_R(M \otimes_R N, R) \cong \text{Hom}_R(M, N^*) \cong M^* \otimes_R N^*$$

So if an algebra A is Cauchy as a module, then we have the following structure:

- $\delta : A^* \xrightarrow{\mu^*} (A \otimes_R A)^* \cong A^* \otimes_R A^* \cong A^*$,
- $\varepsilon : A^* \xrightarrow{\eta^*} R^* \cong R$.

Let us modify the notion of a dual algebra instead of restricting it.

An ideal I of an algebra A is called *coCauchy* when the quotient algebra A/I is Cauchy. Define:

$$A^0 = \{u \in A^* \mid u \text{ is zero on some coCauchy ideal of } A\}.$$

Proposition 7.1. Let R be a field,

1. A^0 is a submodule of A^* .
2. If $f \in \mathbf{Alg}_R(A, B)$, then $f^* : B^* \rightarrow A^*$ takes B^0 to A^0 .
3. For any R -algebra B the canonical morphism $A^* \otimes B^* \rightarrow (A \otimes B)^*$ induces an isomorphism $A^0 \otimes B^0 \xrightarrow{\cong} (A \otimes B)^0$.
4. There is a unique $\delta : A^0 \rightarrow A^0 \otimes A^0$ such that the following diagram commutes:

$$\begin{array}{ccccc} A^0 & \xrightarrow{\delta} & A^0 \otimes A^0 & \xrightarrow{\cong} & (A \otimes A)^0 \\ \downarrow & & & & \downarrow \\ A^* & \xrightarrow{\mu^*} & & & (A \otimes A)^* \end{array}$$

Proof. 1. If $u \in A^0$ and $r \in R$, then $\ker u \subseteq \ker(ru)$, then $ru \in A^0$. Take $u, v \in A^0$ zero on coCauchy ideals I and J respectively. One can find subspaces U, V, W of A such that:

$$A = (I \cap J) \oplus U \oplus V \oplus W$$

and $I = (I \cap J) \oplus U$ and $J = (I \cap J) \oplus V$. Thus $A/I \cong V \oplus W$ and $A/J \cong U \oplus W$ are finite dimensional. Therefore $A/I \cap J \cong U \oplus V \oplus W$ is finite dimensional. Therefore $I \cap J$ is coCauchy on which $u + v$ is zero.

2. Take $v \in B^0$ zero on coCauchy J in B . Then $f^{-1}J \subseteq \ker(vf) = \ker f^*(v)$ is an ideal of A . $f^{-1}(J)$ is the kernel of $A \xrightarrow{f} B \rightarrow B/J$, so $A/f^{-1}(J)$ is isomorphic to a subspace B/J . Thus $f^{-1}(J)$ is coCauchy.
3. Note that for any coCauchy ideal K in $A \otimes B$, one has the following ideals of A , B and $A \otimes B$ respectively:

$$\begin{aligned} I &= \{a \in A \mid a \otimes 1 \in K\} = (A \xrightarrow{-\otimes 1} A \otimes B)^*(K) \\ J &= \{b \in B \mid 1 \otimes b \in K\} = (B \xrightarrow{1 \otimes -} A \otimes B)^*(K) \\ A \otimes_R J + I \otimes_R B &= \ker(A \otimes_R B \rightarrow (A/I) \otimes_R (B/J)). \end{aligned}$$

Take $w \in (A \otimes B)^0$ which is zero on some coCauchy K . Then w is zero on $A \otimes J + I \otimes B \subseteq K$. Therefore:

$$A \otimes B / (A \otimes J + I \otimes B) \cong A/I \otimes B/J$$

so there is a unique $\bar{w}$:

$$\begin{array}{ccc} A \otimes B & \xrightarrow{w} & R \\ & \searrow \rho \otimes \rho \quad \nearrow \bar{w} & \\ & (A/I) \otimes (B/J) & \end{array}$$

Since A/I and B/J are finite dimensional, therefore we have:

$$(A/I)^* \otimes (B/J)^* \xrightarrow{\cong} (A/I \otimes B/J)^*$$

and thus there is some $\Sigma \bar{h}_i \otimes \bar{k}_i$ corresponding to $\bar{w}$. For $x \in A/I$ and $y \in B/J$ we have

$$\bar{w}(x \otimes y) = \sum_i \bar{h}_i(x) \bar{k}_i(y).$$

for $x \in A/I$ and $y \in B/J$. Define the composite $h_i = \bar{h}_i \circ \rho : A \rightarrow A/I \rightarrow R$ and similarly define $k_i = \bar{k}_i \circ \rho : B \rightarrow B/J \rightarrow R$. These are in A^0 and B^0 respectively since they are zero on I and J . Thus:

$$\sum_i h_i \otimes k_i \in A^0 \otimes B^0.$$

Conversely, if $h \in A^0$ and $k \in B^0$ vanish on coCauchy I and J (ideals of A and B), then $h \otimes k$ vanishes $A \otimes J + I \otimes B$, which is a coCauchy ideal of $A \otimes B$.

4. Suppose $u \in A^0$ vanishes on a coCauchy ideal I . Then

$$\mu^*(u)(a \otimes b) = (u\mu)(a \otimes b) = u(ab),$$

so $\mu^*(u)$ vanishes on $A \otimes I + I \otimes A$ which is a coCauchy ideal of $A \otimes A$. So μ^* takes A^0 into $(A \otimes A)^0$ and δ does exist. $\square$

Corollary 7.1. Let R be a field. For each algebra A a coalgebra structure on A^0 is given by the δ in the previous statement and $\varepsilon = (A^0 \hookrightarrow A^* \xrightarrow{\eta^*} R^* \cong R)$. Each algebra morphism $f : A \rightarrow B$ induces a coalgebra morphism $f^0 : B^0 \rightarrow A^0$ given by restriction of f^* .

For each algebra A , one obtains a left and right A -module structure A^* given the following way for $a \in A$ and $u \in A^*$:

- $(au)(x) = u(xa),$
- $(ua)(x) = u(ax).$

For any $f \in A^*$ we write Af , fA and AfA for the R -submodules of A^* consisting of those elements of the form af , fa and afb respectively with $a, b \in A$.

Proposition 7.2. Let R be a field and $f \in A^*$, then the following are equivalent:

1. $f \in A^0$,
2. $\mu^*(f)$ is in the image of $A^* \otimes A^* \rightarrow (A \otimes A)^*$,
3. $\mu^*(f)$ is in the image of $(A \otimes A)^0 \hookrightarrow (A \otimes A)^*$;
4. Af is Cauchy,
5. fA is Cauchy,
6. AfA is Cauchy.

Corollary 7.2. For any coalgebra C the canonical injection $d : C \rightarrow C^{**}$ given by $d(c)(u) = u(c)$ has image in $(C^*)^0$.

Proof. Take $c \in C$, then $C^*d(c) = \{ud(c) \mid u \in C^*\} \subseteq C^{**}$. Therefore:

$$(ud(c))(v) = d(c)(v * u) = (v \otimes u)\delta(c) = (v \otimes u) \sum_{(c)} c_{(1)} \otimes c_{(2)} = \sum_{(c)} v(c_{(1)})u(c_{(2)})$$

by multiplication $v * u$ in C . Thus

$$ud(c) = \sum_{(c)} u(c_{(2)})d(c_{(1)})$$

which is in the subspace spanned by $d(c_{(1)})$. Thus $C^*d(c)$ has finite dimension, so $d(c) \in (C^*)^0$. $\square$

Theorem 7.1. Let R be a field, then there is the following bijection between algebras A and coalgebras C :

- $\mathbf{Alg}_R(A, C^*) \cong \mathbf{Cog}_R(C, A^0)$,
- $(A \xrightarrow{f} C) \mapsto (C \xrightarrow{d} (C^*)^0 \xrightarrow{f^0} A^0)$.

Proof. The inclusion $i : A^0 \rightarrow A^*$ induces $i^* : A^{**} \rightarrow A^{0*}$, while the inverse to $f \mapsto f^0 \circ d$ takes $g \in \mathbf{Cog}_R(C, A^0)$ to the composite:

$$A \xrightarrow{d} A^{**} \xrightarrow{i^*} A^{0*} \xrightarrow{g^*} C^*.$$

$\square$

8 Hopf algebras

An R -Hopf algebra is an R -bialgebra equipped with an R -module morphism $\nu : H \rightarrow H$ called *the antipode* satisfying the following diagram:

$$\begin{array}{ccccc} H & \xrightarrow{\delta} & H \otimes H & \xrightleftharpoons[1 \otimes \nu]{\nu \otimes 1} & H \otimes H & \xrightarrow{\mu} & H \\ & \searrow \varepsilon & & & \nearrow \eta & & \\ & & R & & & & \end{array}$$

For any Hopf algebra H , H^{op} denotes the Hopf algebra obtained by replacing μ with $\mu \circ \sigma : H \otimes H \xrightarrow{\sigma} H \otimes H \xrightarrow{\mu} H$ and replacing δ with $\sigma \circ \delta : H \xrightarrow{\delta} H \otimes H \xrightarrow{\sigma} H \otimes H$, whilst η , ε and ν remain the same.

There is a bialgebra H' obtained by replacing δ with $\sigma \circ \delta$, whilst the rest of the operations remain the same, but H' does not have to be a Hopf algebra.

Proposition 8.1. Let H be a Hopf algebra, then:

1. The antipode ν is uniquely determined.
2. $\nu : H^{op} \rightarrow H$ is a bialgebra morphism.
3. H' is a Hopf algebra if and only if ν is bijective.
4. If H is commutative or cocommutative, then $\nu \circ \nu = 1_H$.

Proof.

1. H is both an algebra and a coalgebra, so we have the convolution algebra structure on $\text{Hom}_R(H, H)$. An antipode is an inverse for $1_H \in \text{Hom}_R(H, H)$. In every monoid, an inverse is unique.
2. To show that $\nu : H^{op} \rightarrow H$ preserves multiplication, we must show that the following diagram commutes:

$$\begin{array}{ccccc} H \otimes H & \xrightarrow{\sigma} & H \otimes H & \xrightarrow{\nu \otimes \nu} & H \otimes H \\ \mu \downarrow & & & & \downarrow \mu \\ H & \xrightarrow{\nu} & H & & H \end{array}$$

We do this by showing that, under convolution, the left-hand side is a left inverse for $\mu \in \text{Hom}_R(H \otimes H, H)$, while the right-hand side is a right

inverse.

$$\begin{array}{c}
\begin{array}{c}
H \otimes H \xrightarrow{(\nu \circ \mu) * \mu} H \\
\downarrow \varepsilon \otimes \varepsilon \quad \searrow \delta \otimes \delta \quad \searrow \mu \quad \downarrow \mu \otimes \mu \\
R \xrightarrow{\eta} H \\
\downarrow \varepsilon \quad \downarrow \delta \quad \downarrow \nu \otimes 1 \quad \downarrow \mu \\
H \xrightarrow{\delta} H^{\otimes 2} \xrightarrow{\nu \otimes 1} H^{\otimes 2} \xrightarrow{\mu} H
\end{array}
\end{array}$$

$$\begin{array}{c}
H^{\otimes 2} \xrightarrow{\delta \otimes \delta} H^{\otimes 4} \xrightarrow{1 \otimes \sigma \otimes 1} H^{\otimes 4} \xrightarrow{\mu \otimes 1 \otimes 1} H^{\otimes 3} \\
\downarrow \delta \otimes 1 \quad \downarrow 1 \otimes \sigma \quad \downarrow 1 \otimes \delta \otimes 1 \quad \downarrow \mu \otimes 1 \otimes 1 \quad \downarrow 1 \otimes \sigma \\
H^{\otimes 3} \xrightarrow{1 \otimes \sigma} H^{\otimes 3} \xrightarrow{1 \otimes \delta \otimes 1} H^{\otimes 4} \xrightarrow{\mu \otimes 1 \otimes 1} H^{\otimes 3} \\
\downarrow 1 \otimes 1 \otimes \varepsilon \quad \downarrow 1 \otimes \varepsilon \otimes 1 \quad \downarrow 1 \otimes 1 \otimes \nu \otimes 1 \quad \downarrow 1 \otimes \nu \otimes 1 \\
H \xrightarrow{\delta} H^{\otimes 2} \xrightarrow{1 \otimes \eta \otimes 1} H^{\otimes 3} \xrightarrow{\mu \otimes 1} H^{\otimes 3} \\
\downarrow 1 \otimes 1 \quad \downarrow 1 \otimes \mu \quad \downarrow 1 \otimes \mu \quad \downarrow 1 \otimes \mu \quad \downarrow \mu \\
R \xrightarrow{\eta} H \\
\downarrow \varepsilon \quad \downarrow \delta \quad \downarrow \nu \otimes 1 \quad \downarrow \mu \\
H \xrightarrow{\delta} H^{\otimes 2} \xrightarrow{\nu \otimes 1} H^{\otimes 2} \xrightarrow{\mu} H
\end{array}$$

Thus ν preserves multiplication. The following proves that ν preserves

unit.

$$\begin{array}{ccccc}
 R & \xrightarrow{\eta} & H & \xrightarrow{\nu} & H \\
 \parallel & \searrow \eta & \searrow \eta \otimes \eta & & \swarrow 1 \otimes \eta \\
 & & H & \xrightarrow{\delta} & H^{\otimes 2} & \xrightarrow{\nu \otimes 1} & H^{\otimes 2} \\
 & \swarrow \varepsilon & & & \searrow \mu & & \\
 R & & & & & & H \\
 & \xrightarrow{\eta} & & & & &
 \end{array}$$

3. ν' is a right inverse for ν iff $\nu \circ \nu' = 1_H$ iff $\nu \circ \nu'$ is a convolution left inverse for ν iff the following commutes:

$$\begin{array}{ccccccc}
 H & \xrightarrow{\delta} & H \otimes H & \xrightarrow{1 \otimes \nu'} & H \otimes H & \xrightarrow{\nu \otimes \nu} & H \otimes H & \xrightarrow{\mu} & H \\
 & \searrow \varepsilon & & & & & \nearrow \eta & & \\
 & & & & R & & & &
 \end{array}$$

iff the following commutes

$$\begin{array}{ccccccc}
 H & \xrightarrow{\delta} & H \otimes H & \xrightarrow{1 \otimes \nu'} & H \otimes H & \xrightarrow{\sigma} & H \otimes H & \xrightarrow{\mu} & H & \xrightarrow{\nu} & H \\
 & \searrow \varepsilon & & & & & \nearrow \eta & & & & \\
 & & & & R & & & & & &
 \end{array}$$

iff the following commutes

$$\begin{array}{ccccccc}
 H & \xrightarrow{\delta} & H \otimes H & \xrightarrow{\sigma} & H \otimes H & \xrightarrow{1 \otimes \nu'} & H \otimes H & \xrightarrow{\mu} & H & \xrightarrow{\nu} & H \\
 & \searrow \varepsilon & & & & & \nearrow \eta & & & & \\
 & & & & R & & & & & &
 \end{array}$$

4. If H is commutative, then $H' = H$, then H' is a Hopf algebra with antipode $\nu' = \nu$, and thus $\nu \circ \nu = 1_H$.

□

Proposition 8.2. Let H and K be Hopf algebras, then each bialgebra morphism $f : H \rightarrow K$ preserves antipode, i.e., the following square commutes:

$$\begin{array}{ccc}
 H & \xrightarrow{f} & K \\
 \nu \downarrow & & \downarrow \nu \\
 H & \xrightarrow{f} & K
 \end{array}$$

Proof. Clearly $f : D \rightarrow C$ and $g : A \rightarrow B$ are coalgebra and algebra morphisms respectively, then the following morphism

$$\mathrm{Hom}(C, A) \rightarrow \mathrm{Hom}(D, B)$$

realised as $u \mapsto g \circ u \circ f$ is a monoid morphism for the convolution structures. We also have the following monoid morphisms

$$_ \circ f, f \circ _ : \mathrm{Hom}(H, H) \rightarrow \mathrm{Hom}(H, K)$$

that both take 1_H to f , but monoid morphisms take inverses to inverses. $\square$

Recall that H is an algebra, then H^0 is a coalgebra. If H is a bialgebra, then H^0 becomes a bialgebra using the multiplication:

$$H^0 \otimes H^0 \cong (H \otimes H)^0 \xrightarrow{\delta^0} H^0$$

and unit

$$R \cong R^0 \xrightarrow{\varepsilon^0} H^0.$$

Further, if H is a Hopf algebra, so is H^0 with antipode $\nu^0 : H^0 \rightarrow H^0$. What we have here is a contravariant “self-adjoint” functor

$$(_)^0 : \mathbf{Hopf}_R^{op} \rightarrow \mathbf{Hopf}_R.$$

What “self-adjoint” means in this context is that:

$$\mathbf{Big}_R(H, K^0) \cong \mathbf{Big}_R(K, H^0).$$

Proposition 8.3. If H is a Hopf algebra, then the monoid $\mathcal{D}(H)$ of set-like elements is a group.

Proof. Take any $g \in \mathcal{D}(H)$, then

$$\begin{aligned} \nu(g)g &= (\mu \circ (\nu \otimes 1))(g \otimes g) \\ &= (\mu \circ (\nu \otimes 1))\delta(g) \\ &= (\mu \circ (\nu \otimes 1) \circ \delta)(g) = \eta(\varepsilon(g)) = \eta(1) = 1 \end{aligned}$$

$\square$

An *A-point* of a Hopf algebra H is an algebra morphism $f : H \rightarrow A$.

Proposition 8.4. Let H be a Hopf algebra, then:

1. If $f, g : H \rightarrow A$ are commuting A -points of H (i.e. $[f(h), g(k)] = 0$ for each $h, k \in H$), then $f * g : H \rightarrow A$ is an A -point of H .
2. If $f : H \rightarrow A$ is an A -point of H , then f has a convolution inverse $f \circ \nu : H^{op} \rightarrow A$, which is an A -point of H^{op}

Example 8.1. For a monoid G , the monoid algebra $R(G)$ is a bialgebra. If G is a group, then the group algebra $R(G)$ becomes a Hopf algebra with antipode $\nu : R(G) \rightarrow R(G)$ given by $\nu(g) = g^{-1}$.

Example 8.2. Let L be a Lie algebra, L^{op} stands for the Lie algebra with the same underlying module L , but with the Lie bracket β^{op} given by $\beta^{op}(x, y) = \beta(y, x)$. We have $(A^{op})_L = (A_L)^{op}$. Thus it follows that we have a canonical algebra isomorphism:

$$\mathcal{U}(L^{op}) \cong \mathcal{U}(L)^{op}.$$

We have a Lie algebra isomorphism $L \rightarrow L^{op}$ taking x to $-x$. So we define $\nu : \mathcal{U}(L) \rightarrow \mathcal{U}(L)^{op}$ by

$$\begin{array}{ccc} L & \xrightarrow{-} & L^{op} \\ i \downarrow & & \downarrow i \\ \mathcal{U}(L) & \xrightarrow{\nu} & \mathcal{U}(L)^{op} \cong \mathcal{U}(L^{op}) \end{array}$$

One can also check that for $x_1, \dots, x_n \in L$

$$\nu(i(x_1) \cdots i(x_n)) = (-1)^n i(x_1) \cdots i(x_n)$$

$\mathcal{U}(L)$ becomes a Hopf algebra with antipode.

Example 8.3. The matrix algebra $M(n)$ is not a Hopf algebra. One needs to adjoin an inverse for the determinant. Recall that $M(n) = R[X] = \mathcal{S}(\mathcal{F}_R(X))$ for $X = \{x_{ij} \mid i, j \in 1, \dots, n\}$. Define

$$\det(X) = \sum_{\xi \in \mathcal{S}_n} (-1)^{|\xi|} x_{1\xi(1)} x_{2\xi(2)} \cdots x_{n\xi(n)}$$

where $|\xi|$ is the smallest number of simple transpositions required to obtain a permutation ξ . Let us form the following commutative polynomial R -algebra:

$$R[X \cup \{t\}] = R[(x_{ij}), t] = \mathcal{S}(\mathcal{F}_R(X \cup t))$$

in $n^2 + 1$ commuting indeterminates t and x_{ij} for $1 \leq i, j \leq n$. Put:

$$\mathrm{GL}(n) = R[X \cup \{t\}] / (t \det(X) - 1)$$

as a commutative R -algebra. We make $\mathrm{GL}(n)$ a bialgebra the following way:

- $\delta(x_{ij}) = \sum_{k=1}^n x_{ik} \otimes x_{kj},$
- $\delta(t) = t \otimes t,$
- $\varepsilon(x_{ij}) = \delta_{ij},$
- $\varepsilon(t) = 1$

modulo $(t \det(X) - 1)$. Put $X_{ij} = \{x_{rs} \mid r \neq i, s \neq j\}$. Let us define the morphism $\nu : \mathrm{GL}(n) \rightarrow \mathrm{GL}(n)$ by putting:

- $\nu(x_{ij}) = t \det(X_{ji})$,
- $\nu(t) = \det(X)$

modulo $(t \det(X) - 1)$. Then $\mathrm{GL}(n)$ becomes a Hopf algebra.

Let A be a commutative R -algebra, then we have a canonical isomorphism of groups:

$$\mathbf{Alg}_R(\mathrm{GL}(n), A) \cong \mathrm{GL}(n, A).$$

Example 8.4. This example is a quantum deformation of the previous one. Take $X = \{x_{ij} \mid i, j \in \{1, \dots, n\}\}$. We form the free algebra $R\langle X \rangle = T(\mathcal{F}_R(X))$ on the non-commuting indeterminates x_{ij} . Let $M_q(n)$ denote the quotient of $R\langle X \rangle$ by the ideal generated by the following elements:

- $x_{ir}x_{jk} - x_{jk}x_{ir}$ for $i < j$ and $k < r$,
- $x_{ir}x_{jk} - x_{jk}x_{ir} - (q - q^{-1})x_{ik}x_{jr}$ for $i < j$ and $k < r$,
- $x_{ik}x_{jk} - qx_{jk}x_{ik}$ for $i < j$,
- $x_{ik}x_{ir} - qx_{ir}x_{ik}$ for $k < r$.

Thus we have a coalgebra with comultiplication (modulo the ideal):

$$\delta(x_{ij}) = \sum r = 1^n x_{ir} \otimes x_{rj}$$

and counit $\varepsilon(x_{ij}) = \delta_{ij}$, i.e. Kronecker delta. “The quantum determinant” is defined the following way

$$\det_q(X) = \sum_{\xi \in \mathcal{S}_n} (-q)^{|\xi|} x_{1(\xi(1))} x_{2(\xi(2))} \dots x_{n(\xi(n))}$$

which is a central element of $M_q(n)$. The quantum general linear group is defined by:

$$GL_q(n) = M_q(n)[t]/(t \det_q(X) - 1).$$

We modify the multiplication and counit of $M_q(n)$ by putting $\delta(t) = t \otimes t$ and $\varepsilon(t) = 1$. Thus we have a bialgebra epimorphism $\rho : M_q(n) \rightarrow GL_q(n)$. Define $\nu : GL_q(n) \rightarrow GL_q(n)$ by $\nu(x_{ij}) = t \det_q(X_{ji})$ and $\nu(t) = \det_q(X)$.

8.1 Exercise

Exercise 8.1. Assume that the base ring R is a field. An ideal in an algebra A is a submodule I such that $\mu(I \otimes A + A \otimes I) \subseteq I$. Then A/I becomes an algebra. A *coideal* of a coalgebra C is a submodule I such that $\delta(I) \subseteq I \otimes C + C \otimes I$ and $\varepsilon(I) \subseteq 0$. Show the following:

1. If I is a coideal of a coalgebra C , describe a coalgebra structure on C/I for which $\rho : C \rightarrow C/I$ becomes a coalgebra morphism. If C is a bialgebra and I is an ideal, show that C/I is a bialgebra. What condition on I ensures that C/I has an antipode if C has?
2. Check that the polynomial R -algebra $B = R\langle x, y, z \rangle$ on three non-commuting indeterminates is a bialgebra with:
 - $\delta(x) = x \otimes x$,
 - $\delta(y) = y \otimes y$,
 - $\delta(z) = 1 \otimes z + z \otimes 1$,
 - $\varepsilon(x) = \varepsilon(y) = 1$,
 - $\varepsilon(z) = 0$.
3. Verify that the ideal $(xy - 1, yx - 1)$ is a coideal in B . Let H denote the quotient algebra further.
4. Show that H is a Hopf algebra with antipode ν given by (modulo the ideal $(xy - 1, yx - 1)$):
 - $\nu(x) = y$,
 - $\nu(y) = x$,
 - $\nu(z) = -zy$.

Show also that $\nu^{2n}(z) = x^n z y^n$ and $\nu^{2n+1}(x) = -x^n z y^{n+1}$. Therefore, this ν has infinite order.

5. Show that the ideals $I_n = (x^n z y^n - z)$ and $J_n = (x^n - 1)$ are Hopf algebras in H . Show that the antipodes of both H/I_n and H/J_n have order $2n$.

Proof.

1. Let I be a coideal of C , so I is a submodule satisfying $\delta(I) \subseteq I \otimes C + C \otimes I$. The composite:

$$C \xrightarrow{\delta} C \otimes C \xrightarrow{\rho \otimes \rho} C/I \otimes C/I$$

maps I to 0 since:

$$\begin{aligned} (\rho \otimes \rho)\delta(I) &\subseteq (\rho \otimes \rho)(I \otimes C + C \otimes I) = \\ &\rho(I) \otimes \rho(C) + \rho(C) \otimes \rho(I) = \\ &0 \otimes \rho(C) + \rho(C) \otimes 0 = 0 \end{aligned}$$

Thus there is a unique module morphism $\delta : C/I \rightarrow C/I \otimes C/I$ such that:

$$\begin{array}{ccc} C & \xrightarrow{\rho} & C/I \\ \delta \downarrow & & \downarrow \delta \\ C \otimes C & \xrightarrow{\rho \otimes \rho} & C/I \otimes C/I \end{array}$$

Similarly, $\varepsilon(I) = 0$ implies that there is a unique $\varepsilon : C/I \rightarrow R$ such that:

$$\begin{array}{ccc} C & \xrightarrow{\varepsilon} & R \\ & \searrow \rho & \nearrow \varepsilon \\ & C/I & \end{array}$$

All those facts about δ and ε will mean that $\rho : C \rightarrow C/I$ is a coalgebra morphism if one ensures that C/I is a coalgebra. To show that $\delta : C/I \rightarrow C/I \otimes C/I$ is coassociative, one needs to take the coassociativity diagram for C/I and precompose it with $\rho : C \rightarrow C/I$:

$$\begin{array}{ccc} C & \xrightarrow{\rho} & C/I \\ \downarrow \delta & & \downarrow \delta \\ C \otimes C & \xrightarrow{\rho \otimes \rho} & C/I \otimes C/I \\ \downarrow 1 \otimes \delta \quad \downarrow \delta \otimes 1 & & \downarrow 1 \otimes \delta \quad \downarrow \delta \otimes 1 \\ C \otimes C \otimes C & \xrightarrow{\rho \otimes \rho \otimes \rho} & C/I \otimes C/I \otimes C/I \end{array}$$

And this commutes by coassociativity of C . But ρ is also surjective, so the coassociativity diagram for C/I (the bottom one) also commutes. Similarly, we take the unitality diagram for C/I and conclude its commutativity from the following one:

$$\begin{array}{ccc} C & \xrightarrow{\rho} & C/I \\ \downarrow \delta & & \downarrow \delta \\ C \otimes C & \xrightarrow{\rho \otimes \rho} & C/I \otimes C/I \\ \downarrow 1 \otimes \varepsilon \quad \downarrow \varepsilon \otimes 1 & & \downarrow 1 \otimes \varepsilon \quad \downarrow \varepsilon \otimes 1 \\ C & \xrightarrow{\rho} & C/I \end{array}$$

1 1

Therefore $\varepsilon : C/I \rightarrow R$ is a counit and C/I is a bialgebra. The rest is to check the bialgebra conditions, the most complicated one is the condition

that δ commutes with multiplication:

$$\begin{array}{ccc}
C/I \otimes C/I & \xrightarrow{\mu} & C/I \\
\delta \otimes \delta \downarrow & & \downarrow \delta \\
C/I \otimes C/I \otimes C/I \otimes C/I & \xrightarrow{1 \otimes \sigma \otimes 1} C/I \otimes C/I \otimes C/I \otimes C/I \xrightarrow{\mu \otimes \mu} & C/I \otimes C/I
\end{array}$$

To show that, one precomposes the above diagram with $\rho \otimes \rho : C \otimes C \rightarrow C/I \otimes C/I$ and use the corresponding condition on C and the fact that $\rho \otimes \rho$ is surjective.

2. Put R -algebra $B = R\langle x, y, z \rangle$, then their equations define algebra morphisms $\delta : B \rightarrow B \otimes B$ and $\varepsilon : B \rightarrow R$ since B is free an algebra. Let us check that those morphisms make B a bialgebra. The coassociativity is showed the following way for x and similarly for y :

$$\begin{aligned}
(\delta \otimes 1)\delta(x) &= (\delta \otimes 1)(x \otimes x) \\
&= x \otimes x \otimes x \\
&= (1 \otimes \delta)(x \otimes x) = (1 \otimes \delta)\delta(x)
\end{aligned}$$

The coassociativity for z :

$$\begin{aligned}
(\delta \otimes 1)\delta(x) &= (\delta \otimes 1)(1 \otimes z + z \otimes x) \\
&= 1 \otimes 1 \otimes z + (1 \otimes z + z \otimes x) \otimes x \\
&= 1 \otimes (1 \otimes z + z \otimes x) + z \otimes x \otimes x \\
&= (1 \otimes \delta)(1 \otimes z + z \otimes x) = (1 \otimes \delta)\delta(z)
\end{aligned}$$

The counit conditions for x (and similarly for y) and for z :

$$\begin{aligned}
(\varepsilon \otimes 1)\delta(x) &= (\varepsilon \otimes 1)(x \otimes x) = \varepsilon(x)(x) = x \\
&= (1 \otimes \varepsilon)(x \otimes x) = (1 \otimes \varepsilon)\delta(x).
\end{aligned}$$

$$\begin{aligned}
(\varepsilon \otimes 1)\delta(z) &= (\varepsilon \otimes 1)(1 \otimes z + z \otimes x) = z + 0x = z \\
&= 0 + z = (\varepsilon \otimes 1)(1 \otimes z + z \otimes x) = (\varepsilon \otimes 1)\delta(z)
\end{aligned}$$

3. One has

$$\begin{aligned}
\delta(xy - 1)(x \otimes x)(y \otimes y) - 1 \otimes 1 &= xy \otimes xy - 1 \otimes 1 \\
&= (xy - 1) \otimes xy + 1 \otimes xy - 1 \otimes 1 \\
&= (xy - 1) \otimes xy + 1 \otimes (xy - 1) \subseteq I \otimes B + B \otimes I.
\end{aligned}$$

4. We must check that the following is satisfied:

$$\begin{array}{ccccc}
H & \xrightarrow{\delta} & H \otimes H & \xrightleftharpoons[1 \otimes \nu]{\nu \otimes 1} & H \otimes H & \xrightarrow{\mu} & H \\
& & \searrow \varepsilon & & \nearrow \eta & & \\
& & R & & & &
\end{array}$$

Let us check this condition for x (the calculation for y is the same) and z :

$$\begin{aligned}\mu(\nu \otimes 1)\delta(x) &= \mu(\nu \otimes 1)(x \otimes x) = \mu(y \otimes x) = yx \equiv 1 \\ &= \nu(\varepsilon(x)) \equiv xy = \mu(x \otimes y) = \mu(1 \otimes \nu)(x \otimes x) = \mu(1 \otimes \nu)\delta(x)\end{aligned}$$

$$\begin{aligned}\mu(\nu \otimes 1)\delta(z) &= \mu(\nu \otimes 1)(1 \otimes z + z \otimes x) = \mu(1 \otimes z + (-zy) \otimes x) \\ &= z - zyx \equiv 0 = \eta\varepsilon(z) = 0 \\ &= -zy + zy = \mu(1 \otimes (-zy) + z \otimes y) \\ &= \mu(1 \otimes \nu)(1 \otimes z + z \otimes x) = \mu(1 \otimes \nu)\delta(z)\end{aligned}$$

Thus H is a Hopf algebra. We know that ν reverses both multiplication and comultiplication.

Let us show that $\nu^{2n}(z) = x^n zy^n$ and $\nu^{2n+1}(z) = -x^n zy^{n+1}$ for each $n < \omega$. The statement is trivial for $n = 0$. Let $n = k + 1$ and assume that $\nu^{2k}(z) = x^k zy^k$ and $\nu^{2k+1}(z) = -x^k zy^{k+1}$. We need $\nu^{2k+2}(z) = x^{k+1} zy^{k+1}$ and $\nu^{2(k+1)+1}(z) = -x^{k+1} zy^{k+2}$. First of all, observe that:

$$\nu^{2k+1}(z) = -x^k zy^{k+1} = -x^k zyy^k = x^k (-zy)y^k = \nu(y)^k \nu(z) \nu(x)^k$$

Thus:

$$\begin{aligned}\nu^{2(k+1)}(z) &= -\nu(y)^{n+1} \nu(z) \nu(x)^n \\ &\quad - x^{n+1} (-zy)y^n = x^{n+1} zy^{n+1}\end{aligned}$$

Similarly:

$$\begin{aligned}\nu^{2k+3}(z) &= \nu^{2(k+1)+1}(z) \\ &= -x^{k+1} zy^{k+2} = -x^{k+1} zyy^{k+1} \\ &= x^{k+1} (-zy)y^{k+1} = \nu(y)^{k+1} \nu(z) \nu(x)^{k+1}\end{aligned}$$

Therefore:

$$\begin{aligned}\nu^{2(k+1)+1}(z) &= \nu(y)^{k+1} \nu(z) \nu(x)^{k+1} \\ &= \nu(x)^{k+1} (-zy) \nu(y)^{k+1} = -\nu(x)^{k+1} z \nu(y)^{k+2}\end{aligned}$$

5. Consider the ideals $I_n = (x^n zy^n - z)$ and $J_n = (x^n - 1)$. Let us evaluate ν , δ and ε . First of all:

$$\varepsilon(x^n zy^n - z) = \varepsilon(z(x^n y^n - 1)) = \varepsilon(z)\varepsilon(x^n y^n - 1) = 0$$

Further,

$$\begin{aligned}\delta(x^n zy^n - z) &= (x^n \otimes x^n)(1 \otimes z + z \otimes x)(y^n \otimes y^n) - 1 \otimes z - z \otimes x \\ &= (x^n \otimes z + x^n z \otimes x^{n+1})(y^n \otimes y^n) - 1 \otimes z - z \otimes x \\ &= x^n y^n \otimes zy^n + x^n zy^n \otimes x^{n+1} y^n - 1 \otimes z - z \otimes x \\ &= 1 \otimes x^n zy^n + x^n zy^n \otimes x - 1 \otimes z - z \otimes x \\ &= 1 \otimes (x^n zy^n - z) + (x^n zy^n - z) \otimes x \in I_n \otimes H + H \otimes I_n.\end{aligned}$$

And also:

$$\begin{aligned}\nu(x^n zy^n - z) &= \nu(x^n zy^n) - \nu(z) \\ &= x^n(-zy)y^n + zy = -x^n zy^{n+1} + zy = (x^n zy^n - z)(-y) \in I_n\end{aligned}$$

Consider $J_n = (x^n - 1)$ similarly:

$$\varepsilon(x^n - 1) = 0$$

Then further:

$$\delta(x^n - 1) = x^n \otimes x^n - 1 \otimes 1 = (x^n - 1) \otimes x^n + 1 \otimes (x^n - 1) \in J_n \otimes H + H \otimes J_n$$

And finally:

$$\nu(x^n - 1) = y^n - 1 \equiv -y^n(x^n - 1) \pmod{J_n}.$$

□

9 Representation of Quantum Groups

Note that if $f : E \rightarrow A$ is a ring morphism, then each (left) A -module M becomes a (left) E -module via the following action for $e \in E$ and $m \in M$:

$$em = f(e)m$$

This action is called *restriction of scalars along f* .

Let A be an R -algebra, then each module is automatically an R -module via restriction of scalars along the unit $\eta : R \rightarrow A$. Alternatively, one can view an A -module as an R -module M with a ring morphism $\hat{\mu} : A \rightarrow \text{End}_R(M)$. Let us elaborate on how the concept of an A -module is dualised.

An R -module M with a module morphism $\mu : A \otimes_R M \rightarrow M$, called the *action* of A on M , satisfying the following:

$$\begin{array}{c} A \otimes_R A \otimes_R M \xrightleftharpoons[1 \otimes \mu]{\mu \otimes 1} A \otimes_R M \xrightarrow{\mu} M \\ \begin{array}{ccc} & A \otimes_R M & \\ \nearrow & & \searrow \\ M & \xlongequal{\quad} & M \end{array} \end{array}$$

We write $\mathbf{Mod}_R(A)$ for $\mathbf{Mod}_A$ to emphasise that we build it up from $\mathbf{Mod}_R$.

Suppose M and N are (left) modules over the R -module A . Regard $M : A \not\rightarrow R$ and $N : R \not\rightarrow A^{op}$. One can see that $M \otimes_R N : A \not\rightarrow A^{op}$, thus, $M \otimes_R N$ becomes an $A \otimes A$ -module. If A is a bialgebra, then we can restrict

along $\delta : A \rightarrow A \otimes_R A$ to obtain an A -module structure on $M \otimes_R N$. We give the action explicitly as the following composite:

$$A \otimes_R M \otimes_R N \xrightarrow{\delta \otimes 1 \otimes 1} A \otimes_R A \otimes_R M \otimes_R N \xrightarrow{1 \otimes \sigma \otimes 1} A \otimes_R M \otimes_R A \otimes_R N \xrightarrow{\mu \otimes \mu} M \otimes_R N$$

With M and N left A -modules, one can regard $M : R \nrightarrow A^{op}$ and $N : R \nrightarrow A^{op}$, so that $\text{Hom}_R(M, N) : A^{op} \nrightarrow A^{op}$. Or, in other words, $\text{Hom}_R(M, N)$ is an $A^{op} \otimes A$ -module. If $A = H$ is a Hopf algebra, then one can restrict scalars along the R -algebra morphism:

$$H \xrightarrow{\delta} H \otimes_R H \xrightarrow{\nu \otimes 1} H^{op} \otimes_R H$$

to make $\text{Hom}_R(M, N)$ into an H -module. The action of H on $\text{Hom}_R(M, N)$ is given explicitly as the following composite:

$$\begin{array}{c} H \otimes_R \text{Hom}_R(M, N) \\ \downarrow \delta \otimes 1 \\ H \otimes_R H \otimes_R \text{Hom}_R(M, N) \\ \downarrow 1 \otimes \sigma \\ H \otimes_R \text{Hom}_R(M, N) \otimes_R H \\ \downarrow \mu_1 \otimes \nu \\ \text{Hom}_R(M, N) \otimes_R H \\ \downarrow \mu_2 \\ \text{Hom}_R(M, N) \end{array}$$

where μ_1 and μ_2 are the left and the right actions:

$$\begin{aligned} H \otimes_R \text{Hom}_R(M, N) &\xrightarrow{\hat{\mu} \otimes 1} \text{Hom}_R(N, N) \otimes_R \text{Hom}_R(M, N) \xrightarrow{\circ} \text{Hom}_R(M, N) \\ \text{Hom}_R(M, N) \otimes_R H &\xrightarrow{\hat{\mu} \otimes 1} \text{Hom}_R(M, N) \otimes_R \text{Hom}_R(M, M) \xrightarrow{\circ} \text{Hom}_R(M, N) \end{aligned}$$

given as:

$$\begin{aligned} \mu_1 : h \otimes f &\mapsto (m \mapsto h(fm)) \\ \mu_2 : h \otimes f &\mapsto (m \mapsto f(hm)) \end{aligned}$$

Proposition 9.1. For left modules M and N over the Hopf algebra H , the canonical R -modules morphisms

- $e : \text{Hom}_R(M, N) \otimes_R M \rightarrow N$ with $e : f \otimes m \mapsto f(m)$,
- $d : M \rightarrow \text{Hom}_R(N, M \otimes_R N)$ with $m \mapsto (n \mapsto m \otimes n)$

are left H -module morphisms.

Proof. Routine. □

Corollary 9.1. For modules M , N and L over a Hopf algebra H , the canonical isomorphism:

$$\mathrm{Hom}_R(M \otimes_R N, L) \cong \mathrm{Hom}_R(M, \mathrm{Hom}_R(N, L))$$

restrict to an isomorphism

$$\mathrm{Hom}_H(M \otimes_R N, L) \cong \mathrm{Hom}_H(M, \mathrm{Hom}_R(N, L)).$$

The tensor and the hom are also preserved by the functor $\mathbf{Mod}_R(H) \rightarrow \mathbf{Mod}_R$, which is given by the H -action.

Let C be an R -coalgebra, a (left) C -comodule is an R -module M equipped with a module morphism $\delta : M \rightarrow C \otimes_R M$, called the coaction of C on M such that:

$$\begin{array}{c} M \xrightarrow{\delta} C \otimes_R M \xrightleftharpoons[1 \otimes \delta]{\delta \otimes 1} C \otimes_R C \otimes_R M \\ \\ \begin{array}{ccc} & C \otimes_R M & \\ \delta \nearrow & & \searrow \varepsilon \otimes 1 \\ M & \xlongequal{\quad\quad\quad} & M \end{array} \end{array}$$

$\mathbf{Com}_R(C)$ stands for the category of C -comodules and *comodule* morphisms, i.e. R -module morphisms $f : M \rightarrow N$ such that

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \delta \downarrow & & \downarrow \delta \\ C \otimes_R M & \xrightarrow{1 \otimes f} & C \otimes_R N \end{array}$$

Each C -comodule becomes a C^* -module with the following action:

$$C^* \otimes_R M \xrightarrow{1 \otimes \delta} C^* \otimes_R C \otimes_R M \xrightarrow{e \otimes 1} M.$$

Recall that by the fundamental theorem of Morita theory, if C is a Cauchy module, then there is a bijection between C -coactions δ and C^* -actions μ on each R -module M , and we recover δ as the composite:

$$M \xrightarrow{d \otimes 1} C \otimes_R C^* \otimes_R M \xrightarrow{1 \otimes \mu} C \otimes_R M$$

Thus if C is Cauchy, then the following categories are isomorphic:

$$\mathbf{Com}_R(C) \cong \mathbf{Mod}_R(C^*)$$

If C is an R -bialgebra, but necessarily Cauchy, then a coaction on the tensor product of M and N over R is given by the composite:

$$M \otimes_R N \xrightarrow{\delta \otimes \delta} C \otimes_R M \otimes_R C \otimes_R N \xrightarrow{\sigma} C \otimes_R C \otimes_R M \otimes_R N \xrightarrow{\mu \otimes 1 \otimes 1} C \otimes_R M \otimes_R N$$

The empty tensor product R has the coaction $\eta : R \rightarrow C \otimes_R R$.

Proposition 9.2. Each Cauchy R -module M lifts to an R -coalgebra $M \otimes_R M^*$ with counit $e : M \otimes_R M^* \rightarrow R$ and comultiplication:

$$1 \otimes d \otimes 1 : M \otimes_R M^* \rightarrow M \otimes_R M^* \otimes_R M \otimes_R M^*.$$

For any R -coalgebra C , the assignment

$$\hat{\delta} = (M \otimes_R M^* \xrightarrow{\delta \otimes 1} C \otimes_R M \otimes_R M^* \xrightarrow{1 \otimes e} C)$$

determines a bijection between coactions $\delta : M \rightarrow C \otimes_R M$ of C on M and coalgebra morphisms $\hat{\delta} : M \otimes_R M^* \rightarrow C$.

Proof. The following composite

$$M \otimes_R M^* \xrightarrow{\cong} M^* \otimes M \xrightarrow{\cong} \text{Hom}_R(M, M)$$

has the universal of Hom under reversal of arrows; so the diagrammatic proof that $\text{End}_R(M)$ is an algebra and that an action is an algebra morphism of the kind $A \rightarrow \text{End}_R(M)$, dualises. $\square$

Take $M = R^n$ in the above proposition and let $e_1, \dots, e_n$ be the standard basis. Now let $e_1^*, \dots, e_n^*$ be the dual basis for R^{n*} , so $e_i^*(e_j) = \delta_{ij}$ (Kronecker delta). A coaction of C on R^n amounts to a coalgebra morphism

$$\hat{\delta} : R^n \otimes_R R^{n*} \rightarrow C.$$

$\hat{\delta}$ is determined by its values on the basis of elements $e_i \otimes e_j^*$ of $R^n \otimes_R R^{n*}$:

$$\hat{\delta}(e_i \otimes e_j^*) = x_{ij} \in C.$$

C -comodule structures on R^n are in bijection with multiplicative matrices in C , i.e., matrices $\vec{x} = (x_{ij})$ in C such that:

$$\begin{aligned} \delta(x_{ij}) &= \sum_k x_{ik} \otimes x_{kj}, \\ \varepsilon(x_{ij}) &= \delta_{ij}. \end{aligned}$$

One can write the last two equations the following way:

$$\begin{aligned} \delta(\vec{x}) &= \vec{x} \otimes \vec{x} \\ \varepsilon(\vec{x}) &= \vec{1}, \end{aligned}$$

where $\vec{1}$ is the identity matrix and $\vec{x} \otimes \vec{x} = \sum_k x_{ik} \otimes x_{kj}$ is *not* the usual tensor product of matrices.

Proposition 9.3.

1. $\hat{\delta} : M \otimes_R M^* \rightarrow H$ is a coalgebra morphism if and only if the composite

$$M^* \otimes_R M \xrightarrow{\sigma} M \otimes_R M^* \xrightarrow{\hat{\delta}} H^{op}$$

is a coalgebra morphism.

2. Suppose M is an H -module and put

$$\hat{\delta}_k = (M \otimes_R M^* \xrightarrow{\hat{\delta}} H \xrightarrow{\nu^k} H).$$

For k even, $\hat{\delta}_k$ is a coalgebra morphism and has convolution inverse $\hat{\delta}_{k+1}$ in $\text{Hom}_R(M \otimes_R M^*, H)$.

Proof.

1. This part is proved with the following commutative diagrams:

$$\begin{array}{ccccc}
M^* \otimes_R M & \xrightarrow{\sigma} & M \otimes_R M^* & \xrightarrow{\hat{\delta}} & H \\
1 \otimes d \otimes 1 \downarrow & & \downarrow 1 \otimes d \otimes 1 & & \downarrow \sigma \\
M^* \otimes_R M^* \otimes_R M \otimes_R M & \xrightarrow{\sigma_{4231}} & M \otimes_R M^* \otimes_R M \otimes_R M^* & \xrightarrow{\hat{\delta} \otimes \hat{\delta}} & H \otimes H \\
1 \otimes \sigma \otimes 1 \downarrow & & \downarrow \sigma_{3412} & & \downarrow \sigma \\
M^* \otimes_R M \otimes_R M^* \otimes_R M & \xrightarrow{\sigma \otimes \sigma} & M \otimes_R M^* \otimes_R M \otimes_R M^* & \xrightarrow{\hat{\delta} \otimes \hat{\delta}} & H \otimes H
\end{array}$$

$$\begin{array}{ccccc}
M^* \otimes_R M & \xrightarrow{\sigma} & M \otimes_R M^* & \xrightarrow{\hat{\delta}} & H \\
\sigma \downarrow & \nearrow & \searrow e & & \downarrow \varepsilon \\
M \otimes_R M^* & \xrightarrow{e} & & & R
\end{array}$$

2. $\nu : H \rightarrow H^{op}$ is a coalgebra morphism, so is $\nu^k : H \rightarrow H$ for k even. Thus $\hat{\delta}_k = \nu^k \circ \hat{\delta}$ is a coalgebra morphism. Thus that the right composition with $\hat{\delta}_k$ preserves convolution. 1_H and ν are convolution inverses, so are $1_H \circ \hat{\delta}_k$ are $\nu \circ \hat{\delta}_k$, that is, so are $\hat{\delta}_k$ and $\hat{\delta}_{k+1}$.

□

10 Monoidal Categories

A *braiding* for a monoidal category $\mathcal{V}$ is a family of natural isomorphisms:

$$c_{A,B} : A \otimes B \rightarrow B \otimes A$$

such that the following diagrams commute:


$$\begin{array}{ccccc}
& & A \otimes (B \otimes C) & \xrightarrow{c_{A,B \otimes C}} & (B \otimes C) \otimes A \\
& \nearrow a_{A,B,C} & & & \searrow a_{B,C,A} \\
(A \otimes B) \otimes C & & & & B \otimes (C \otimes A) \\
& \searrow c_{A,B} \otimes 1_C & (B \otimes A) \otimes C & \xrightarrow{a_{B,A,C}} & B \otimes (A \otimes C) \\
& & & & \nearrow c_{A,C} \otimes 1 \\
& & & & B \otimes (C \otimes A)
\end{array}$$

$$\begin{array}{ccccc}
& & (A \otimes B) \otimes C & \xrightarrow{c_{A \otimes B,C}} & C \otimes (A \otimes B) \\
& \nearrow a_{A,B,C}^{-1} & & & \searrow a^{-1}_{C,A,B} \\
A \otimes (B \otimes C) & & & & (C \otimes A) \otimes B \\
& \searrow 1 \otimes c_{B,C} & A \otimes (C \otimes B) & \xrightarrow{a_{A,C,B}^{-1}} & (A \otimes C) \otimes B \\
& & & & \nearrow c_{A,C} \otimes 1 \\
& & & & (C \otimes A) \otimes B
\end{array}$$

A *braided monoidal category* is a monoidal category with a choice of braiding. A *symmetry* for a tensor category is a braiding satisfying the extra condition:

$$\begin{array}{ccc}
A \otimes B & \xlongequal{\quad} & A \otimes B \\
& \searrow c_{A,B} & \nearrow c_{B,A} \\
& B \otimes A &
\end{array}$$


Example 10.1. The *braid category* $\mathcal{B}$ has as objects the set of natural numbers and as arrows $\alpha : n \rightarrow n$ the braids on n strings. No arrows $m \rightarrow n$ for $n \neq m$. A braid $\alpha : n \rightarrow n$:




on n strings can be viewed as an element of the Artin braid group $\mathcal{B}_n$ with generators $s_1, \dots, s_{n-1}$ and the following relations:

- $s_i s_j = s_j s_i$ for $j < i - 1$,
- $s_{i+1} s_i s_{i+1} = s_i s_{i+1} s_i$


where s_i is a braid visualised the following way:




Composition of braids is multiplication is given by multiplication in $\mathcal{B}_n$:



The tensor product of braids adds the numbers of strings by placing one braid next to the other longitudinally:



Therefore $\mathcal{B}$ is a strict monoidal category. A braiding $c_{m,n} : m + n \rightarrow n + m$ is given by crossing the first m strings over the remaining n :



Example 10.2. The category $\mathbf{Mod}_R$ of modules over a commutative ring R is a symmetric monoidal category with tensor product $\otimes_R$ and symmetry $\sigma_{A,B} : A \otimes_R B \rightarrow B \otimes_R A$.

Example 10.3. Let A be an R -algebra, if M and N are A -modules, there is an A -module structure on $M \otimes_R N$ is given by:

$$a \cdot (m \otimes n) = \sum_{(a)} a_{(1)} m \otimes a_{(2)} n.$$

Thus $\mathbf{Mod}_R(A)$ is a monoidal category with $\otimes_R$. $\mathbf{Mod}_R(A)$ is symmetric if A is commutative, but we will be interested in a non-commutative A . A braiding $c_{M,N} : M \otimes_R N \rightarrow N \otimes_R M$, in particular, gives a morphism $c_{A,A} : A \otimes_R A \rightarrow A \otimes_R A$, which, in turn, gives an element $\gamma = c_{A,A}(1 \otimes 1) \in A \otimes A$.

Conversely, each element $\gamma = \sum_i u_i \otimes v_i \in A \otimes A$ determines a natural transformation with components $c_{M,N} : M \otimes_R N \rightarrow N \otimes_R M$ via the formula:

$$c_{M,N}(m \otimes n) = \sum_i (u_i n) \otimes (v_i m).$$

This is a bijection, as can be seen from the below diagram, where $\hat{m} : A \rightarrow M$ is the unique module morphism such that $\hat{m}(1) = m$:

$$\begin{array}{ccccc} R \otimes_R R & \xrightarrow{\eta} & A \otimes_R A & \xrightarrow{c_{A,A}} & A \otimes_R A \\ & & \hat{m} \otimes \hat{n} \downarrow & & \downarrow \hat{n} \otimes \hat{m} \\ & & M \otimes_R N & \xrightarrow{c_{M,N}} & N \otimes_R M \end{array}$$

To make each $c_{M,N}$ to be an isomorphism we need $\gamma \in A \otimes_R A$. We also need each c to be a module morphism:

$$c(a \cdot (m \otimes n)) = c\left(\sum_{(a)} (a_{(1)} m) \otimes (a_{(2)} n)\right) = \sum_{(a)} \sum_i (u_i a_{(2)} n) \otimes (v_i a_{(1)} m)$$

to be equal to

$$a \cdot c(m \otimes n) = a \cdot \sum_i (u_i n) \otimes (v_i m) = \sum_{(a)} \sum_i (a_{(1)} u_i n) \otimes (a_{(2)} v_i m).$$

Regarding $\gamma \in A \otimes_R A$ as a morphism $\gamma : R \rightarrow A \otimes_R A$, whose value at $1 \in R$ is the given γ , we can express this condition diagrammatically the following way (the B0 condition):

$$A \xrightarrow{\gamma \otimes \delta} A^{\otimes 4} \xrightarrow[\sigma_{3142}]{\sigma_{1423}} A^{\otimes 4} \xrightarrow{\mu \otimes \mu} A^{\otimes 2}$$

For a braiding, we require two more conditions:

$$\begin{aligned} c_{M,N \otimes L}(m \otimes n \otimes l) &= (1_N \otimes c_{M,L})(c_{M,N} \otimes 1_L)(m \otimes n \otimes l), \\ c_{M \otimes N,L}(m \otimes n \otimes l) &= (c_{M,L} \otimes 1_N)(1_M \otimes c_{N,L})(m \otimes n \otimes l). \end{aligned}$$

that is,

$$\begin{aligned} \sum_{i,(u_i)} u_{i(1)}n \otimes u_{i(2)}l \otimes v_i m &= \sum_{i,j} u_i n \otimes u_j l \otimes v_j v_i m, \\ \sum_{i,(v_i)} u_i l \otimes v_{i(1)}m \otimes v_{i(2)}n &= \sum_{i,j} u_j u_i l \otimes v_j m \otimes v_i n. \end{aligned}$$

These are equivalent to the following conditions respectively:

$$\begin{aligned} \sum_{i,(u_i)} u_{i(1)}n \otimes u_{i(2)}l \otimes v_i m &= \sum_{i,j} u_i \otimes u_j \otimes v_j v_i, \\ \sum_{i,(v_i)} u_i l \otimes v_{i(1)}m \otimes v_{i(2)}n &= \sum_{i,j} v_j v_i \otimes u_j \otimes u_j. \end{aligned}$$

Diagrammatically, those conditions correspond to the following ones ((B1) and (B2) respectively).

$$\begin{array}{ccc} R & \xrightarrow{\gamma \otimes \gamma} & A^{\otimes 4} \xrightarrow{\sigma_{1342}} A^{\otimes 4} \\ \gamma \downarrow & & \downarrow 1 \otimes 1 \otimes \mu \\ A^{\otimes 2} & \xrightarrow{\delta \otimes 1} & A^{\otimes 3} \end{array}$$

$$\begin{array}{ccc} R & \xrightarrow{\gamma \otimes \gamma} & A^{\otimes 4} \xrightarrow{\sigma_{3142}} A^{\otimes 4} \\ \gamma \downarrow & & \downarrow \mu \otimes 1 \otimes 1 \\ A^{\otimes 2} & \xrightarrow{1 \otimes \delta} & A^{\otimes 3} \end{array}$$

Thus one can define a braiding element for a bialgebra A to be an invertible element $\gamma \in A \otimes_R A$.

A *braided bialgebra*, or a *quasitriangular bialgebra*, is a bialgebra equipped with a braiding element $\gamma \in A \otimes_R A$. A braiding element γ is called a *symmetry element* when $\gamma^2 = 1 \in A \otimes_R A$. Symmetry elements are in bijection with symmetries on $\mathbf{Mod}_R(A)$. A *symmetric bialgebra*, or a *triangular algebra*, is a bialgebra equipped with a symmetry element.

Let us point out that the conditions (B1) and (B2) can be reworded differently if A is Cauchy as an R -module. In this case, elements $\gamma = \sum_i u_i \otimes v_i \in A \otimes_R A$ are in bijection with R -module morphisms $g : A^* \rightarrow A$ with the following formula:

$$\gamma = (R \xrightarrow{d} A^* \otimes_R A \xrightarrow{g \otimes 1_A} A \otimes_R A)$$

(B1) says that g preserves comultiplication, whilst (B2) says that g preserves multiplication. In fact, if γ is a braiding element, $g : A^* \rightarrow A'^{op}$ is a bialgebra morphism; preservation of unit and counit follows from $c_{M,I} = c_{I,M} = 1_M$.

Let us consider the translation of (B2) to g . Let us start with the following

diagram

$$\begin{array}{ccccc}
R & \xrightarrow{d} & A^* \otimes_R A & \xrightarrow{1 \otimes d \otimes 1} & A^* \otimes_R A^* \otimes_R A \otimes_R A \\
\downarrow d & & & & \downarrow \delta^* \otimes 1 \otimes 1 \\
A^* \otimes_R A & \xrightarrow{1 \otimes \delta} & A^* \otimes_R A \otimes_R A & &
\end{array}$$

for δ , which is the multiplication for A^* . To prove g preserves multiplication is to prove:

$$\begin{array}{ccccc}
A^* \otimes_R A^* & \xrightarrow{g \otimes g} & A \otimes_R A & \xrightarrow{\sigma} & A \otimes_R A \\
d^* \downarrow & & & & \downarrow \mu \\
A^* & \xrightarrow{g} & A & &
\end{array}$$

This is equivalent to proving the legs are equal after applying $_ \otimes_R A \otimes_R A$ and composing with

$$R \xrightarrow{d} A^* \otimes_R A^* \xrightarrow{1 \otimes d \otimes 1} A^* \otimes_R A^* \otimes_R A \otimes_R A.$$

Thus we have:


$$\begin{array}{ccccccc}
R & \xrightarrow{d} & A^* \otimes_R A^* & \xrightarrow{1 \otimes d \otimes 1} & A^* \otimes_R A^* \otimes_R A \otimes_R A & \xrightarrow{g \otimes g \otimes 1 \otimes 1} & A^{\otimes 4} \xrightarrow{\sigma} A^{\otimes 4} \\
& & \downarrow 1 \otimes \delta & & & & \downarrow \mu \otimes 1 \otimes 1 \\
& & A^* \otimes_R A \otimes_R A & \xrightarrow{g \otimes 1 \otimes 1} & A^{\otimes 3} & &
\end{array}$$

Let $\mathcal{V}$ be a braided tensor category, a twist for $\mathcal{V}$ is a family of isomorphisms $\theta_A : A \rightarrow A$ such that $\theta_I = 1_I$ and


$$\begin{array}{ccc}
A \otimes B & \xrightarrow{c_{A,B}} & B \otimes A \\
\theta_{A \otimes B} \downarrow & & \downarrow \theta_B \otimes \theta_A \\
A \otimes B & \xleftarrow{c_{B,A}} & B \otimes A
\end{array}$$

A *balanced tensor category* is a braided monoidal category with a choice of a twist.

Example 10.4. The braid category $\mathcal{B}$ is canonically balanced. The twist $\theta_n : n \rightarrow n$ is obtained by taking n vertical parallel strings with ends tied to two horizontal parallel rods, and rotating the bottom rod through a full 2π twist in the right-hand screw direction with thumb vertical. Then θ_0, θ_1 are identities, while θ_2 is:




Example 10.5. There is a monoidal category $\tilde{\mathcal{B}}$, which is defined similarly to $\mathcal{B}$, except that the arrows are braids on ribbons (instead of strings), and it is permissible to twist the ribbons through full 2π turns:




The homsets $\tilde{\mathcal{B}}(n, n) = \tilde{\mathcal{B}}_n$ are groups under composition. A presentation of this group $\tilde{\mathcal{B}}_n$ is given by generators $s_1, \dots, s_n$, where $s_1, \dots, s_{n-1}$ satisfy the relations from the Artin braid group $\mathcal{B}_n$, whereas $s_1, \dots, s_n$ satisfy the extra relation

$$s_{n-1}s_ns_{n-1}s_n = s_ns_{n-1}s_ns_{n-1}$$

where s_n is depicted as follows:



Composition in $\tilde{\mathcal{B}}$ is vertical stacking of diagrams, and tensor product for $\tilde{\mathcal{B}}$ is horizontal placement of diagrams, much as for $\mathcal{B}$. The braiding $c_{m,n} : m + n \rightarrow n + m$ for $\tilde{\mathcal{B}}$ is obtained by placing the first m ribbons over the remaining n without introducing any twists. Now the twist $\theta_n : n \rightarrow n$ for $\tilde{\mathcal{B}}$ is obtained by regarding the two boundary edges of the ribbons as extra strings and taking $\theta_{2n} : 2n \rightarrow 2n$ in $\mathcal{B}$. Therefore, we have the following $\tilde{\mathcal{B}}$:



Example 10.6. Let A and B be Abelian groups and let $f : A \times A \rightarrow B$ be a bilinear function. Then there is a balanced strict monoidal category $\mathcal{C}_f$ is constructed the following way: the objects are elements of A , the homset $\mathcal{C}_f(x, y)$ is empty unless $x = y$ and $\mathcal{C}_f(x, x) = B$. The tensor product is given by

$$(x \xrightarrow{\alpha} x) \otimes (y \xrightarrow{\beta} y) = (x + y \xrightarrow{\alpha+\beta} x + y).$$

The braiding is $c_{x,y} = f(x, y) : x + y \rightarrow y + x$ and the twist is given by $\theta_X = f(x, x) : x \rightarrow x$.

Example 10.7. Let A be a braided R -algebra with braiding element:

$$\gamma = \sum_{i \in I} u_i \otimes v_i \in A \otimes_R A.$$

A *twist element* for A is an invertible central element $\tau \in A$ such that $\varepsilon(\tau) = 1$

$$\delta(\tau) = \sum_{i,j} (u_i \tau v_j) \otimes (v_i \tau u_j).$$

The above equation is diagrammatically presented the following way:

$$\begin{array}{ccccc} R & \xrightarrow{\gamma \otimes \tau \otimes \tau \otimes \gamma} & A^{\otimes 6} & \xrightarrow{\sigma_{136245}} & A^{\otimes 6} \\ \downarrow \tau & & & & \downarrow \mu \otimes 1 \otimes \mu \otimes 1 \\ & & & & A^{\otimes 4} \\ & & & & \downarrow \mu \otimes \mu \\ A & \xrightarrow{\delta} & A^{\otimes 2} & & \end{array}$$

Twist elements τ for A are in bijection with twists θ for the braided monoidal category $\mathbf{Mod}_R(A)$. Naturality of $\theta_M : M \rightarrow M$ means it has the form of $\theta_M(m) = \tau m$ for some $\tau \in A$. To make θ_M an A -module morphism, τ should be central, i.e. $\tau \cdot a = a \cdot \tau$ for all $a \in A$.

A *balanced algebra* is a braided algebra with a twist.

11 Internal homs and duals

Let $\mathcal{V}$ be a monoidal category and let A and B be objects in $\mathcal{V}$. A *left internal hom* for A and B is an object $[A, B]$ along with the *evaluation arrow* $e_A :$

$[A, B] \otimes A \rightarrow B$ such that for all arrows $f : C \otimes A \rightarrow B$ there exists a unique arrow $\hat{f} : C \rightarrow [A, B]$ such that the following holds:

$$\begin{array}{ccc} [A, B] \otimes A & \xrightarrow{e_A} & B \\ \hat{f} \otimes 1_A \uparrow & \nearrow f & \\ C \otimes A & & \end{array}$$

Thus there is a natural bijection:

$$\begin{aligned} \mathcal{V}(C, [A, B]) &\cong \mathcal{V}(C \otimes A, B) \\ g &\mapsto e_A \circ (g \otimes 1_A) \end{aligned}$$

A monoidal category is *left-closed* if each pair of objects has a left internal hom. If $f : C \rightarrow A$ and $g : B \rightarrow D$ are arrows in $\mathcal{V}$, provided $\mathcal{V}$ is monoidal closed, then there is a unique arrow:

$$[f, g] : [A, B] \rightarrow [C, D]$$

such that:

$$\begin{array}{ccc} [A, B] \otimes C & \xrightarrow{[f, g] \otimes 1} & [C, D] \otimes C \\ 1 \otimes f \downarrow & & \downarrow e_C \\ [A, B] \otimes A & \xrightarrow{e_A} B \xrightarrow{g} & D \end{array}$$

Therefore the internal hom is a functor $[_, _] : \mathcal{V}^{op} \otimes \mathcal{V} \rightarrow \mathcal{V}$.

The internal hom for A and B is unique up to isomorphism. An internal hom for I and A always exists since we have $[I, A] = A$ with $e_A = r_A : A \otimes I \rightarrow A$.

If B, C and $A \otimes B, C$ have internal homs, so do $A, [B, C]$:

$$[A, [B, C]] = [A \otimes B, C]$$

with $a_A = e_{A \otimes B} : [A \otimes B, C] \otimes A \rightarrow [B, C]$.

There is a composition arrow: $[B, C] \otimes [A, B] \rightarrow [A, C]$ in $\mathcal{V}$ (whenever the corresponding internal homs do exist), which corresponds to the following composite:

$$[B, C] \otimes [A, B] \otimes A \xrightarrow{1 \otimes e_A} [B, C] \otimes B \xrightarrow{e_B} C.$$

A right internal hom is defined dually and is denoted as $[A, B]'$. The right evaluation arrow has the form of $e'_A : A \otimes [A, B]' \rightarrow B$.

A monoidal category is *closed* if it has all left and right internal homs.

Example 11.1. The symmetric monoidal category $\mathbf{Mod}_R$ of modules over a commutative ring R is closed:

$$[M, N] = [M, N]' = \text{Hom}_R(M, N)$$

Example 11.2. Let H be an R -Hopf algebra. Then the tensor category $\mathbf{Mod}_R(H)$ of H -modules with tensor product $\otimes_R$ is left-closed:

$$[M, N] = \text{Hom}_R(M, N)$$

Assume the antipode ν for H is invertible, then H' becomes a Hopf algebra with antipode ν^{-1} . Let $\text{Hom}'_R(M, N)$ denote $\text{Hom}_R(M, N)$ as a left H' -module. The forgetful functor $\mathbf{Mod}_R(H) \rightarrow \mathbf{Mod}_R$ preserves tensors and left/right internal homs.

Example 11.3. Let $\mathbf{Ban}$ be the category of Banach spaces over $\mathbb{C}$ and arrows $f : A \rightarrow B$ are linear functions such that

$$\|f(a)\| \leq \|a\|.$$

One can make $\mathbf{Ban}$ into a symmetric tensor category by taking tensor products as vector space. The internal hom exists for all Banach spaces, which is the Banach space of bounded functions with the usual norm. Thus $\mathbf{Ban}$ is a symmetric monoidal category.

Let $\mathcal{V}$ be a monoidal category. An object J is *left dualising* when, for each A , internal homs $[A, J]$ and $[A, J]'$ exist, and the arrow $\omega_A : A \rightarrow [[A, J]', J]$ is invertible. Thus $\mathcal{V}$ is left-closed with

$$[A, B] = [A \otimes [B, J]', J]$$

since

$$\mathcal{V}(C, [A \otimes [B, J]', J]) \cong \mathcal{V}(C \otimes A \otimes [B, J]', J) \cong \mathcal{V}(C \otimes A, [[B, J]', J]) \cong \mathcal{V}(C \otimes A, B).$$

The concept of right dualising object is defined similarly.

Example 11.4. Let $\mathbf{k}$ be a field. A *quadratic algebra* is a pair (V, R) where V is a finite-dimensional vector space and R is a subspace of $V \otimes V$. A quadratic algebra morphism $f : (V, R) \rightarrow (W, S)$ is a linear function $f : V \rightarrow W$ such that

$$(f \otimes f)(R) \subseteq S$$

Write $\mathcal{QA}$ for the category of quadratic algebras. Each quadratic algebra (V, R) determines an actual algebra:

$$T(V)/R$$

where $T(V)$ is the tensor algebra on V . $\mathcal{QA}$ is an SMC with a symmetric tensor product given:

$$(V, R) \otimes (W, S) = (V \otimes W, \sigma_{1324}(R \otimes S))$$

The unit object is $I = (\mathbf{k}, \mathbf{k} \otimes \mathbf{k})$. The claim is that $J = (\mathbf{k}, 0)$ is a dualising object. Observe that:

$$[(V, R), J] = (V^*, R^\perp)$$

where $V^* = \text{Hom}_{\mathbf{k}}(V, \mathbf{k})$ and $R^\perp$ is the kernel of the composite surjection:

$$V^* \otimes V^* \xrightarrow{\cong} (V \otimes V)^* \xrightarrow{i^*} R^*$$

where $i : R \rightarrow V \otimes V$ being inclusion. It follows that R is the kernel of

$$V \otimes V \xrightarrow{\omega} V^{**} \otimes V^{**} \xrightarrow{\cong} (V^* \otimes V^*)^* \xrightarrow{i^*} R^{\perp*}$$

so we have an isomorphism:

$$\omega : (V, R) \cong (V^{**}, R).$$

The quadratic algebra which we identify with the quantum plane $\mathbb{A}_q^{2|0}$ is

$$\mathbb{A}_q^{2|0} = (\mathbf{k}^2, (y \otimes x - qx \otimes y))$$

where $x = (1, 0)$, $y = (0, 1) \in \mathbf{k}^2$. Let $\xi, \eta \in \mathbf{k}^{2*}$ be the dual basis given by $\xi(x) = \eta(y) = 1$, $\xi(y) = \eta(x) = 0$. Then:

$$(y \otimes x - qx \otimes y)^\perp = (\xi \otimes \xi, \eta \otimes \eta, \xi \otimes \eta + q\eta \otimes \xi)$$

as a subspace of $\mathbf{k}^{2*} \otimes \mathbf{k}^{2*}$. Hence the quantum superplane arises as the quadratic algebra:

$$\mathbb{A}_q^{0|2} = (\mathbf{k}^{2*}, (\xi \otimes \xi, \eta \otimes \eta, \xi \otimes \eta + q\eta \otimes \xi)) = [\mathbb{A}_q^{0|2}, J].$$

Notice also that:

$$\mathbb{A}_q^{0|2} \otimes \mathbb{A}_q^{2|0} = (\mathbf{k}^{2*} \otimes \mathbf{k}^2, (b \otimes a - qa \otimes b, d \otimes c - qc \otimes d, q^{-1}b \otimes c - qc \otimes b + d \otimes a - a \otimes d)).$$

Let $\mathcal{V}$ be a monoidal category. Write $d, e : B \dashv A$, or just $B \dashv A$, for $A, B \in \mathcal{V}$ and arrows $e : B \otimes A \rightarrow I, d : I \rightarrow A \otimes B$ such that the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{d \otimes 1_A} & A \otimes B \otimes A \\ & \searrow & \downarrow 1_A \otimes e \\ & & A \end{array} \quad \begin{array}{ccc} B & \xrightarrow{1_B \otimes d} & B \otimes A \otimes B \\ & \searrow & \downarrow e \otimes 1_B \\ & & B \end{array}$$

B is called a *left dual* for A , and A is called a *right dual* for B . e is called *the counit*, d is called *the unit*. Duals are defined uniquely up to isomorphism.

A monoidal category is called *left autonomous* when each $A \in \mathcal{V}$ has a left dual A^* . Each arrow $f : A \rightarrow B$ determines a unique arrow $f^* : B^* \rightarrow A^*$ given by the composite:

$$B^* \xrightarrow{1 \otimes d} B^* \otimes A \otimes A^* \xrightarrow{1 \otimes f \otimes 1} B^* \otimes B \otimes A^* \xrightarrow{e \otimes 1} A^*$$

So one makes left dual into a functor $(_)^* : \mathcal{V}^{op} \rightarrow \mathcal{V}$. We also have $(A \otimes B)^* \cong B^* \otimes A^*$ and $I^* \cong I$. The monoidal category is *autonomous* when each A has both a left dual A^* and right dual $A^\vee$.

If $\mathcal{V}$ is braided, then each left dual A^* is also a right dual with counit and unit:

$$\begin{aligned} A \otimes A^* &\xrightarrow{c_{A,A^*}} A^* \otimes A \xrightarrow{e} I \\ I &\xrightarrow{d} A \otimes A^* \xrightarrow{c_{A,A^*}^{-1}} A^* \otimes A \end{aligned}$$

Therefore, $A \cong A^{**}$.

A *tortile monoidal category* is an autonomous balanced monoidal category, in which twist is related to the dual via the condition:

$$\theta_{A^*} = \theta_A^* : A^* \rightarrow A^*$$

A left dual A^* for A gives a left internal hom $[A, B] = B \otimes A^*$ with evaluation:


$$e_A = 1 \otimes e : B \otimes A^* \otimes A \rightarrow B$$

for each $B \in \mathcal{V}$. A right dual $A^\vee$ gives a right internal hom $[A, B]^\vee = A^\vee \otimes B$. Thus a left/right autonomous monoidal category is left/right-closed.


Example 11.5. Let R be a commutative ring R , an object M of $\mathbf{Mod}_R$ a (left) dual if and only if it is Cauchy. $\mathbf{Prf}_R$ is the full subcategory of $\mathbf{Mod}_R$ consisting of Cauchy R -modules. $\mathbf{Prf}_R$ is closed under $\otimes$, so $\mathbf{Prf}_R$ is an autonomous monoidal category.

Example 11.6. Let H be an R -Hopf algebra. An object M of the monoidal category $\mathbf{Mod}_R(H)$ has a left dual iff it is Cauchy as an R -module; $M^* = \text{Hom}_R(M, R)$. If H has an invertible antipode, then each M has a right dual $M^\vee = \text{Hom}'_R(M, R)$. Write $\mathbf{Prf}_R(H)$ for the full subcategory $\mathbf{Mod}_R(H)$ consisting of those H -modules M which are Cauchy viewed as R -modules. If H has an invertible antipode, then $\mathbf{Prf}_R(H)$ is an autonomous monoidal category.


Example 11.7. Let P be a Euclidean plane. A *geometric tangle* T is a compact 1-dimensional oriented submanifold of $[0, 1] \times P$, which is tamely embedded and whose boundary ∂T is equal to $T \cap \partial([0, 1] \times P)$. Therefore a geometric tangle T is a disjoint union of directed (topological) circles contained in $(0, 1) \times P$ and of directed paths connecting two points on the boundary $\partial([0, 1] \times P)$. The *target* of T is the subset $\partial T \cap (\{1\} \times P)$ as an oriented 0-dimensional manifold. The *source* of T is the subset $\partial T \cap (\{0\} \times P)$, but with orientation reserved. A *geometric tangle* is pictured as follows:



Now we define the autonomous braided monoidal category $\mathcal{T}$ of tangles. The objects are functions $A : \{1, 2, \dots, n\} \rightarrow \{+, -\}$ for $n \geq 0$ called *signed sets*. The arrows of $\mathcal{T}$ are the tangles which have these signed sets as sources and targets. Composition and tensor-product are for braids. The braiding is visualised as follows:



The left dual A^* of a signed set A is given by reversing the order and the sign of the points. That is, $A^*(i)$ and $A(n - i + 1)$ are opposite signs for $1 \leq i \leq n$. The counit and unit are illustrated below for $A = \{-, +, -\}$:



12 Monoidal Functors and Yang-Baxter operators

Let $\mathcal{C}$ and $\mathcal{V}$ be monoidal categories. A *monoidal functor* $F : \mathcal{C} \rightarrow \mathcal{V}$ consists of a functor $F : \mathcal{C} \rightarrow \mathcal{V}$ along with a natural isomorphism $\phi_{A,B} : FA \otimes FB \xrightarrow{\cong}$

$F(A \otimes B)$ and an isomorphism $\varphi_0 : I \xrightarrow{\cong} FI$ such that the following is satisfied:

$$\begin{array}{ccc}
FA \otimes FB \otimes FC & \xrightarrow{\phi_{A,B}} & F(A \otimes B) \otimes FC \\
\downarrow 1 \otimes \phi_{B,C} & & \downarrow \phi_{A \otimes B, C} \\
FA \otimes F(B \otimes C) & \xrightarrow{\phi_{A, B \otimes C}} & F(A \otimes B \otimes C)
\end{array}
\quad
\begin{array}{ccc}
FA & \xrightarrow{1 \otimes \varphi_0} & FA \otimes FI \\
\downarrow \varphi_0 \otimes 1 & \searrow & \downarrow \varphi_{A, I} \\
FI \otimes FA & \xrightarrow{\varphi_{I, A}} & FA
\end{array}$$

If all the ϕ 's are not invertible, then F is a *lax monoidal functor*; if ϕ 's are identities, then F is a *strict monoidal functor*.

If $\mathcal{C}$ and $\mathcal{V}$ are braided (symmetric), then a monoidal functor is braided (symmetric) if the following square commutes:

$$\begin{array}{ccc}
FA \otimes FB & \xrightarrow{\varphi_{A,B}} & F(A \otimes B) \\
\downarrow c_{FA, FB} & & \downarrow c_{FB, FA} \\
FB \otimes FA & \xrightarrow{\varphi_{B,A}} & F(B \otimes A)
\end{array}$$

If $\mathcal{C}$ and $\mathcal{V}$ are balanced, then F is balanced if it is braided and $F(\theta_A) = \theta_{FA} : FA \rightarrow FA$.

Let $F : \mathcal{C} \rightarrow \mathcal{V}$ is a lax monoidal functor and let $\mathcal{C}$ and $\mathcal{V}$ be left-closed, then the composite:

$$F[A, B] \otimes FA \xrightarrow{\varphi_{[A, B], A}} F([A, B] \otimes A) \xrightarrow{Fe_A} FB$$

corresponds to an arrow:

$$\phi_{A,B}^{\sim} : F[A, B] \rightarrow [FA, FB]$$

F is a left-closed monoidal functor whenever each $\phi_{A,B}^{\sim}$ is invertible. Right-closed and closed are defined in a similar fashion.

Note that monoidal functors preserve duals: if $F : \mathcal{C} \rightarrow \mathcal{V}$ is a monoidal functor and $d, e : B \dashv A$ in $\mathcal{C}$, then $FB \dashv FA$ in $\mathcal{V}$ with unit:

$$I \xrightarrow{\phi_0} FI \xrightarrow{Fd} F(B \otimes A) \xrightarrow{\phi_{[A, B], A}^{-1}} FA \otimes FB$$

and counit

$$FB \otimes FA \xrightarrow{\phi_{[B, A]}} F(B \otimes A) \xrightarrow{Fe} FI \xrightarrow{\phi_0^{-1}} I$$

Example 12.1. The category **Set** of sets is a monoidal category with $\otimes = \times$. Let R be a commutative ring, then the free module functor $\mathcal{F}_R : \mathbf{Set} \rightarrow \mathbf{Mod}_R$ is monoidal. But it is not closed since:

$$\mathrm{Hom}_R(\mathcal{F}_R X, \mathcal{F}_R Y) \cong (\mathcal{F}_R Y)^X \not\cong Y^X$$

The functor $|_| : \mathbf{Mod}_R \rightarrow \mathbf{Set}$ taking a module M to its underlying set $|M|$ is an example of a lax monoidal functor since the following functions:

- $\phi_{M,N} : |M| \times |N| \rightarrow |M \otimes N|$ with $(m, n) \mapsto m \otimes n$,
- $\phi_0 = \eta : 1 \mapsto |R|$.

Example 12.2. The universal enveloping algebra functor $\mathcal{U} : \mathbf{Lie}_R \rightarrow \mathbf{Alg}_R$ is a monoidal functor.


Example 12.3. Let $\mathcal{V}$ be a monoidal category. A *Yang-Baxter operator* on an object $A \in \mathcal{V}$ is an invertible arrow $y : A \otimes A \rightarrow A \otimes A$ such that the following hexagon commutes:

$$\begin{array}{ccccc}
 & & A \otimes A \otimes A & \xrightarrow{1 \otimes y} & A \otimes A \otimes A \\
 & \nearrow y \otimes 1 & & & \searrow y \otimes 1 \\
 A \otimes A \otimes A & & & & A \otimes A \otimes A \\
 & \searrow 1 \otimes y & & & \nearrow 1 \otimes y \\
 & & A \otimes A \otimes A & \xrightarrow{y \otimes 1} & A \otimes A \otimes A
 \end{array}$$


For instance, the object 1 of $\mathcal{B}$ admits the following YB-operator:

$$c_{1,1} = s_1 : 1 + 1 \rightarrow 1 + 1$$

which is the element of the braid group $\mathcal{B}_2$:



Then the YB-hexagon becomes the following identity:



As far as monoidal functor $F : \mathcal{B} \rightarrow \mathcal{V}$ preserves tensors up to isomorphism, one obtains a YB-operator y on $F(1) = A$:

$$y : A \otimes A \xrightarrow{\phi_{1,1}} F(1 + 1) \xrightarrow{Fs_1} F(1 + 1) \xrightarrow{\phi_{1,1}^{-1}} A \otimes A$$

Given a YB-operator y on $A \in \mathcal{V}$, one can determine a monoidal functor $F : \mathcal{B} \rightarrow \mathcal{V}$ such that $F(1) = A$ and y is the above composite. Moreover, F is unique up to isomorphism. As far as F is a monoidal functor, we also have:


$$F(n) = F(1 + 1 + \dots + 1) \cong A^{\otimes n}$$

Each generator s_i of $\mathcal{B}_n$ can be written as

$$s_i = \mathbf{1}_{i-1} \otimes s_1 \otimes \mathbf{1}_{n-i-1} \in \mathcal{B}$$

Thus, the definition of $F s_i : A^{\otimes n} \rightarrow A^{\otimes n}$ is forced.

Example 12.4. A tensor functor from ribbons $F : \tilde{\mathcal{B}} \rightarrow \mathcal{V}$ determines a YB-operator y on $F1 = A$ as in the above example. The twist $\theta_1 : 1 \rightarrow 1$ in $\tilde{\mathcal{B}}$ gives an isomorphism $z = F\theta_1 : A \rightarrow A$. Due to the following equalities in $\tilde{\mathcal{B}}$:



A YB-operator y on an object $A \in \mathcal{V}$ is *balanced* when it is equipped with an isomorphism $z : A \rightarrow A$ such that the following squares commute:

$$\begin{array}{ccc} A \otimes A & \xrightarrow{y} & A \otimes A \\ \downarrow 1 \otimes z & & \downarrow z \otimes 1 \\ A \otimes A & \xrightarrow{y} & A \otimes A \end{array} \quad \begin{array}{ccc} A \otimes A & \xrightarrow{y} & A \otimes A \\ \downarrow z \otimes 1 & & \downarrow 1 \otimes z \\ A \otimes A & \xrightarrow{y} & A \otimes A \end{array}$$

Each balanced YB-operator determines a unique monoidal functor $F : \tilde{\mathcal{B}} \rightarrow \mathcal{V}$ from which A , y and z are recovered. Therefore, $(\tilde{\mathcal{B}}, 1, c_{1,1}, \theta_1)$ is the free tensor category containing an object equipped with a balanced YB-operator.

Let us recall the following observations:

- Monoidal functors take YB-operators to YB-operators.
- A braiding on each monoidal category gives a YB-operator $c_{A,A} : A \otimes A \rightarrow A \otimes A$.

Besides, monoidal functors take balanced YB-operators to balanced YB-operators.

A YB-operator y on A is called (*left-*) *dualisable* when A has a left dual A^* and both the arrows $u, v : A^* \otimes A \rightarrow A^*$ are given by the composites:

$$A^* \otimes A \xrightarrow{1 \otimes 1 \otimes d} A^* \otimes A \otimes A \otimes A^* \xrightarrow[1 \otimes y^{-1} \otimes 1]{1 \otimes y \otimes 1} A^* \otimes A \otimes A \otimes A^* \xrightarrow{e \otimes 1 \otimes 1} A \otimes A^*$$

are invertible. It follows that the composite w given by:

$$A^* \otimes A^* \xrightarrow{1 \otimes 1 \otimes d} A^* \otimes A^* \otimes A \otimes A^* \xrightarrow{1 \otimes u \otimes 1} A^* \otimes A \otimes A^* \otimes A^* \xrightarrow{e \otimes 1 \otimes 1} A^* \otimes A^*$$

has inverse w^{-1} given by the following composite:

$$A^* \otimes A^* \xrightarrow{1 \otimes 1 \otimes d} A^* \otimes A^* \otimes A \otimes A^* \xrightarrow{1 \otimes v \otimes 1} A^* \otimes A \otimes A^* \otimes A^* \xrightarrow{e \otimes 1 \otimes 1} A^* \otimes A^*$$

Proposition 12.1. In a braided monoidal category, if an object A has a dual, then $y = c_{A,A}$ is a dualisable YB-operator on A with:

$$u = c_{A,A^*}^{-1}, v = c_{A^*,A}^{-1}, w = c_{A^*,A^*}^{-1}.$$

Proof. $c_{A,A^*}^{-1} \circ u = 1_{A^* \otimes A}$, it suffices to show that equality holds after applying $A \otimes _$ to both sides and composing with $d \otimes 1_A$. The following diagram gives the first equation:

$$\begin{array}{ccccc}
A \otimes A^* \otimes A \otimes A \otimes A^* & \xrightarrow{1 \otimes 1 \otimes c_{A,A} \otimes 1} & & & A \otimes A^* \otimes A \otimes A \otimes A^* \\
\uparrow 1 \otimes 1 \otimes 1 \otimes d & \swarrow d \otimes 1 \otimes 1 & & \searrow d \otimes 1 \otimes 1 & \downarrow 1 \otimes e \otimes 1 \otimes 1 \\
A \otimes A^* \otimes A & & A \otimes A \otimes A^* & \xrightarrow{c_{A,A} \otimes 1} & A \otimes A \otimes A^* \\
\uparrow d \otimes 1 & \nearrow 1 \otimes d & & \searrow c_{A,A \otimes A^*} & \downarrow 1 \otimes c_{A,A^*} \\
A & \xrightarrow{d \otimes 1} & A & & A \otimes A^* \otimes A
\end{array}$$

The other equalities are shown similarly. $\square$

Monoidal functors preserve dualisability: if $\mathcal{C}$ is braided and X has a dual in $\mathcal{V}$, we obtain a dualisable YB-operator FX in $\mathcal{V}$.

A YB-operator on A is called *tortile* when it is balanced, dualisable and the following diagram commutes:

$$\begin{array}{ccccc}
& & A \otimes A \otimes A^* & & \\
& \nearrow y \otimes 1 & & \searrow 1 \otimes v^{-1} & \\
A \otimes A \otimes A^* & & & & A \otimes A^* \otimes A \\
\nearrow 1 \otimes d & & & & \searrow 1 \otimes e \\
A & \xrightarrow{z} & A & \xrightarrow{z} & A
\end{array}$$

Proposition 12.2. In a balanced monoidal category, if an object A has a dual then the pair $(c_{A,A}, \theta_A)$ is a tortile YB-operator precisely when:

$$\theta_{A^*} = (\theta_A)^*.$$

Proof. The following equation:

$$(\theta_{A^*})^* \theta_A = (1 \otimes e) \circ (1 \otimes v^{-1}) \circ (y \otimes 1) \circ (1 \otimes d)$$

is proved by the following diagram:

$$\begin{array}{ccccccc}
A & \xrightarrow{\theta_A} & A & \xrightarrow{1 \otimes d} & A \otimes A \otimes A^* & \xrightarrow{1 \otimes c_{A^*,A}^{-1}} & A \otimes A^* \otimes A \\
\downarrow 1 \otimes d & & & & & \nearrow \theta_A \otimes 1 \otimes 1 & \downarrow 1 \otimes \theta_{A^*} \otimes 1 \\
A \otimes A \otimes A^* & \xrightarrow{1 \otimes c_{A^*,A}^{-1}} & A \otimes A^* \otimes A & \xrightarrow{\theta_A \otimes \theta_{A^*} \otimes 1} & A \otimes A^* \otimes A & & \\
\downarrow y \otimes 1 = c_{A,A} \otimes 1 & \searrow c_{A^*,A}^{-1} & \swarrow c_{A^*,A}^{-1} \otimes 1 & & & & \\
& A^* \otimes A \otimes A & \xrightarrow{\theta_{A^*} \otimes A \otimes 1} & A^* \otimes A \otimes A & & & \\
& \downarrow 1 \otimes c_{A,A} & \searrow e \otimes 1 & \swarrow c_{A^*} \otimes A, A & & & \\
& A^* \otimes A \otimes A & \xrightarrow{c_{A^*,A} \otimes 1} & A & \xlongequal{\quad} & A & \\
& \swarrow c_{A^*,A}^{-1} \otimes A \otimes A & \searrow c_{A^*,A} \otimes 1 & & & & \\
A \otimes A \otimes A^* & \xrightarrow{1 \otimes v^{-1} = 1 \otimes c_{A^*,A}^{-1}} & A \otimes A^* \otimes A & \xrightarrow{1 \otimes e} & A & & \\
& & & & & & \downarrow e \otimes 1 \\
& & & & & & A
\end{array}$$

Thus the balanced YB-operator (y, z) is tortile if and only if $(\theta_{A^*})^* \theta_A = z^2$, i.e., if and only if

$$(\theta_{A^*})^* \theta_A = (\theta_A)^2 \Leftrightarrow (\theta_{A^*})^* = \theta_A \Leftrightarrow \theta_{A^*} = (\theta_A)^*.$$

□

Therefore, in a tortile monoidal category each object A is equipped with a tortile YB-operator $(c_{A,A}, \theta_A)$.

Example 12.5. As far as $\tilde{\mathcal{T}}$ is a tortile monoidal category, we obtain a tortile YB-operator $(c_{+,+}, \theta_+)$ on the object $+$ of $\tilde{\mathcal{T}}$. Thus, each monoidal functor $F : \tilde{\mathcal{T}} \rightarrow \mathcal{C}$ yields a tortile YB-operator on $F(+)$ in $\mathcal{V}$.

$(\tilde{\mathcal{T}}, +, c_{+,+}, \theta_+)$ is the free monoidal category equipped with a tortile YB-operator. This means the following thing. Let (y, z) be a tortile YB-operator on

A in a monoidal category $\mathcal{V}$, then there is a unique up to isomorphism monoidal functor $F : \tilde{\mathcal{T}} \rightarrow \mathcal{V}$ such that:

$$F : (+, c_{+,+}, \theta_+) \mapsto (A, y, z).$$

This is how F is defined in terms of A, y, z . For example:

- $F(+ - - - + -) = A \otimes A^* \otimes A^* \otimes A^* \otimes A \otimes A^*$,
- $F(c_{+,-}) = u^{-1}$,
- $F(c_{-,-}) = w$,
- $F(\theta_+) = z$,
- $F(\theta_-) = z^*$.
- etc.

Any tangle of ribbons can be decomposed, using composition and tensor product in $\tilde{\mathcal{T}}$, into single crossings ($c_{+,+}$, $c_{+,-}$, $c_{-,+}$, $c_{-,-}$ and their inverses), turnings e and d and twists (θ_+ , θ_- and their inverses). Thus the value of F is forced.