h-adic topology and profinite linear lambda terms

Daniel Rogozin

Contents

1 Profinite Lambda terms and related concepts **1**
1.1 The Explicit Description of Profinite Topology 3
1.2 Profinite Lambda Terms 4
1.3 The cartesian closed category of profinite lambda terms 6

2 h-adic topology **6**

3 On Dialogue Categories and their internal language **8**
3.1 Term Assignment for Tensorial Logic 9
3.2 Parametricity for Tensorial Logic 10

1 Profinite Lambda terms and related concepts

One starts with the observation that if we have a work over a finite alphabet Σ:

$$W = a_{w_1} \dots a_{w_n} \in \Sigma^*$$

then one can represent W as a well-typed term in the simply typed lambda calculus:

$$a_{w_1} : q \to q, \dots, a_{w_n} : q \to q \vdash W : q \to q$$

where q is a base type that we informally interpret as a type of states, so we have a closed lambda term

$$\lambda a^{q\to q}_{w_1} \dots \lambda a^{q\to q}_{w_n} \lambda x^q . a_{w_k}(\dots(a_{w_1}\ q)\dots) : (q \to q) \dots (q \to q) \to q \to q$$

encoding a word w via the instance of the Church encoding. Further, we will call the type of such terms Church_Σ. Further, we have the monoid isomorphism:

$$\Sigma^* \cong \{M \text{ is a lambda term } | M : \mathrm{Church}_\Sigma \}/\beta\eta$$

In other words, words over a finite alphabet can be viewed as lambda terms of a particular type factorised module the $\beta\eta$-equivalence. Generally, there is amount literature studying both syntactic and semantic aspects of the Church encoding of the finite words.

The claim made in [vGMM23] connects the finitary interpretation of STLC in the category of finite sets **FinSet** with deterministic finite automata as follows. Take any finite set Q, so for any type A, we define a set $[\![A]\!]_Q$ inductively defined as

- $[\![p]\!]_Q = Q$

- $[\![A \to B]\!]_Q = [\![A]\!]_Q \Rightarrow [\![B]\!]_Q$

so every term $x_1 : A_1, \ldots, x_n : A_n \vdash M : A$ transports to the function between finite sets $[\![M]\!] : [\![A_1]\!]_Q \times \ldots \times [\![A_n]\!]_Q \to [\![A]\!]_Q$. On closed terms, the interpretation induces a function $[\![.]\!]_Q : \Lambda_{\beta\eta}(A) \to [\![A]\!]_Q$ mapping every closed term M of type A to its interpretation $[\![M]\!]_Q$. Further, a word $W = a_{w_1} \ldots a_{w_k} \in \Sigma^*$ is mapped to the following morphism in **FinSet**: $[\![W]\!]_Q : (Q \Rightarrow Q) \times \ldots \times (Q \Rightarrow Q) \to (Q \Rightarrow Q)$, i.e.,

$$[\![W]\!]_Q : f_{w_1}, \ldots, f_{w_k} \mapsto f_{w_k} \circ \ldots \circ f_{w_1}$$

Recall that a *deterministic finite state automaton* on a finite alphabet $\Sigma = \{a_1, \ldots, a_n\}$ is a structure $\mathcal{A} = (Q, \delta, q_0, Acc)$, where Q is a finite set of states, $\delta : \Sigma \times Q \to Q$ is a transition function, $q_0 \in Q$ is the initial state and $Acc \subseteq Q$ is the set of accepting states. The transition function δ induces a family of functions $\delta_{a_i}(q_k) = \delta(a_i, q_k) : Q \to Q$ for each $a_i \in \Sigma$ and $q_k \in Q$. $\delta_{a_i}(q_k) = q_{k'}$ means that $\mathcal{A}$ transitions from q_k to q'_k encountering a_i. Now one can apply the interpretation $[\![W]\!]_Q$ to the functions $\delta_{a_{w_i}}$ for $1 \leq i \leq k$, so we obtain

$$\delta_w : [\![W]\!]_Q(\delta_{a_{w_1}}, \ldots, \delta_{a_{w_k}}) : Q \to Q$$

Therefore, given a deterministic finite automaton $\mathcal{A} = (Q, \delta, q_0, Acc)$, the tuple (Q, δ, q_0) induces the evaluation function

$$\mathrm{eval}_{\delta, q_0} : [\![\mathrm{Church}_\Sigma]\!]_Q \to Q$$

transporting every functional $F \in [\![\mathrm{Church}_\Sigma]\!]_Q$ to the state $F(\delta_{a_{w_1}}, \ldots, \delta_{a_{w_k}})$ in Q, therefore we have

$$\Sigma^* \cong \Lambda_{\beta\eta}(\mathrm{Church}_\Sigma) \to [\![\mathrm{Church}_\Sigma]\!]_Q \xrightarrow{\mathrm{eval}_{\delta, q_0}} Q$$

so we associate every finite word $w \in \Sigma^*$ with the final state $q_f = \delta_w(q_0)$. The inverse image $Acc \subseteq Q$ under this composite is called a *regular language* $L_\mathcal{A}$, the set of finite words recognisable by $\mathcal{A}$.

Regular languages $L_\mathcal{A}$ form a Boolean algebra $\mathrm{Reg}_Q(\mathrm{Church}_\Sigma)$ of regular languages of lambda terms of type Church_Σ recognised by Q. The algebra can be defined as the image of the Boolean algebra homomorphism $[\![.]\!]_Q^{-1} : 2^{[\![\mathrm{Church}_\Sigma]\!]} \to 2^{\Sigma^*}$ obtained by applying the contravariant powerset functor $2^{(.)} : \mathbf{FinSet}^{op} \to \mathbf{BA}$ to the semantic bracket $[\![.]\!]_Q$. The Boolean algebra $\mathrm{Reg}_Q(A)$ of regular languages of lambda terms of any higher order type A recognisable by a finite set Q is defined as the image of the Boolean algebra homomorphism $[\![.]\!]_Q^{-1} : 2^{[\![A]\!]_Q} \to 2^{\Lambda_{\beta\eta}(A)}$. That is, a set L of lambda terms of type A is recognisable by the finite set Q exactly when it has the form $[\![.]\!]_Q^{-1}(\mathrm{Acc}) = \{M \in \Lambda_{\beta\eta}(A) \,|\, [\![M]\!]_Q \in \mathrm{Acc}\}$ for some choice $\mathrm{Acc} \subseteq [\![A]\!]_Q$ of accepting elements. Thus, the collection $\mathrm{Reg}(A) \subseteq 2^{\Lambda_{\beta\eta}(A)}$ of lambda terms of type A as follows

$$\mathrm{Reg}(A) = \bigcup \{\mathrm{Reg}_Q(A) \mid Q \in \mathbf{FinSet}\}.$$


The monoid $\widehat{\Sigma^*}$ is the free profinite monoid generated by Σ. $\widehat{\Sigma^*}$ can be constructed as the limit, computed in the category of monoids **Mon**, of the projective system of finite monoid homomorphisms

$$(\Sigma^*/\phi \to \Sigma^*/\phi')_{\phi \subseteq \phi'}$$

where ϕ and ϕ' are finite index congruences on the free monoid Σ^* such that $\phi \subseteq \phi'$. Every finite index congruence ϕ, in turn, induces a surjective homomorphism $h : \Sigma^* \to M$ for the quotient monoid $M = \Sigma^*/\phi$. The surjectivity condition on h can be relaxed to show that $\widehat{\Sigma^*}$ of profinite words is the codirected limit of the composite functor

$$\Sigma^*/\mathbf{FinMon} \xrightarrow{\pi} \mathbf{FinMon} \hookrightarrow \mathbf{Mon}$$

$\Sigma^*/\mathbf{FinMon}$ is the slice category whose objects (M, h) are pairs consisting of a finite monoid M and a homomorphism $h : \Sigma^* \to M$ and whose morphisms $(M, h) \to (M', h')$ are homomorphisms $f : M \to M'$ such that the following triangle commutes:



The projection functor $\pi : \Sigma^*/\mathbf{FinMon} \to \mathbf{FinMon}$ transports (M, h) to its underlying monoid M. One obtains $\widehat{\Sigma^*}$ as the limit of a codirected diagram of monoid homomorphisms $(M \to M')_{(M,h)\to(M',h')}$ extending the diagram formed from the family $(\Sigma^*/\phi \to \Sigma^*/\phi')_{\phi\subseteq\phi'}$ to all homomorphisms $h : \Sigma^* \to M$ to a finite monoid M.

Recall that every homomorphism $h : \Sigma^* \to M$ induces a deterministic finite automaton with $Q := M$, $\delta(a, q) = q \cdot_M a$ for every $q \in M$ and $a \in \Sigma$. So every homomorphism $h : \Sigma^* \to M$ (assume $M = \{q_1, \ldots, q_m\}$) induces a decomposition of Σ^* into m components $L_{q_i} = h^{-1}(q_i)$ for $1 \leq i \leq m$ where each L_{q_i} is a regular language. Let $\mathrm{Reg}_{(M,h)}(\Sigma)$ denote the Boolean algebra of languages of the form $L_q = h^{-1}(q)$ for $q \in M$. We have a functor:

$$\mathrm{Reg}_{(.)}(\Sigma) : (\Sigma^*/\mathbf{FinMon})^{op} \to \mathbf{BA}$$

The Boolean algebra $\mathrm{Reg}_{(M,h)}(\Sigma)$ coincides with 2^h, the image of $h : \Sigma^* \to M$ applied to the contravariant powerset functor. Also, the following directed diagram in $\mathbf{BA}$ associated to $\mathrm{Reg}_{(.)}(\Sigma)$:

$$(\mathrm{Reg}_{(M',h')}(\Sigma) \to \mathrm{Reg}_{(M,h)}(\Sigma))_{(M,h)\to(M',h')}$$

can be obtained by applying the contravariant powerset functor to the above trianlge and $(M \to M')_{(M,h)\to(M',h')}$.

1.1 The Explicit Description of Profinite Topology

This description is based on [Pin10, C.X.2]. Let Σ be a finite alphabet and let M be a finite monoid, given $u, v \in \Sigma^*$, a monoid homomorphism $f : \Sigma^* \to M$ *separates* u and v if $f(u) \neq f(v)$. A monoid M separates u and v if there is a homomorphism $\Sigma^* \to M$ separating u and v. The important fact is that every u and v are separated by some monoid M.

A metric on Σ^* is defined as follows: two words are closed if they are separated by the large monoid. Take $u, v \in \Sigma^*$, put:

- $r(u, v) = \min\{|M| \mid M \text{ separates } u \text{ and } v\}$,
- $d(u, v) = 2^{-r(u,v)}$.

Therefore, $d(u, v) < 2^n$ iff u and v cannot be separated by a monoid smaller than n and $d(u, v) < \varepsilon$ iff u and v cannot be separated by a monoid of size n, where n is the unique natural number such that $2^{-(n+1)} < \varepsilon \leq 2^{-n}$. d is a metric on Σ^* such that (Σ^*, d) is an ultrametric space such that, for each $u, v, w \in \Sigma^*$:

- $d(uw, vw) \leq d(uv)$,
- $d(wu, wv) \leq d(uv)$.

As a topological space, (Σ^*, d) is discrete, but open balls are of more interest. For any $n \geq 2$, consider the congruence relation $\sim_n$ on Σ^* defined as:

$$u \sim_n v \Leftrightarrow \forall M\ (|M| \leq n \to (\forall f : \Sigma^* \twoheadrightarrow M) \to f(u) = f(v))$$

Clearly $\sim_n$ is a congruence of finite index as they are finitely many surjections $\Sigma^* \twoheadrightarrow M$ for a fixed finite M. Thus $d(u, v) < 2^{-n}$ iff they cannot be separated by a monoid of size smaller than n.

Proposition 1.1. Let Σ be a finite alphabet,

1. The concatenation function $(u, v) \mapsto uv$ is uniformly continuous,
2. Every open ball of (Σ^*, d) is a recognisable language and every recognisable language is a finite union of open balls.
3. (Σ^*, d) is totally bounded: for every $n > 0$, Σ^* is covered by a finite number of open balls of radius strictly smaller than 2^{-n}.

Proposition 1.2. Let Σ_1 and Σ_2 be finite alphabets, then $f : \Sigma_1 \to \Sigma_2$ is uniformly continuous iff for every recognisable language of Σ_2 L, $f^{-1}(L)$ is also recognisable.

Let $\widehat{\Sigma^*}$ denote the completion of Σ^*, whose points are called profinite words of Σ.

Proposition 1.3. Let M be a finite monoid and let $f : \Sigma^* \to M$ be uniformly continuous, then f uniquely extends to $\widehat{f} : \widehat{\Sigma^*} \to M$.

The following is the universal property of the free profinite monoid:

Proposition 1.4. Let Σ be a finite alphabet, then if $f : \Sigma \to M$ is a function to a finite monoid M, there exists a unique uniformly continuous monoid morphism $\widehat{f} : \widehat{\Sigma^*} \to M$ such that for each $a \in \Sigma$, $\widehat{f}(a) = f(a)$. $\widehat{f}$ is onto iff $f(\Sigma)$ generates M.

Proposition 1.5. Let Σ_1 and Σ_2 be finite alphabets, then every uniformly continuous morphism $f : \Sigma_1^* \to \Sigma_2^*$ uniquely extends to uniformly continuous $\widehat{f} : \widehat{\Sigma_1^*} \to \widehat{\Sigma_2^*}$.

The metric on $\widehat{\Sigma^*}$ is defined similarly. Let M be a finite monoid, then a morphism $f : \Sigma^* \twoheadrightarrow M$ separates profinite words $u, v \in \widehat{\Sigma^*}$ if $\widehat{f}(u) = \widehat{f}(v)$. For $u, v \in u, v \in \widehat{\Sigma^*}$, we set:

- $r(u, v) = \min\{|M| \mid M \text{ separates } u \text{ and } v\}$,
- $d(u, v) = 2^{-r(u,v)}$.

Proposition 1.6. The profinite topology on $\widehat{\Sigma^*}$ is the least topology making every continuous every morphism Σ^* onto a finite discrete monoid.

1.2 Profinite Lambda Terms

One moves from profinite words to profinite lambda terms in the following way. For any finite set Q, one has the interpretation of any type A as the finite set $[\![A]\!]_Q$. Given another finite set Q' with a relation $R \subseteq Q \times Q'$, one can relate different interpretations of A by constructing a relation $[\![A]\!]_R \subseteq [\![A]\!]_Q \times [\![A]\!]_{Q'}$. Such relations are called *logical relations*. Lambda terms are parametric: that is, if M has type A and $R \subseteq Q \times Q'$, then $([\![M]\!]_Q, [\![M]\!]_{Q'}) \in [\![A]\!]_R$.

Now let us discuss how one can define the exponential of relations. Let $S \subseteq P \times P'$ and $R \subseteq Q \times Q'$ be relations on finite sets, the exponential $S \Rightarrow R \subseteq (P \Rightarrow Q) \times (P' \Rightarrow Q')$ is given as:

$$S \Rightarrow R := \{(g, h) \mid \forall x \in P \forall y \in P' \; (xSy \to g(x)Rh(y))\}$$

Therefore, for any relation $R \subseteq Q \times Q'$ between finite sets Q and Q', one constructs a relation $[\![A]\!]_R \subseteq [\![A]\!]_Q \times [\![A]\!]_{Q'}$ defined by induction on A as follows:

- $[\![p]\!] = R$,
- $[\![A \Rightarrow B]\!]_R = [\![A]\!]_R \Rightarrow [\![B]\!]_R$

The *fundamental lemma of logical relations* states that for any $M \in \Lambda_{\beta\eta}(A)$ for any A and for any $R \subseteq Q \times Q'$, the interpretations of M are related as follows:

$$([\![M]\!]_Q, [\![M]\!]_{Q'}) \in [\![A]\!]_R$$

Further, we will be using partial surjections since they are stable under taking relational exponentials. Also, if $f : Q \twoheadrightarrow Q'$ is a partial surjection, so $[\![A]\!]_f \subseteq [\![A]\!]_Q \times [\![A]\!]_{Q'}$ for any A. Besides, we have the inclusion of Boolean algebras $\mathrm{Reg}_{Q'}(A) \subseteq \mathrm{Reg}_Q(A)$, which is proved using the fundamental lemma of logical relations.

Theorem 1.1. The diagram of Boolean algebras $\mathrm{Reg}_{(.)}(A) : \mathbf{FinPSurj}^{op} \to \mathbf{BA}$:

$$(\mathrm{Reg}_{Q'}(A) \hookrightarrow \mathrm{Reg}_Q(A))_{f:Q \to Q' \in \mathbf{FinPSurj}}$$


is directed, and its colimit in **BA** coincides with the Boolean algebra $\mathrm{Reg}(A)$ of regular languages for a higher-order type A.

Also, for any simple type A, the set of regular languages $\mathrm{Reg}(A)$ is closed under finite intersections.

Now let us discuss how the space of profinite lambda terms is constructed. The idea is that for any simple type A as parametric families of semantic elements, so they form a Stone space dual to the Boolean algebra $\mathrm{Reg}(A)$.

Proposition 1.7. Let Q, Q' be finite sets,

1. The set of atoms $\mathrm{At}_Q(A)$ of the finite Boolean algebra $\mathrm{Reg}_Q(A)$ is in bijection with the set $[\![A]\!]^\bullet_Q$ of definable elements of simple type A, given by the function $q \mapsto [\![.]\!]_Q^{-1}(\{q\}) : \mathrm{At}_Q(A) \to [\![A]\!]^\bullet_Q$.
2. For partial surjection $f : Q \twoheadrightarrow Q'$ and for any simple type A, the set $[\![A]\!]^\bullet_Q \subseteq [\![A]\!]_Q$ of definable elements is contained in the domain of $[\![A]\!]_f$, and the restriction of $[\![A]\!]_f$ to $[\![A]\!]^\bullet_Q$ is the unique function $p_{Q,Q'}$ such that the following triangle commutes:



Note that if $f, g : Q \to Q'$ are partial surjections, then, while the semantic interpretations $[\![A]\!]_f, [\![A]\!]_g : [\![A]\!]_Q \to [\![A]\!]'_Q$ are different, their restrictions to $[\![A]\!]^\bullet_Q$ are both equal to $p_{Q,Q'}$.

One defines profinite lambda terms of simple type A as follows.

Definition 1.1. Let A be a simple type, the set of *profinite lambda terms* is defined as the limit $\widehat{\Lambda}_{\beta\eta}(A)$ in **Set** of the diagram:

$$([\![A]\!]^\bullet_f : [\![A]\!]^\bullet_Q \twoheadrightarrow [\![A]\!]^\bullet_{Q'})_{f:Q\twoheadrightarrow Q'\in\mathbf{FinPSurj}}$$

In particular, by the definition of a limit in **Set**, it means that a profinite lambda term in $\widehat{\Lambda_{\beta\eta}(A)}$ is a family θ of definable terms $\theta_Q \in [\![A]\!]_Q$ for every finite set Q such that for every partial surjection $f : Q \twoheadrightarrow Q'$ one has $[\![A]\!]^\bullet_f(\theta_Q) = \theta_{Q'}$. By Proposition 1.7, the above condition is equivalent to the condition that for every term $M : A$ for simple A and any finite sets Q and Q':

$$\theta_Q = [\![M]\!]_Q \wedge |Q'| \leq |Q| \Rightarrow \theta_{Q'} = [\![M]\!]_{Q'}$$

$\widehat{\Lambda_{\beta\eta}(A)}$ is equipped with a natural topology making it a Stone dual of the Boolean algebra $\mathrm{Reg}(A)$. One can define this topology as inherited from the inclusion

$$\widehat{\Lambda_{\beta\eta}(A)} \hookrightarrow \prod_Q [\![A]\!]^\bullet_Q$$

into the product space $\prod_Q [\![A]\!]^\bullet_Q$ where each $[\![A]\!]^\bullet_Q$ is considered a space with the discrete topology. Concretely, for any finite set Q and $q \in [\![A]\!]^\bullet_Q$, let $U_{Q,q}$ denote the set of profinite lambda terms taking value at Q:

$$U_{Q,q} := \{\theta \in \widehat{\Lambda_{\beta\eta}(A)} \mid \theta_Q = q\}$$

Now we take the collection $U_{Q,q}$ as the basis on $\widehat{\Lambda_{\beta\eta}(A)}$ for any Q and q.

The following is similar to [**?**, Section 4.2]:

Proposition 1.8. The space $\widehat{\Lambda_{\beta\eta}(A)}$ is a Stone dual to the Boolean algebra $\mathrm{Reg}(A)$.

1.3 The cartesian closed category of profinite lambda terms

2 h-adic topology

The following summary is based on [Kas12, Chapter XVI].

Consider the complex algebra $K = \mathbb{C}[[h]]$ of complex formal series in one variable h. Every element $f \in \mathcal{K}$ has the form

$$f = \sum_{n=0}^{\infty} a_n h^n$$

where $a_n \in \mathbb{C}$ for each $n \in \mathbb{N}$. Every polynomial $f \in \mathbb{C}[h]$ is clearly a formal series. K form a commutative ring with operations given as

- $f + g = \sum\limits_{n=0}^{\infty} a_n h^n + \sum\limits_{n=0}^{\infty} b_n h^n = \sum\limits_{n=0}^{\infty} (a_n + b_n) h^n$
- $f \cdot g = (\sum\limits_{n=0}^{\infty} a_n h^n) \cdot (\sum\limits_{n=0}^{\infty} b_n h^n) = \sum\limits_{n=0}^{\infty} (\sum\limits_{n=p+q} a_p b_q) h^n$

A formal series f is invertible iff its every coefficient is non-zero.

For any $n > 0$, consider the algebra $K_n = \mathbb{C}[h]/(h^n)$ of truncated polynomials obtained as the quotient of the ring of complex polynomials generated by the ideal (h^n). There is a morphism

$$\pi_n : K \to K_n$$

sending each formal series $\sum_{n=0}^{\infty} a_n h^n$ to the class of $\sum_{k=0}^{n-1} a_k h^k$ modulo (h^n). π_n is onto and its kernel is the ideal $h^n K$ generated by h^n in the ring of formal series. Therefore, we have the following isomorphism:

$$\mathbb{C}[[h]]/(h^n) \cong \mathbb{C}[h]/(h^n)$$

For $n > 0$, there is a surjective map of algebras $p_n : K_n \to K_{n-1}$ generated by the inclusion of ideals $(h^n) \subset (h^{n-1})$. So we have the inverse system of algebras $(K_n, p_n)_{n>0}$ and its inverse limit $\varprojlim_n K_n$. One has $p_n \circ \pi_n = \pi_{n-1}$ for each n. Also, there is a unique morphism of algebras from K to $\varprojlim_n K_n$ whose composition with the projection of the inverse limit is equal to π_n. Besides, the map $\pi : \mathbb{C}[[h]] \twoheadrightarrow \varprojlim_n \mathbb{C}[[h]]/(h^n)$ is the isomorphism, which is partly the instance of Birkhoff's subdirect representation theorem, [SB81, Theorem 8.6].

Let us equip K with the *h-adic topology*. The family of of subsets $\pi_n^{-1}(U_n)$ for $n > 0$ and $U_n \subseteq K_n$ form an open basis on K. $\{0\}$ is a family of open neighbourhoods in the discrete set K_n, the family $\pi_n^{-1}(0)$ is a family of open neighbourhoods of 0 in K for the h-adic topology. There is also the structure of a metric space on K given as follows. Given a formal series $f = \sum_{n=0}^{\infty} a_n h^n$, let $o(f)$ denote the unique natural number such that $a_{o(f)} \neq 0$ and $a_k = 0$ for each $k < o(f)$. For $f = 0$, set $o(f) = +\infty$ and extend $(\mathbb{N}, <)$ by adding the top element $+\infty$. It is easy to see that:

$$(h^n) = \{f \in K \mid o(f) > n - 1\}$$

and therefore

$$\bigcap_{n>0} (h^n) = \{0\}$$

We also have for any series $f, g \in K$:

$$o(f + g) \geq \min(o(f), o(g))$$

Now define $|.| : K \to \mathbb{R}_{\geq 0}$ as

$$|f| = 2^{-o(f)}$$

Then if we put $d(f, g) = |f - g|$, then $(\mathbb{C}[[h]], d)$ becomes an ultrametric space.

Let M be a left K-module, consider the family $(h^n M)_{n>0}$ of submodules of M with the canonical K-linear projections:

$$p_n : M_n = M/h^n M \to M_{n-1} = M/h^{n-1} M$$

They form an inverse system of K-modules, so we can consider the inverse limit

$$\widetilde{M} = \varprojlim_n M_n$$

having a natural structure of a left K-module. $\widetilde{M}$ has a natural topology as well, the inverse limit topology, for which the family of submodules $(h^n \widetilde{M})_n$ form an open basis. The projections

$i_n : M \to M_n$ induce a unique K-linear map $i : M \to \widetilde{M}$ such that $\pi_n \circ i = i_n$ for each n. The kernel of i is given as

$$\mathrm{Ker}(i) = \bigcap_{n>0} h^n M$$

A K-module M is *separated* if $\mathrm{Ker}(i) = \{0\}$. It is *complete* if i is injective.

For any module K, the module $M/\mathrm{Ker}(i)$ is separated and the completion $\widetilde{M}$ of M is complete.

Generally, any separated complete K-module M is equipped with the h-adic topology induced by the inverse limit topology on $\widetilde{M}$ via $M \cong \widetilde{M}$.

Now let us describe the following class of separated complete K-modules. Let V be a vector space over $\mathbb{C}$, let $V[[h]]$ denote the set of all formal series in variable h

$$f = \sum_{n=0}^{\infty} a_n h^n$$

where a's range over elements of V. There is a left K-module on $V[[h]]$, and such modules are called *topologically free modules*. One can observe that any topologically free module is separated and complete. Also, if $\{e_i\}_{i\in I}$ is a basis of V, then the K-submodule generated by $\{e_i\}_{i\in I}$ is dense in V for the h-adic topology. Besides, let N be a complete separated K-module, there is a natural isomorphism in the category of K-modules and K-linear maps:

$$\mathrm{Hom}_K(V[[h]], N) \cong \mathrm{Hom}_K(V, N)$$

Further, a K-module M is topologically free iff it is complete, separated and torsion-free, that is, there is such that $mh \neq 0$ for any non-zero $m \in M$.

Let M and N be left modules over the algebra $K = \mathbb{C}[[h]]$. The K-module $M \otimes_K N$ is the quotient of the vector space $M \otimes N$ obtained by spanning all elements of the form $fm \otimes n - m \otimes fn$ for $f \in K$, $m \in M$ and $n \in N$.

Definition 2.1. The topological tensor product $M\widetilde{\otimes}N$ of M and N is the h-adic completion of $M \otimes_K N$:

$$M\widetilde{\otimes}N := \widetilde{M \otimes_K N} = \varprojlim_{n>0} (M \otimes_K N)/h^n(M \otimes_K N)$$

By definition, $M\widetilde{\otimes}N$ is complete.

Take $m \in M$ and $n \in N$, let $m\widetilde{\otimes}n$ denote the image of $m \otimes n$ under the following image:

$$M \otimes N \to M \otimes_K N \to M\widetilde{\otimes}N$$

The subspace of the topological tensor product spanned by all elements of this form is dense in $M \otimes N$.

Proposition 2.1.

1. The class left modules over the algebra $K = \mathbb{C}[[h]]$ with $\widetilde{\otimes}$ is endowed with the structure of a symmetric monoidal category
2. If M and N be topologically free modules, so is $M\widetilde{\otimes}N$, that is:

$$V[[h]]\widehat{\otimes}W[[h]] = (V \otimes W)[[h]].$$

3 On Dialogue Categories and their internal language

Definition 3.1. A dialogue category is a monoidal category $(\mathcal{C}, \otimes, e)$ equipped with an object $\perp$ along with a family of isomorphisms

$$\varphi_{A,B} : \mathcal{C}(A \otimes B, \perp) \cong \mathcal{C}(B, A \multimap \perp)$$

natural in x and y; along with a family of isomorphisms

$$\psi_{A,B} : \mathcal{C}(A \otimes B, \perp) \cong \mathcal{C}(A, \perp \circ\!\!- B)$$

natural in x and y as well.

3.1 Term Assignment for Tensorial Logic

Figure 1: Term Calculus for Tensorial Logic

$$\frac{}{\vdash \mathbf{1} : \mathbb{1}}\ \mathbb{1}\mathbf{I} \qquad \frac{}{x : A \vdash x : A}\ \mathbf{ax} \qquad \frac{\Gamma \vdash M : \mathbb{1} \qquad \Delta \vdash N : B}{\Gamma, \Delta \vdash \mathbf{let}\ \mathbf{1} = M\ \mathbf{in}\ N : B}\ \mathbb{1}\mathbf{E}$$

$$\frac{x : A, \Gamma \vdash M : \perp}{\Gamma \vdash \nabla_l x.M : A \multimap \perp}\ \multimap \perp_{\mathbf{I}} \qquad \frac{\Delta \vdash N : A \qquad \Gamma \vdash M : A \multimap \perp}{\Gamma, \Delta \vdash N \triangleleft M : \perp}\ \multimap \mathbf{E}$$

$$\frac{\Gamma, x : A \vdash M : \perp}{\Gamma \vdash \nabla_r x.M : \perp \circ\!\!- A}\ \perp \circ\!\!-_{\mathbf{I}} \qquad \frac{\Gamma \vdash M : A \circ\!\!- \perp \qquad \Delta \vdash N : A}{\Gamma, \Delta \vdash M \triangleright N : \perp}\ \multimap \mathbf{E}$$

$$\frac{\Gamma \vdash M : A \qquad \Delta \vdash N : B}{\Gamma, \Delta \vdash M \otimes N : A \otimes B}\ \otimes\mathbf{I} \qquad \frac{\Gamma \vdash M : A \otimes B \qquad \Delta, x : A, y : B \vdash N : C}{\Delta, \Gamma \vdash \mathbf{let}\ x \otimes y\ =\ M\ \mathbf{in}\ N : C}\ \otimes\mathbf{E}$$

Definition 3.2. Reduction rules for tensorial logic:

$(\multimap \perp\beta)$ $N \triangleleft (\nabla_l x.M) \to M[x := N]$,

$(\perp \circ\!\!- \beta)$ $(\nabla_r x.M) \triangleright N \to M[x := N]$,

$(\mathbb{1}\beta)$ $\mathbf{let}\ \mathbf{1} = \mathbf{1}\ \mathbf{in}\ N \to N$,

$(\otimes\beta)$ $\mathbf{let}\ x \otimes y = M \otimes N\ \mathbf{in}\ P \to P[x := M, y := N]$,

$(\multimap \perp\eta)$ $(\nabla_l x.x \triangleleft M) \to M$ if x is not free in M,

$(\perp \circ\!\!- \eta)$ $(\nabla_r x.M \triangleright x) \to M$ if x is not free in M,

$(\otimes\eta)$ $\mathbf{let}\ x \otimes y = M\ \mathbf{in}\ N[z := x \otimes y] \to N[z := M]$,

$(\otimes_{\mathbf{nat}})$ $P[w := \mathbf{let}\ x \otimes y = M\ \mathbf{in}\ N] \to \mathbf{let}\ x \otimes y = M\ \mathbf{in}\ P[w := N]$,

$(\mathbb{1}_{\mathbf{nat}})$ $P[w := \mathbf{let}\ \mathbf{1} = M\ \mathbf{in}\ N] \to \mathbf{let}\ \mathbf{1} = M\ \mathbf{in}\ P[w := N]$.

Questions:

- Do we need resource modalities? If so, then have a look at [MT10], [MTT18] and [MT07]. Exponential negation? Note sure what's more essential for the profinite setting.
- Do we need sums? Sums are essential for the resource-sensitive CPS as they model non-deterministic choice.
- When we model the profinite lambda terms, we use the category of finite sets as the starting point: intuitively, **FinSet** is the cartesian closed category of finite types. What's the analogue of **FinSet** for tensorial terms? $\mathbb{F}$-weighted automaton over Galois fields? Some references: [BR11], [MN23], [AMU18], [UACM16], [BBB$^+$12], [VZ14].

3.2 Parametricity for Tensorial Logic

The following subsection develops parametricity for the tensorial calculus in the fashion of [AMP25] and [ZZZ10].

References

[AMP25] CB Aberlé, Chris Martens, and Frank Pfenning. Substructural parametricity. *arXiv preprint arXiv:2503.03153*, 2025.

[AMU18] Jiří Adámek, Stefan Milius, and Henning Urbat. A categorical approach to syntactic monoids. *Logical Methods in Computer Science*, 14, 2018.

[BBB$^+$12] Filippo Bonchi, Marcello Bonsangue, Michele Boreale, Jan Rutten, and Alexandra Silva. A coalgebraic perspective on linear weighted automata. *Information and Computation*, 211:77–105, 2012.

[BR11] Jean Berstel and Christophe Reutenauer. *Noncommutative rational series with applications*. Number 137. Cambridge University Press, 2011.

[Kas12] Christian Kassel. *Quantum groups*. Springer Science & Business Media, 2012.

[MN23] Vincent Moreau and Lê Thành Dũng Nguyên. Syntactically and semantically regular languages of lambda-terms coincide through logical relations. *arXiv preprint arXiv:2308.00198*, 2023.

[MT07] Paul-André Melliès and Nicolas Tabareau. Resource modalities in game semantics. In *22nd Annual IEEE Symposium on Logic in Computer Science (LICS 2007)*, pages 389–398. IEEE, 2007.

[MT10] Paul-André Melliès and Nicolas Tabareau. Resource modalities in tensor logic. *Annals of Pure and Applied Logic*, 161(5):632–653, 2010.

[MTT18] Paul-André Melliès, Nicolas Tabareau, and Christine Tasson. An explicit formula for the free exponential modality of linear logic. *Mathematical Structures in Computer Science*, 28(7):1253–1286, 2018.

[Pin10] Jean-Éric Pin. Mathematical foundations of automata theory. *Lecture notes LIAFA, Université Paris*, 7:73, 2010.

[SB81] Hanamantagouda P Sankappanavar and Stanley Burris. A course in universal algebra. *Graduate Texts Math*, 78:56, 1981.

[UACM16] Henning Urbat, Jiří Adámek, Liang-Ting Chen, and Stefan Milius. Eilenberg theorems for free. *arXiv preprint arXiv:1602.05831*, 2016.

[vGMM23] Sam van Gool, Paul-André Melliès, and Vincent Moreau. Profinite lambda-terms and parametricity. *Electronic Notes in Theoretical Informatics and Computer Science*, 3, 2023.

[VZ14] Benoît Valiron and Steve Zdancewic. Modeling simply-typed lambda calculi in the category of finite vector spaces. *Scientific Annals of Computer Science*, 24(2):325, 2014.

[ZZZ10] Jianzhou Zhao, Qi Zhang, and Steve Zdancewic. Relational parametricity for a polymorphic linear lambda calculus. In *Asian Symposium on Programming Languages and Systems*, pages 344–359. Springer, 2010.