

# Notes on filtrations for logics that contain **K5**

Daniel Rogozin

## 1 Preliminaries

**Definition 1.** A normal modal logic is a set of formulas that contains all Boolean tautologies, formulas  $\Diamond p \vee \Diamond q \leftrightarrow \Diamond(p \vee q)$  and  $\Diamond \perp \leftrightarrow \perp$ , and is closed under Modus Ponens, Substitution, and Monotonicity: from  $\varphi \rightarrow \psi$  infer  $\Diamond \varphi \rightarrow \Diamond \psi$ .

**Definition 2.** An Kripke model is a triple  $\mathcal{M} = \langle W, R, \vartheta \rangle$ , where  $R \subseteq W \times W$ ,  $\vartheta : PV \rightarrow 2^W$ , and the connectives have the following semantics:

1.  $\mathcal{M}, w \models p \Leftrightarrow w \in \vartheta(p)$
2.  $\mathcal{M}, w \models \neg \varphi \Leftrightarrow \mathcal{M}, w \not\models \varphi$
3.  $\mathcal{M}, w \models \varphi \vee \psi \Leftrightarrow \mathcal{M}, w \models \varphi$  or  $\mathcal{M}, w \models \psi$
4.  $\mathcal{M}, w \models \Diamond \varphi \Leftrightarrow \exists v \in R(w) \mathcal{M}, v \models \varphi$

A modal logic  $\mathcal{L}$  is called locally finite, if every fragment of  $\mathcal{L}$  with a finite number of variables is finite. Equivalently, its weak canonical model generated by  $k < \omega$  variables is finite. Moreover, if  $\mathcal{L}$  is locally finite, then  $\mathcal{L} = \text{Log}(\mathcal{F}_{\mathcal{L}} \upharpoonright k \mid k < \omega)$ , where  $\mathcal{F}_{\mathcal{L}} \upharpoonright k$  is a weak canonical frame generated by  $k$  variables.

**Definition 3.** The modal depth of a formula is defined recursively by the following function:

$$\begin{aligned} md(p_i) &= 0 \\ md(\neg \varphi) &= md(\varphi) \\ md(\varphi \vee \psi) &= \max(md(\varphi), md(\psi)) \\ md(\Diamond \varphi) &= 1 + md(\varphi) \end{aligned}$$

Let  $\mathcal{L}$  be a modal logic, then the modal depth of  $\mathcal{L}$ , denoted as  $md(\mathcal{L})$ , is the minimal  $n$  such that any  $\varphi \in \mathcal{L}$  is equivalent to some  $\psi$  such that  $md(\psi) = n$ . If such an  $n$  does not exist, then  $md(\mathcal{L}) = \infty$ .

### 1.1 Filtrations

Let  $\mathcal{M} = \langle W, R_1, \dots, R_n, \vartheta \rangle$  be a Kripke model and  $\Gamma$  a set of formulas closed under subformulas. An equivalence relation  $\sim$  is set to have a finite index if the quotient set  $W / \sim$  is finite. The equivalence relation  $\sim_{\Gamma}$  induced by  $\Gamma$  is defined as

$$w \sim_{\Gamma} v \Leftrightarrow \forall \varphi \in \Gamma (\mathcal{M}, w \models \varphi \Leftrightarrow \mathcal{M}, v \models \varphi).$$

If  $\Gamma$  is finite, then  $\sim_{\Gamma}$  has a finite index. An equivalence relation  $\sim$  respects  $\sim_{\Gamma}$ , if  $w \sim v$  implies  $w \sim_{\Gamma} v$ .

The following definition of a filtration is due to, e.g., [9].

**Definition 4.** Let  $\mathcal{M} = \langle W, R_1, \dots, R_n, \vartheta \rangle$  be a Kripke model and  $\Gamma$  be a Sub-closed set formulas. A  $\Gamma$ -filtration of  $\mathcal{M}$  is a model  $\widehat{\mathcal{M}} = \langle \widehat{W}, \widehat{R}_1, \dots, \widehat{R}_n, \widehat{\vartheta} \rangle$  such that:

1.  $\widehat{W} = W / \sim$ , where  $\sim$  is an equivalence relation that respects  $\Gamma$
2.  $\widehat{\vartheta}(p) = \{[x]_{\sim} \mid x \in W \ \& \ x \in \vartheta(p)\}$
3. For each  $i \in I$  one has  $\widehat{R}_i^{\min} \subseteq \widehat{R}_i \subseteq \widehat{R}_i^{\max}$ .  $\widehat{R}_{i,\sim}^{\min}$  is the  $i$ -th minimal filtered relation on  $\widehat{W}$  defined as

$$\hat{x} \widehat{R}_{i,\sim}^{\min} \hat{y} \Leftrightarrow \exists x' \sim x \exists y' \sim y \ x R_i y$$

$\widehat{R}_{\Gamma,i}^{\max}$  is the  $i$ -th maximal filtered relation on  $\widehat{W}$  induced by  $\Gamma$  defined as

$$\hat{x} \widehat{R}_{\Gamma,i}^{\max} \hat{y} \Leftrightarrow \forall \Box_i \varphi \in \Gamma \ (\mathcal{M}, x \models \Box_i \varphi \Rightarrow \mathcal{M}, y \models \varphi)$$

Alternatively, one may reformulate the condition of the maximal filtered relations using  $\Diamond$ 's as follows. We will use this formulation occasionally:

$$\hat{x} \widehat{R}_{\Gamma,i}^{\max} \hat{y} \Leftrightarrow \forall \Diamond_i \varphi \in \Gamma \ (\mathcal{M}, y \models \varphi \Rightarrow \mathcal{M}, x \models \Diamond_i \varphi)$$

If  $\Phi$  is an extension of  $\Gamma$  and  $\sim = \sim_{\Phi}$ , then  $\widehat{\mathcal{M}}$  is a definable  $\Gamma$ -filtration of  $\mathcal{M}$  through  $\Phi$ . If  $\sim = \sim_{\Gamma}$ , then such a filtration by means of the definition above is called *strict*.

A class of models  $\mathbb{M}$  admits strict filtrations for models (ASF), if for every Sub-closed set  $\Gamma$  and for every  $\mathcal{M} \in \mathbb{M}$  there exists a model  $\widehat{\mathcal{M}}$  such that  $\widehat{\mathcal{M}} \in \mathbb{M}$  and  $\widehat{\mathcal{M}}$  is a filtration of  $\mathcal{M}$  through  $\Gamma$ .

A class of frames  $\mathbb{F}$  admits strict filtrations for frames, if for every Sub-closed set  $\Gamma$  and for every frame  $\mathcal{F} \in \mathbb{F}$  and every model  $\mathcal{M}$  over  $\mathcal{F}$  there exists a  $\Gamma$  filtration of  $\mathcal{M}$ , and the underlying frame of this filtration belongs to  $\mathbb{F}$ .

If  $\mathcal{L}$  is canonical, then the ASF property for frames and ASF property for models are equivalent, see [7, Theorem 2.10].

The key lemma about filtrations is the following, see [1, Theorem 2.39]:

**Lemma 1.** Let  $\Gamma$  be a finite set of formulas closed under subformulas and  $\widehat{\mathcal{M}}$  a filtration of  $\mathcal{M}$  through  $\Gamma$ , then for each  $x \in W$  and for each  $\varphi \in \Gamma$  one has

$$\mathcal{M}, x \models \varphi \Leftrightarrow \widehat{\mathcal{M}}, \hat{x} \models \varphi$$

**Definition 5.**

1. Let  $\mathbb{F}$  be a class of Kripke frames and  $\Gamma$  a finite set of formulas closed under subformulas. If for every model  $\mathcal{M}$  over  $\mathcal{F} \in \mathbb{F}$  there exists a model that is a  $\Gamma$ -definable filtration of  $\mathcal{M}$ , then  $\mathbb{F}$  admits definable filtration.
2. A class of models  $\mathbb{M}$  admits definable filtration if for every  $\mathcal{M} \in \mathbb{M}$  there exists a model belonging to the same class that is a definable  $\Gamma$ -filtration of  $\mathcal{M}$ .

**Lemma 2.**

1. Let  $\mathcal{L}$  be a complete normal modal logic. If  $\text{Frames}(\mathcal{L})$  admits filtration, then  $\mathcal{L}$  has the finite model property.
2. If the class of models  $\text{Mod}(\mathcal{L})$  admits filtration, then  $\mathcal{L}$  has the finite model property and it is Kripke complete as well.

## 1.2 Horn closure

**Definition 6.** A first-order formula is called *Horn* if it has the following form (see [3]):

$$\forall x_1, \dots, x_n (x_{i_1} R x_{j_1} \wedge \dots \wedge x_{i_s} R x_{j_s} \rightarrow A), \text{ where } A \text{ is either } x_k R x_l \text{ or } \perp.$$

**Definition 7.** Let  $H$  be a Horn property and  $\langle W, R \rangle$  a Kripke frame. A Horn closure of a binary relation  $R$  is the minimal relation  $R^H$  containing  $R$  and satisfying  $H$ .

**Lemma 3.**  $R^H = \bigcup_{n < \omega} R_n$  where

1.  $R_0 = R$ .
2.  $R_{n+1} = R_n \cup \{(a, b) \in W \mid \exists \vec{c} \in W \text{ } P(a, b, \vec{c})\}$ , where  $P$  is a premise of  $H$ .

## 1.3 Bisimulations

A model  $\mathcal{M}$  is called *weak* model, if its valuation is restricted to some finite subset of variables, namely  $PV \upharpoonright k$ , if the cardinality of this subset equals  $k$ .

**Definition 8.** Let  $\mathcal{M}_1 = \langle W_1, R_1, \vartheta_1 \rangle$  and  $\mathcal{M}_2 = \langle W_2, R_2, \vartheta_2 \rangle$  be weak Kripke models (with the same variables),  $x \in W_1$ , and  $y \in W_2$ , then  $x$  and  $y$  are 0-equivalent, if the following holds:

$$x \equiv_0 y \text{ iff for each } p \in PV \upharpoonright k \text{ (or for each } p \in PV \text{) one has } \mathcal{M}_1, x \models p \Leftrightarrow \mathcal{M}_2, y \models p$$

**Definition 9.** Let  $\mathcal{M}_1 = \langle W_1, R_1, \vartheta_1 \rangle$   $\mathcal{M}_2 = \langle W_2, R_2, \vartheta_2 \rangle$  be Kripke models, then a bisimulation between  $\mathcal{M}_1$  and  $\mathcal{M}_2$  is a binary relation  $E \subseteq W_1 \times W_2$  between two models such that:

- (0)  $x_1 E x_2$  implies  $x_1 \equiv_0 x_2$
- (**zig**)  $x_1 E y_1$  and  $x_1 R_1 z$  implies  $y_1 R_2 y_2$  and  $z E y_2$  for some  $y_2 \in W_2$
- (**zag**)  $x_1 E y_1$  and  $y_1 R_2 y_2$  implies  $x_1 R_1 x_2$  and  $x_2 E y_2$  for some  $x_2 \in W_1$

Models with designated points  $\langle \mathcal{M}_1, x \rangle$  and  $\langle \mathcal{M}_2, y \rangle$  are bisimilar, if there exists a bisimulation  $E$  between them with  $x E y$ . We denote that as  $\langle \mathcal{M}_1, x \rangle \Leftrightarrow \langle \mathcal{M}_2, y \rangle$ . Moreover, we have

$$\langle \mathcal{M}_1, x \rangle \Leftrightarrow \langle \mathcal{M}_2, y \rangle \text{ implies } Th(\mathcal{M}_1, x) = Th(\mathcal{M}_2, y).$$

**Definition 10.** Let  $\langle \mathcal{M}_1, x \rangle$  and  $\langle \mathcal{M}_2, y \rangle$  be models with designated points and  $r < \omega$ . The  $r$ -round bisimulation game  $\mathcal{G}_r(\mathcal{M}_1, \mathcal{M}_2, x, y)$  is played by two players,  $\forall$  (Abelard, man) and  $\exists$  (Heloise, woman). The rules of the game are the following:

1. (**Round 0**)  $\exists$  wins, if  $x \equiv_0 y$ . Otherwise,  $\forall$  wins.
2. (**Round  $n+1$** ) Let  $n+1 < r$ . Suppose rounds  $0, \dots, n$  have been played. The  $n$ -th position is a pair  $(x_n, y_n)$ .  $\forall$  chooses  $x_{n+1}$  (or  $y_n \in W_2$ ) with  $x_n R_1 x_{n+1}$  (or  $y_n R_1 y_{n+1}$ ).  $\exists$  chooses  $y_{n+1}$  ( $x_{n+1}$ ) with  $y_n R_2 y_{n+1}$  (or  $y_n R_2 y_{n+1}$ ) and  $x_{n+1} \equiv_0 y_{n+1}$ .  $\forall$  ( $\exists$ ) wins, if  $\exists$  ( $\forall$ ) cannot move.

$\exists$  wins the game after  $r$  rounds.  $\forall$  wins the game, if there exists  $k < r$  such that  $\forall$  wins in Round  $k$ .

There are two  $n$ -equivalences for formulas and games (the first one is due to Fine [4]).

**Definition 11.**

- $(\mathcal{M}_1, x) \equiv_n (\mathcal{M}_2, y)$  iff for every  $\varphi$  such that  $md(\varphi) \leq n$  we have  $\mathcal{M}_1, x \models \varphi$  iff  $\mathcal{M}_2, y \models \varphi$ .
- $(\mathcal{M}_1, x) \equiv_n (\mathcal{M}_2, y)$  iff  $\exists$  has a winning strategy in  $\mathcal{G}_n(\mathcal{M}_1, \mathcal{M}_2, x, y)$ .

**Lemma 4** (See [6]). *The definitions of  $n$ -equivalence are equivalent for weak Kripke models.*

**Lemma 5.** (*Stabilisation lemma*)

*Let  $n$  be a natural number and  $\mathcal{L}$  a modal logic. If  $\equiv_n = \equiv_{n+1}$  in a weak canonical model  $\mathcal{M}_{\mathcal{L}} \upharpoonright k$ , then  $md(\mathcal{L} \upharpoonright k) \leq n$ .*

**Theorem 1** (See [5]). *Let  $\mathcal{L}$  be a modal logic such that  $md(\mathcal{L}) = n$  for some  $n < \omega$ , then  $\mathcal{L}$  is locally finite.*

## 2 Filtrations for K5

The **K5**-closure (the Euclidean Horn closure of a binary relation) has the following equivalent definitions:


**Lemma 6.** *Let  $\mathcal{F} = \langle W, R \rangle$  be a Kripke frame. The following are equivalent:*

1.  $R^{\mathbf{K5}}$  is the smallest Euclidean relation containing  $R$ , that is, the Horn closure of  $R$ .
2.  $R^{\mathbf{K5}} = \bigcup_{i < \omega} R_i$ , where
  - $R_0 = R$
  - $R_{n+1} = R_n \cup (R_n^{-1} \circ R_n)$
3.  $xR^{\mathbf{K5}}y$  iff there exists  $n < \omega$  such that either  $xRy$  or  $\exists z_1, \dots, z_n$  with  $z_1Rx$  and  $z_{n-1}Ry$  and for each  $1 < i \leq n$  one has either  $z_{i-1}Rz_i$  or  $z_iRz_{i-1}$ .
4.  $R^{\mathbf{K5}} = R \cup \bigcup_{i < \omega} (R^{-1} \circ (R \cup R^{-1})^n \circ R)$ .

**Theorem 2.** *K5 does not admit strict filtrations.*

*Proof.* Let us consider a **K5** model whose Euclidean closure of the minimal filtration does not give us a filtration.

Let us consider a frame called  $\mathcal{F}_{bad}$ . We define this frame with the following graph:



Let us define a valuation  $\vartheta$  such that  $\vartheta(p) = \{w_5\}$  and  $\vartheta(q) = \{w_1\}$ . Let us consider a minimal filtration of  $\mathcal{M}_{bad}$  through the Sub-closure of  $\Gamma = \{\neg p, \neg \Diamond p\}$ .

Clearly that  $w_2 \sim_{\Gamma} w_3$ , since  $\neg p$  and  $\neg \Diamond p$  are true both at  $w_2$  and  $w_3$ .

Moreover,  $R_{min} \cup (R_{min}^{-1} \circ R_{min})$  is not a subset of  $R_{max}$  since  $(\hat{w}_1, \hat{w}_5) \in (R_{min}^{-1} \circ R_{min})$ , but  $\Diamond p$  is not true at  $w_5$ .

Let us also note that strict filtrations of this model is not Euclidean. Suppose by contrary that  $\hat{R}^{\mathcal{E}}$  is a strict filtration of that model. So  $R_{min}^{\mathcal{E}} \subseteq \hat{R}^{\mathcal{E}}$ , since  $R_{min}^{\mathcal{E}}$  is the minimal Euclidean relation containing  $R_{min}$ . On the other hand,  $R_{min}^{\mathcal{E}} \not\subseteq R_{max}$ , so is not  $\hat{R}^{\mathcal{E}}$ .  $\square$

**Theorem 3.** **K5** admits definable filtrations.

**Theorem 4.** **K45** admits strict filtrations.

*Proof.* Let  $\mathcal{M} = \langle W, R, \vartheta \rangle$  be a transitive Euclidean model and  $\overline{\mathcal{M}} = \langle \overline{W}, \overline{R}, \overline{\vartheta} \rangle$  its minimal filtration through  $\Gamma$ , where  $\Gamma$  is finite and Sub-closed. Let us put  $\hat{R} = \overline{R}^+ \cup \overline{R}^{\mathbf{K5}}$ . Let us show that  $\overline{R}^+ \cup \overline{R}^{\mathbf{K45}} \subseteq \overline{R}^{max}$ .

That is, if  $\mathcal{M}, y \models \varphi$  for  $\Diamond \varphi \in \Gamma$  and  $\hat{x}\hat{R}\hat{y}$ , then  $\mathcal{M}, x \models \Diamond \varphi$ .

Let  $\hat{x}\hat{R}\hat{y}$ .

1. Suppose  $(\hat{x}, \hat{y}) \in \overline{R}$ , then  $\mathcal{M}, x \models \Diamond \varphi$  holds trivially by the definition of the minimal filtration.
2. Let us consider the case when  $(\hat{x}, \hat{y}) \in \overline{R}^{\mathbf{K5}}$ . The second alternative is the same as for the **K4**-case, see [2, p. 141].

Suppose the statement holds  $\overline{R}_n$  and  $(\hat{x}, \hat{y}) \in \overline{R}_{n+1} = \overline{R}_n \cup (\overline{R}_n^{-1} \circ \overline{R}_n)$ . We consider the case of  $(\hat{x}, \hat{y}) \in (\overline{R}_n^{-1} \circ \overline{R}_n)$ .

Then there exists  $\hat{z}$  such that  $(\hat{z}, \hat{x}), (\hat{z}, \hat{y}) \in \overline{R}_n$ .

By IH,  $\mathcal{M}, z \models \Diamond \varphi$ .

$(\hat{z}, \hat{y}) \in \overline{R}_n$  iff there are  $\hat{u}_1, \dots, \hat{u}_n$  such that

$$\hat{z} \xleftarrow{\hat{R}} \hat{u}_1 \xrightarrow{\hat{R}'} \hat{u}_2 \xrightarrow{\hat{R}'} \dots \xrightarrow{\hat{R}'} \hat{u}_{n-1} \xrightarrow{\hat{R}'} \hat{u}_n \xrightarrow{\hat{R}} \hat{y}$$

where  $\hat{R}'$  is either  $\hat{R}$  or  $\hat{R}^{-1}$ .

As it is known,  $\Diamond \Diamond \varphi \rightarrow \Box \Diamond \varphi \in \mathbf{K45}$ .

$\hat{u}_1 \hat{z}$ , that is,  $u'_1 R z'$  for some  $u'_1 \in \hat{u}_1$  and  $z' \in \hat{z}$ . That is,  $\mathcal{M}, u'_1 \models \Diamond \Diamond \varphi$ , so  $\mathcal{M}, u'_1 \models \Diamond \varphi$  and  $\overline{\mathcal{M}}, \hat{u}_1 \models \Diamond \varphi$ .

We have  $\hat{u}_1 \hat{R}' \hat{u}_2$ . Suppose  $\mathcal{M}, u''_1 \models \Diamond \varphi$  and  $u''_1 R u'_2$ . We also have  $\mathcal{M}, u''_1 \models \Box \Diamond \varphi$ , thus,  $\mathcal{M}, u'_2 \models \Diamond \varphi$ .

Suppose  $\hat{u}_2 \hat{R} \hat{u}_1$  and  $u'_2 R u''_1$ , then  $\mathcal{M}, u'_2 \models \Diamond \varphi$ .

Similarly, we have  $\mathcal{M}, u_i \models \Diamond \varphi$  iff  $\mathcal{M}, u_{i+1} \models \Diamond \varphi$ , whenever  $\hat{u}_i \hat{R}' \hat{u}_{i+1}$ .

Finally, we have  $\hat{u}_n \hat{R} \hat{x}$ . Thus,  $u'_n R x'$  for some  $u'_n \in \hat{u}_n$  and  $x' \in \hat{x}$ .  $\mathcal{M}, u'_n \models \Diamond \varphi$ , so  $\mathcal{M}, u'_n \models \Box \Diamond \varphi$ . Then  $\mathcal{M}, x' \models \Diamond \varphi$ .  $\square$

### 3 i,j-Euclideaness

**Definition 12.** A binary relation  $R \subseteq W \times W$  is called *i, j-Euclidean* for  $i, j < \omega$ , if for each  $x, y, z$  such that  $xR^i y$  and  $xR^j z$  implies  $xRz$ .

**Proposition 1.** Let  $\mathcal{F} = \langle W, R \rangle$ , then  $\mathcal{F}$  is *i, j-Euclidean* iff  $\mathcal{F} \models \Diamond^i p \rightarrow \Box^j \Diamond p$ . As a corollary, the logic  $\mathbf{K5}^{i,j} = \mathbf{K} \oplus \Diamond^i p \rightarrow \Box^j \Diamond p$  is Kripke-complete.

Let  $R$  be a binary relation on  $W \neq \emptyset$ , then the *i, j-Euclidean* closure of  $R$  (where  $i, j < \omega$ ), denoted as  $R^{\mathbf{K5}^{i,j}}$ , is a binary relation defined recursively as follows:

1.  $R_0 = R$

2.  $R_{n+1} = R_n \cup (((R_n)^{-1})^i \circ R_n^j)$
3.  $R^{\mathbf{K5}^{i,j}} = \bigcup_{k < \omega} R_k$

Note that the following property holds not for every  $\mathbf{K5}^{i,j}$ , consider  $\mathbf{KB}$  as an example (clearly,  $\mathbf{KB} = \mathbf{K5}^{0,1}$ ), whose modal depth is  $\infty$ .

**Exercise 1.**  $md(\mathbf{K5}) = 2$ .

*Proof.* For the upper bound, let us show that  $\equiv_2 = \equiv_3$  in every weak canonical model of  $\mathbf{K5}$ , namely  $\mathcal{L}_{\mathbf{K5}} \upharpoonright k$  for  $k < \omega$ . For that, we show that if  $\exists$  can win  $\mathcal{G}_2(\mathcal{L}_{\mathbf{K5}} \upharpoonright k, \Gamma, \Delta)$  (where  $\Gamma$  and  $\Delta$  are maximal consistent  $\mathbf{K5} \upharpoonright k$ -theories), then she can the game  $\mathcal{G}_2(\mathcal{L}_{\mathbf{K5}} \upharpoonright k, \Gamma, \Delta)$  as well.

If  $\exists$  has a winning strategy in  $\mathcal{G}_2(\mathcal{L}_{\mathbf{K5}} \upharpoonright k, \Gamma_0, \Delta_0)$ , then, by Lemma 4,  $\langle \mathcal{L}_{\mathbf{K5}} \upharpoonright k, \Gamma_0 \rangle \equiv_2 \langle \mathcal{L}_{\mathbf{K5}} \upharpoonright k, \Delta_0 \rangle$ .  $\square$

**Exercise 2.** Let  $i = j$ , then  $md(\mathbf{K5}^{i,i}) = i + 1$ . So  $\mathbf{K5}^{i,i}$  is locally finite.

*Proof.*  $\square$

**Theorem 5.**  $md(\mathbf{K5}^{i,j}) = 1 + j + i$ , where  $i > 0$ .

**Theorem 6.**  $\mathbf{K5}^{i,j}$  admits definable filtrations.

*Proof.* Let  $\mathcal{M} = \langle W, R, \vartheta \rangle$  and  $\Gamma$  a finite Sub-closed of formulas. We extend  $\Gamma$  as

$$\Delta = \Gamma \cup \text{Sub}\{\Diamond^i \psi \mid \Box \psi \in \Gamma\} \cup \text{Sub}\{\Box^j \Diamond \psi \mid \Diamond \psi \in \Gamma\}$$

Let  $(\hat{x}, \hat{y}) \in R^{\mathbf{K5}^{i,j}}$ ,  $\mathcal{M}, x \models \Box \psi$  for  $\Box \psi$  in  $\Delta$ . If  $n = 0$ , then the statement is obvious. Suppose  $n = 1$ . Then  $(\hat{x}, \hat{y}) \in R_{min} \cup (((R_{min})^{-1})^i \circ R_{min}^j)$ . Consider the second alternative. Then there exists  $\hat{z}$  such that  $(\hat{x}, \hat{z}) \in ((R_{min})^{-1})^i$  and  $(\hat{z}, \hat{y}) \in R_{min}^j$ .

That is, there are  $\hat{x}_1, \hat{x}_2, \dots, \hat{x}_i$  and  $\hat{y}_1, \hat{y}_2, \dots, \hat{y}_j$  such that

$$\hat{z} \xrightarrow{R_{min}} \hat{x}_1 \xrightarrow{R_{min}} \hat{x}_2 \xrightarrow{R_{min}} \dots \xrightarrow{R_{min}} \hat{x}_i \xrightarrow{R_{min}} \hat{x}$$

$$\hat{z} \xrightarrow{R_{min}} \hat{y}_1 \xrightarrow{R_{min}} \hat{y}_2 \xrightarrow{R_{min}} \dots \xrightarrow{R_{min}} \hat{y}_j \xrightarrow{R_{min}} \hat{y}$$

???

$\square$


## 4 The case of 2-transitivity

Let us define the logic  $\mathcal{L}$  as  $\mathbf{K} \oplus \Diamond \Diamond \Diamond p \rightarrow \Diamond p$ . Let  $R$  be a binary relation, the  $\mathcal{L}$ -closure of  $R$  is defined (denoted as  $R^{\hat{\Diamond}}$ ) as the following union:

$$R^{\hat{\Diamond}} = R \cup R^3 \cup R^5 \cup \dots \cup R^{2k+1} \cup \dots$$

**Theorem 7.**  $\mathcal{L}$  does not admit strict filtrations.

*Proof.* Consider the following frame  $\mathcal{F} = \langle W, R \rangle$ :



Clearly that  $\mathcal{F}$  is an  $\mathcal{L}$ -frame. We define the valuation  $\vartheta$  as follows:

$$\begin{aligned}\vartheta(p) &= \{x\} \\ \vartheta(q) &= \{y, y_1\} \\ \vartheta(r) &= \{z\}\end{aligned}$$

Let us put  $\Gamma = \text{Sub}\{p, q, \Diamond r\}$ . We factorise  $W$  through  $\sim_\Gamma$  and consider a model  $\widehat{\mathcal{M}} = \langle W / \sim_\Gamma, \widehat{R}, \widehat{\vartheta} \rangle$ , where  $\widehat{R} = (\widehat{R}_{\min})^\Delta$ . We have  $(\hat{x}, \hat{z}) \in \widehat{R} \circ \widehat{R} \circ \widehat{R}$ , but  $\Diamond r$  is not true at  $x$ .  $\square$

## 5 Fusion stuff

**Definition 13.** Let  $\mathcal{L}_1$  and  $\mathcal{L}_2$  be modal logics, then the fusion  $\mathcal{L}_1 * \mathcal{L}_2$  is the minimal bimodal logic that contains  $\mathcal{L}_1$  and  $\mathcal{L}_2$  [8].

**Lemma 7.** Let  $\Gamma$  be a finite and Sub-closed set of formulas.

1.  $\mathcal{M} = \langle W, R, \vartheta \rangle$  be a **K5**-model. Consider  $\Gamma' = \Gamma \cup \{\Diamond \Box \psi \mid \Box \psi \in \Gamma\}$ . Let  $\Delta$  be any finite and Sub-closed extension of  $\Gamma'$  then a model  $\widehat{\mathcal{M}} = \langle W / \sim_\Delta, (R_{\Gamma'}^{\min})^{\mathbf{K5}}, \widehat{\vartheta} \rangle$  is a filtration of  $\mathcal{M}$  through  $\Gamma'$ .
2. Let  $\mathcal{L} = \mathbf{K} \oplus \Diamond \Diamond p \rightarrow \Diamond p$ , then we have the similar statement for  $\mathcal{L}$ , where  $\Delta$  is a Sub-closed extension of  $\Gamma'$ .

*Proof.*

Recall that  $(R_\Delta^{\min})^{\mathbf{K5}}$  is defined inductively as:

- (a)  $R_{\Gamma'}^0 = R_{\Gamma'}^{\min}$
- (b)  $R_{\Gamma'}^{n+1} = R_{\Gamma'}^n \cup (R_{\Gamma'}^{n-1} \circ R_{\Gamma'}^n)$
- (c)  $(R_{\Gamma'}^{\min})^E = \bigcup_{k < \omega} R_{\Gamma'}^k$

If  $R_{\Gamma'}^{\min}$  is already a subrelation of  $R^{\max}$ , so the base case is self-evident.

Suppose the statement holds for  $R_{\Gamma'}^n$ ,  $(\hat{x}, \hat{y}) \in R_{\Gamma'}^{n+1}$  such that  $\mathcal{M}, y \models \psi$  for  $\Diamond \psi \in \Gamma'$ .

According to the third item of Lemma 6, this is the same as there exist  $\widehat{z}_0, \widehat{z}_1, \dots, \widehat{z}_{n-1}, \widehat{z}_n$  such that  $\widehat{z}_1 R_{\Gamma'}^{\min} \hat{x}$ ,  $\widehat{z}_n R_{\Gamma'}^{\min} \hat{y}$ , and for each  $i \in n+1$  we have either  $\widehat{z}_i R_{\Gamma'}^{\min} \widehat{z}_{i+1}$  or  $\widehat{z}_{i+1} R_\Delta^{\min} \widehat{z}_i$ .

We visualise this with the following graph:

$$\hat{x} \xleftarrow{R_{\Gamma'}^{\min}} \widehat{z}_0 \xleftrightarrow{R'} \widehat{z}_1 \xleftrightarrow{R'} \dots \xleftrightarrow{R'} \widehat{z}_{n-1} \xrightarrow{R'} \widehat{z}_n \xrightarrow{R_{\Gamma'}^{\min}} \hat{y}$$

where  $R'$  is either  $R_{\Gamma'}^{\min}$  or its converse. We have  $\mathcal{M}, z \models \Box \psi$ ,  $\mathcal{M}, y \models \psi$ , so  $\mathcal{M}, z_n \models \Diamond \psi$ . Since  $\mathcal{M}$  is a **K5**-model, we have  $\mathcal{M}, z_n \models \Box \Diamond \psi$ .

After that we apply the following property of **K5**-models:

Let  $\mathcal{M} \models \mathbf{K5}$  and  $\varphi$  a formula, then for each  $a, b \in \mathcal{M}$  such that  $aRb$  we have  $\mathcal{M}, a \models \Box \Diamond \varphi$  iff  $\mathcal{M}, b \models \Box \Diamond \varphi$

So we have  $\mathcal{M}, z_0 \models \Box \Diamond \varphi$ . Note that we always stay within  $\Gamma'$ . Thus,  $\mathcal{M}, x \models \Diamond \varphi$ .

2. Let us prove the second item. Let  $\mathcal{M}$  be an  $\mathcal{L}$ -model. The  $\mathcal{L}$ -closure of the minimal filtered relation modulo  $\Gamma'$ , namely  $R_{\Gamma'}^{\min \mathcal{L}}$  has the following form:

$$R_{\Gamma'}^{min\mathcal{L}} = \bigcup_{k < \omega} R_{\Gamma'}^{min 2^{k+1}}$$

We reformulate this closure equivalently as follows:

- (a)  $R_0 = R_{\Gamma'}^{min}$
- (b)  $R_{n+1} = R_n \cup ((R_{\Delta}^{min})^2 \circ R_n)$
- (c)  $R_{\Gamma'}^{min\mathcal{L}} = \bigcup_{k < \omega}$

The base case is self-evident. Suppose the statement holds for  $R_n$  and  $(\hat{x}, \hat{y}) \in (R_{\Gamma'}^{min})^2 \circ R_n$ , that is, there exists  $\hat{z}$  such that  $(\hat{x}, \hat{z}) \in (R_{\Gamma'}^{min})^2$  and  $(\hat{z}, \hat{y}) \in R_n$ . By IH, we have  $\mathcal{M}, z \models \Diamond\varphi$ .

We have the following:

$$\hat{x} \xrightarrow{R_{\Delta}^{min}} \hat{y}_1 \xrightarrow{R_{\Gamma'}^{min}} \hat{z}$$

The sequence of implications is the following:

$$\widehat{\mathcal{M}}, \hat{z} \models \Diamond\varphi \Rightarrow \widehat{\mathcal{M}}, \hat{y}_1 \models \Diamond\Diamond\varphi \Rightarrow \mathcal{M}, x \models \Diamond\Diamond\Diamond\varphi \Rightarrow \mathcal{M}, x \models \Diamond\varphi$$

□

**Theorem 8.**

1. **K5** \* **K5** admits definable filtrations.
2. **K5** \*  $\dots$  \* **K5** admits definable filtrations.
3. If  $\mathcal{L}$  admits strict filtrations, then **K5** \*  $\mathcal{L}$  admits definable filtrations
4. If  $\mathcal{L}_1, \dots, \mathcal{L}_n$  admit strict filtrations, then **K5** \*  $\dots$  \* **K5** \*  $\mathcal{L}_1$  \*  $\dots$  \*  $\mathcal{L}_n$
5. Let  $\mathcal{L} = \mathbf{K} \oplus \Diamond\Diamond\Diamond p \rightarrow \Diamond p$  (here and below), then  $\mathcal{L} * \mathcal{L}$  admits definable filtrations.
6. Let  $\mathcal{L} = \mathbf{K} \oplus \Diamond\Diamond\Diamond p \rightarrow \Diamond p$  and  $\mathcal{L}_1$  a logic that admits strict filtrations, then  $\mathcal{L} * \mathcal{L}_1$

*Proof.*

1. Let  $\Gamma$  be a finite Sub-closed set of bimodal formulas,  $\mathcal{F} = \langle W, R_1, R_2 \rangle$  a **K5** \* **K5**-frame, and  $\vartheta$  a valuation on  $\mathcal{F}$ . Denote  $\langle \mathcal{F}, \vartheta \rangle$  as  $\mathcal{M}$ .

We introduce the set of fresh variables  $V = \{p_\psi \mid \psi \in \Gamma\}$  and define a new model  $\mathcal{M}' = \langle \mathcal{F}, \vartheta' \rangle$  as follows:

$$\text{For all } \psi \in \Gamma, \mathcal{M}, x \models \psi \Leftrightarrow \mathcal{M}', x \models \psi \Leftrightarrow \mathcal{M}', x \models p_\psi.$$

Consider these modifications of  $\Gamma$  and  $V$ :

$$\begin{aligned} \Gamma' &= \Gamma \cup \{\Diamond_1 \Box_1 \psi \mid \Box_1 \psi \in \Gamma\} \cup \{\Diamond_2 \Box_2 \psi \mid \Box_2 \psi \in \Gamma\} \\ \Delta &= V \cup \text{Sub}(\{\Diamond \Box p_\psi \mid \Box_i \psi \in \Gamma, in = 1, 2\}) \end{aligned}$$

Let us define an equivalence relation  $\sim_\Delta$  induced by  $\Delta$ .

Consider  $\mathcal{M}_i = \langle W, R_i, \vartheta' \rangle$ , a reduct of  $\mathcal{M}'$ , we have:



- (a)  $\mathcal{M}_i, x \models \Box p_\psi$  iff  $\mathcal{M}, x \models \Box_i \psi$
- (b)  $\mathcal{M}_i, x \models \Diamond \Box p_\psi$  iff  $\mathcal{M}, x \models \Diamond_i \Box_i \psi$

So  $\sim = \sim_{\Gamma'}$  by the construction. Let us put  $\widehat{W} = W / \sim_{\Gamma'}$ . Lemma 7 implies the following claim:

**Claim 1.** *Let  $\widehat{R}_i = (R_{\Delta}^{min})^E$  and  $\widehat{\vartheta}(p) = \{[x]_{\sim_i} \mid \mathcal{M}_i, x \models p\}$  for  $p \in \Delta_1$ , define  $\widehat{\mathcal{M}}_i = \langle \widehat{W}, \widehat{R}_i, \widehat{\vartheta} \rangle$ . Then  $\widehat{\mathcal{M}}_i \models \mathbf{K5}$  and  $\widehat{\mathcal{M}}_i$  is a filtration of  $\mathcal{M}_i$  through  $\Delta$ .*

Finally, we consider a model  $\widehat{\mathcal{M}} = \langle \widehat{W}, \widehat{R}_1, \widehat{R}_2, \widehat{\vartheta} \rangle$ , where  $\widehat{R}_{\Gamma'_i} = R_{i\Gamma'}^{min^E}$  and  $\widehat{\vartheta}(p)$  is defined as usual for  $p \in \Gamma$ .  $\widehat{\mathcal{M}}$  is a filtration of  $\mathcal{M}$  through  $\Gamma'$ .

Let  $\widehat{x}\widehat{R}_{\Gamma'_i}\widehat{y}$  and  $\mathcal{M}, x \models \Box_i \psi$  for  $\Box_i \psi \in \Gamma$ . Then  $\mathcal{M}_i, x \models \Box p_\psi$ , so  $\widehat{\mathcal{M}}_i, \widehat{x} \models \Box p_\psi$ . By the claim above,  $\widehat{\mathcal{M}}_i$  is a filtration of  $\mathcal{M}_i$  through  $\Delta$ , so  $\mathcal{M}_i, y \models p_\psi$ . Then  $\mathcal{M}, y \models \psi$ .

2. Likewise
3. The argument is the same as in the proof of the first item, except for **Claim 1** that has the following formulation: Let  $\widehat{R}_1 = (R_{\Delta}^{min})^E$  and  $\widehat{R}_2 = (R_{\Delta}^{min})^{\mathcal{L}_1}$ . Define a valuation as usual as  $\widehat{\vartheta}(p) = \{[x]_{\sim_i} \mid \mathcal{M}_i, x \models p\}$  for  $p \in \Delta_1$ , define  $\widehat{\mathcal{M}}_1 = \langle \widehat{W}, \widehat{R}_1, \widehat{\vartheta} \rangle$  and  $\widehat{\mathcal{M}}_2 = \langle \widehat{W}, \widehat{R}_2, \widehat{\vartheta} \rangle$ . Then  $\widehat{\mathcal{M}}_1 \models \mathbf{K5}$  and  $\widehat{\mathcal{M}}_1 \models \mathbf{L}$  and  $\widehat{\mathcal{M}}_i$  is a filtration of  $\mathcal{M}_i$  through  $\Delta$ .
4. Likewise
5. The argument is similar to the proof of first item, but filtrations are slightly different. Let  $\mathcal{M} = \langle W, R_1, R_2, \vartheta \rangle$  be a  $\mathcal{L} * \mathcal{L}$  model and  $\Gamma$  a Sub-closed set of formulas. As above, we define a set  $V$  and a model  $\mathcal{M}'$ . Define extensions of  $\Gamma$  and  $V$ :

$$\begin{aligned} \Gamma' &= \Gamma \cup \{\Diamond_1 \Diamond_1 \psi \mid \Diamond_1 \psi \in \Gamma\} \cup \{\Diamond_2 \Diamond_2 \psi \mid \Diamond_2 \psi \in \Gamma\} \\ \Delta &= V \cup \text{Sub}(\{\Diamond \Diamond p_\psi \mid \Diamond \psi \in \Gamma, i = 1, 2\}) \end{aligned}$$

As above  $\sim_{\Delta} = \Gamma'$  and  $\widehat{\mathcal{M}}' = \langle W / \sim_{\Delta}, \widehat{R}_i, \widehat{\vartheta} \rangle$  are filtrations of reducts of  $\mathcal{M}'$  through  $\Delta$ . Then  $\widehat{\mathcal{M}} = \langle W / \sim_{\Delta}, \widehat{R}_1, \widehat{R}_2, \widehat{\vartheta} \rangle$  is a required filtration of the original  $\mathcal{M}$ .

6. Extend  $\Gamma$  with  $\{\Diamond_i \Diamond_i \psi \mid \Diamond_i \psi \in \Gamma, i = 1, 2\}$  and  $V$  with  $\{\Diamond \Diamond p_\psi \mid \Diamond_i \psi, i = 1, 2\}$

□

**Theorem 9.** *Let  $\mathcal{L}_1$  and  $\mathcal{L}_2$  be modal logics that admit definable filtrations. If  $\text{Mod}(\mathcal{L}_1)$  and  $\text{Mod}(\mathcal{L}_2)$  admit definable filtrations, so does  $\text{Mod}(\mathcal{L}_1 * \mathcal{L}_2)$ .*

*Proof.* Let  $\mathcal{M} = \langle W, R_1, R_2, \vartheta \rangle$  be an  $\mathcal{L}_1 * \mathcal{L}_2$ -model. We define a notation  $\nabla = \{\neg \Diamond, \Diamond \neg, \Diamond\}$ .

Both logics admit definable filtrations, so for every finite Sub-closed set  $\Gamma$  and for every  $\mathfrak{M}$ , an  $\mathcal{L}_1$ -model (or an  $\mathcal{L}_2$  one) there exists there exists  $\Delta$ , a extension of  $\Gamma$  having the form:

$$\begin{aligned} \Delta_1 &= \Gamma \cup \text{Sub}\{\nabla_1 \nabla_2 \dots \nabla_n \Diamond \psi \mid \Diamond \psi \in \Gamma\} \text{ (for } \mathcal{L}_1) \\ \Delta_2 &= \Gamma \cup \text{Sub}\{\nabla_1 \nabla_2 \dots \nabla_k \Diamond \psi \mid \Diamond \psi \in \Gamma\} \text{ (for } \mathcal{L}_2) \end{aligned}$$

such that  $\widehat{\mathfrak{M}} = \langle W / \sim_{\Delta_i}, \widehat{R}, \widehat{\vartheta} \rangle$  is a filtration of  $\mathfrak{M}$  through the corresponding  $\Delta_i$ .

Let  $V$  be a set of fresh variables indexed over  $\Gamma$  as in the proof for a fusion of **K5** with something else. Let  $\mathcal{M}'$  be a model defined as previously. We extend  $V$  and  $\Gamma$  in the following way:

$$\begin{aligned}\Gamma' &= \Gamma \cup \text{Sub}\{\nabla_{11}\nabla_{21}\dots\nabla_{n1}\Diamond_1\psi \mid \Diamond_1\psi \in \Gamma\} \cup \text{Sub}\{\nabla_{12}\nabla_{22}\dots\nabla_{n2}\Diamond_2\psi \mid \Diamond_2\psi \in \Gamma\} \\ \Delta &= V \cup \text{Sub}\{\nabla_1\nabla_2\dots\nabla_n\Diamond p_\psi \mid \nabla_{n+1}\psi \in \Gamma'\} \cup \text{Sub}\{\nabla_1\nabla_2\dots\nabla_k\Diamond p_\psi \mid \Diamond_2\psi \in \Gamma\}.\end{aligned}$$

By the construction,  $\sim_{\Gamma'} = \sim_{\Delta}$ . So we have filtrations for the corresponding reducts of  $\mathcal{M}'$  through  $\Delta$  as well as for the original  $\mathcal{M}$ .  $\square$

## References

- [1] Patrick Blackburn, Maarten de Rijke, and Yde Venema. *Modal Logic*. Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, 2001.
- [2] Alexander Chagrov and Michael Zakharyashev. *Modal Logic*. Oxford Logic Guides, Volume 35. Oxford University Press, 1997.
- [3] Chen Chung Chang and H Jerome Keisler. *Model theory*. Elsevier, 1990.
- [4] Kit Fine. Logics containing k4. part i. *The Journal of Symbolic Logic*, 39(1):31–42, 1974.
- [5] Dov M Gabbay and Valentin B Shehtman. Products of modal logics, part 1. *Logic journal of IGPL*, 6(1):73–146, 1998.
- [6] Valentin Goranko and Martin Otto. Model theory of modal logic. In *Studies in Logic and Practical Reasoning*, volume 3, pages 249–329. Elsevier, 2007.
- [7] Stanislav Kikot, Ilya Shapirovsky, and Evgeny Zolin. Completeness of logics with the transitive closure modality and related logics. *arXiv preprint arXiv:2011.02205*, 2020.
- [8] Agi Kurucz. Combining modal logics. *Studies in Logic and Practical Reasoning*, 3:869–924, 2007.
- [9] Valentin B Shehtman. Filtration via bisimulation. *Advances in modal logic*, 5:289–308, 2004.