

# Representable cylindric algebras of dimension $\omega$ : the aspects of canonicity

Daniel Rogozin

## 1 Intro

## 2 The problem itself

Suppose  $\mathcal{C} \in \mathbf{RCA}_\omega$ , whether  $\mathcal{C}^+$  has a complete,  $\omega$ -dimensional representation? [3]

## 3 Boolean algebras with operators and cylindric algebras

**Definition 1.**

1. Let  $\mathcal{B} = \langle B, +, -, 0, 1 \rangle$  be a Boolean algebra. An operator is an  $n$ -ary function  $\Omega : B^n \rightarrow B$  satisfying the following conditions:

- Normality: for all  $b_0, \dots, b_{n-1} \in B$ , if  $b_i = 0$  for some  $i < n$ , then

$$\Omega(b_0, \dots, b_{i-1}, 0, b_{i+1}, \dots, b_{n-1}) = 0$$

- Additivity: for all  $b_0, \dots, b_{n-1}, b, b' \in B$  we have

$$\begin{aligned} \Omega(b_0, \dots, b_{i-1}, (b + b'), b_{i+1}, \dots, b_{n-1}) = \\ \Omega(b_0, \dots, b_{i-1}, b, b_{i+1}, \dots, b_{n-1}) + \Omega(b_0, \dots, b_{i-1}, b', b_{i+1}, \dots, b_{n-1}) \end{aligned}$$

2. Let  $I$  be an index set, a Boolean algebra with operators (BAO) is an algebra  $\langle B, +, -, 0, 1, \{\Omega_i\}_{i \in I} \rangle$  such that  $\langle B, +, -, 0, 1 \rangle$  is a Boolean algebra and for each  $i \in I$   $\Omega_i$  is an operator.

**Definition 2.** Let  $\mathcal{B} = \langle B, +, -, 0, 1, \{\Omega_i\}_{i \in I} \rangle$  be a BAO, then

1. An operator  $\Omega$  is completely additive, if for each  $b_0, \dots, b_{n-1} \in B$  and  $X \subseteq B$ , one has

$$\Omega(b_0, \dots, b_{i-1}, \sum X, b_{i+1}, \dots, b_{n-1}) = \sum_{x \in X} \Omega(b_0, \dots, b_{i-1}, x, b_{i+1}, \dots, b_{n-1})$$

2.  $\mathcal{B}$  is completely additive, if for each  $i \in I$   $\Omega_i$  is additive,
3. A class  $\mathcal{K}$  of BAOs is completely additive, if every  $\mathcal{B} \in \mathcal{K}$  is completely additive.

### 3.1 Atom structures and canonical extensions

**Definition 3.** Let  $I$  be an index set and  $\{\Omega_i\}_{i \in I}$  a set of function symbols

1. An atom structure is a relational structure  $\mathcal{F} = \langle W, \{R_i\}_{i \in I} \rangle$  such that  $R_i$  is a  $n+1$ -ary relation symbol, if  $\Omega_i$  is an  $n$ -ary function symbol,
2. Let  $\mathcal{B}$  be an atomic BAO of the signature  $I$ , the atom structure of  $\mathcal{B}$ , written as  $\mathbf{At}\mathcal{B}$ , is an atom structure  $\langle \mathbf{At}(\mathcal{B}), \{R_i\}_{i \in I} \rangle$  such that for each  $a, b_0, \dots, b_{n+1} \in \mathbf{At}(\mathcal{B})$  and for each  $i \in I$

$$\mathbf{At}\mathcal{B} \models R_i(a, b_0, \dots, b_{n+1}) \text{ iff } \mathcal{B} \models a \leq \Omega_i(b_0, \dots, b_{n+1})$$

3. Let  $\mathcal{F} = \langle W, \{R_i\}_{i \in I} \rangle$  be an atom structure, the complex algebra of  $\mathcal{F}$ , written as  $\mathbf{Cm}\mathcal{F}$ , is a BAO  $\langle \mathcal{P}(W), \cup, -, \emptyset, W, \{\Omega_{R_i}\}_{i \in I} \rangle$  such that for all  $X_0, \dots, X_{n-1} \subseteq W$  and for each  $i \in I$

$$\Omega_{R_i}(X_0, \dots, X_{n-1}) = \{a \in W \mid \exists b_0 \in X_0 \dots \exists b_{n-1} \in X_{n-1} \mathcal{F} \models R_i(a, b_0, \dots, b_{n-1})\}$$

The following duality is due to Thomason [5].

**Fact 1.**

1. Let  $\mathcal{B}$  be a complete atomic BAO, then  $\mathcal{B} \cong \mathbf{Cm}(\mathbf{At}(\mathcal{B}))$ ,
2. Let  $\mathcal{F}$  be an atom structure, then  $\mathcal{F} \cong \mathbf{At}(\mathbf{Cm}(\mathcal{B}))$ .

Let  $A$  be a non-empty subset of a Boolean algebra  $\mathcal{B}$ ,  $A$  is a *filter*, if  $A$  is closed under finite infima and upwardly closed.  $A$  is an *ultrafilter*, if it has no non-trivial extensions. That is, if  $A \subseteq A'$ , then  $A' = \mathcal{B}$ .

**Definition 4.** Let  $\mathcal{B} = \langle \mathcal{B}, +, -, 0, 1, \{\Omega_i\}_{i \in I} \rangle$  be a BAO and  $\mathbf{Uf}(\mathcal{B})$  the set of its ultrafilters. The ultrafilter frame of  $\mathcal{B}$  (or canonical frame) is a relational structure  $\mathcal{F}_{\mathcal{B}} = \langle \mathbf{Uf}(\mathcal{B}), R_{\Omega_i} \rangle$  such that for each ultrafilters  $\beta_0, \dots, \beta_{n-1}, \gamma$  one has

$$\mathbf{Uf}(\mathcal{B}) \models R_{\Omega_i}(\beta_0, \dots, \beta_{n-1}, \gamma) \text{ iff } \{\Omega_i(b_0, \dots, b_{n-1}) \mid b_0 \in \beta_0, \dots, b_{n-1} \in \beta_{n-1}\} \subseteq \gamma.$$

**Definition 5.** Let  $\mathcal{B}$  be a BAO, then

1. The canonical extension of  $\mathcal{B}$  is a complex algebra of the canonical frame  $\mathbf{Cm}(\mathcal{F}_{\mathcal{B}})$  denoted as  $\mathcal{B}^+$ ,
2. The class of BAOs is canonical, if it is closed under canonical extensions.

**Theorem 1.** Let  $\mathcal{A}, \mathcal{B}$  be BAOs,

1. There exists  $\iota : \mathcal{A} \hookrightarrow \mathcal{A}^+$  such that  $\iota : a \mapsto \{\gamma \in \mathbf{Uf}(\mathcal{A}) \mid a \in \gamma\}$ .
2. If  $i : \mathcal{A} \hookrightarrow \mathcal{B}$ , then this embedding might be extended to the embedding  $i^+ : \mathcal{A}^+ \hookrightarrow \mathcal{B}^+$

**Fact 2.**

### 3.2 (Representable) cylindric algebras and cylindric set algebras

**Definition 6.**

A relation algebra is an algebra  $\mathcal{R} = \langle R, 0, 1, +, -, ;, \smile, \mathbf{1}' \rangle$  such that  $\langle R, 0, 1, +, - \rangle$  is a Boolean algebra and the following equations hold, for each  $a, b, c \in R$ :

1.  $a; (b; c) = (a; b); c$
2.  $(a + b); c = (a; c) + (b; c)$
3.  $a; \mathbf{1}' = a$
4.  $a^{\smile\smile} = a$
5.  $(a + b)^{\smile} = a^{\smile} + b^{\smile}$
6.  $(a; b)^{\smile} = b^{\smile}; a^{\smile}$
7.  $a^{\smile}; -(a; b) \leq -b$

where  $a \leq b$  iff  $a + b = b$ . **RA** denotes the class of all relation algebras.

Cylindric algebras provide a generalisation of relation algebras for relations of an arbitrary arity. Let  $\alpha$  be an ordinal. Let  ${}^\alpha U$  be the set of all functions mapping  $\alpha$  to a non-empty set  $U$ . We denote  $x(i) = x_i$  for  $x \in {}^\alpha U$  and  $i < \alpha$ .

**Definition 7.**

1. A subset of  ${}^\alpha U$  is an  $\alpha$ -ry relation on  $U$ . For  $i, j < \alpha$ , the  $i, j$ -diagonal  $D_{ij}$  is the set of all elements of  $U$  such that  $y_i = y_j$ .  
If  $i < \alpha$  and  $X$  is an  $\alpha$ -ry relation on  $U$ , then the  $i$ -th cylindrification  $C_i X$  is the set of all elements of  $U$  that agree with some element of  $X$  on each coordinate except the  $i$ -th one. To be more precise,  $C_i X = \{y \in {}^\alpha U \mid \exists x \in X \forall i < \alpha (i \neq j \Rightarrow y_j = x_j)\}$ .
2. A cylindric set algebra of dimension  $\alpha$  is an algebra consisting of a set  $S$  of  $\alpha$ -ry relation on some base set  $U$  with the constants and operations  $0 = \emptyset$ ,  $1 = {}^\alpha U$ ,  $\cap$ ,  $-$ , the diagonal elements  $\{D_{ij}\}_{i,j < \alpha}$ , the cylindrifications  $\{C_i\}_{i < \alpha}$ .  
A generalised cylindric set algebra of dimension  $\alpha$  is a subdirect of cylindric algebras that have dimension  $\alpha$ .
3. A cylindric algebra of dimension  $\alpha$  is an algebra  $\mathcal{C} = \langle \mathcal{B}, \{c_i\}_{i < \alpha}, \{d_{ij}\}_{i,j < \alpha} \rangle$  such that
  - $\mathcal{B}$  is a Boolean algebra, for each  $i, j < \alpha$   $c_i$  is an operator and  $d_{ij} \in \mathcal{B}$
  - For each  $i < \alpha$ ,  $a \leq c_i a$ ,  $c_i(a \wedge c_i b) = c_i a \wedge c_i b$  and  $d_{ii} = 1$
  - For every  $i, j < \alpha$ ,  $c_i c_j a = c_j c_i a$
  - If  $k \neq i, j < \alpha$ , then  $d_{ij} = c_k(d_{ij} \wedge d_{jk})$
  - If  $i \neq j$ , then  $c_i(d_{ij} \wedge a) \wedge c_i(d_{ij} \wedge -a) = 0$

**CA $_\alpha$**  is the class of all cylindric algebras of dimension  $\alpha$

4. An  $\alpha$ -dimensional cylindric algebra  $C$  is representable, if it is isomorphic to a generalised cylindric set algebra of dimension  $\alpha$ . Such isomorphism is a representation of  $C$ .

**RCA $_\alpha$**  is the class of all representable cylindric algebras that have dimension  $\alpha$ . In particular, we are interested in the case when  $\alpha = \omega$ .

**Definition 8.** Given a cylindric algebra of dimension  $\alpha$   $C$ , let  $x$  be a term of its signature, the substitution operator  $s_j^i$  have the following definition:

$$s_j^i x = \begin{cases} x, & \text{if } i = j \\ c_i(d_{ij} \wedge x), & \text{otherwise} \end{cases}$$

It is well known that  $\mathbf{RCA}_\alpha$  is a variety,  $\mathbf{RCA}_\alpha$  ( $\alpha \leq 2$ ) is finitely axiomatisable and  $\mathbf{RCA}_\alpha$  ( $2 < \alpha < \omega$ ) has no finite axiomatisation, see [2].

Let  $\mathcal{A} \in \mathbf{C}_\omega$ , then  $\mathcal{A}$  has a *complete representation*, if this representation preserves all existing suprema.

Let us concretise the definition of a canonical extension for  $\alpha$ -dimensional cylindric algebras:

**Definition 9.** Let  $\mathcal{C} = \langle C, +, -, 0, 1, \{d_{ij}\}_{i,j < \alpha}, \{c_i\}_{i < \alpha} \rangle$   $\mathcal{A}$  be a BAO of type  $\mathbf{CA}_\alpha$ . Let  $\mathbf{Uf}(\mathcal{C})$  be the set of all ultrafilters of  $\mathcal{B}\mathcal{C}$ , the Boolean part of  $\mathcal{C}$ .

Let us define  $\mathbf{C}_i : \mathbf{Uf}(\mathcal{C}) \rightarrow \mathbf{Uf}(\mathcal{C})$  for each  $i, j < \alpha$  as

1.  $\mathbf{C}_i \mathcal{X} = \{\mathcal{F} \in \mathbf{Uf}(\mathcal{C}) \mid \exists \mathcal{F}' \in \mathbf{Uf}(\mathcal{C}) (a \in \mathcal{F} \Rightarrow c_i a \in \mathcal{F}'R)\},$
2.  $D_{ij} = \{\mathcal{F} \in \mathbf{Uf}(\mathcal{C}) \mid d_{ij} \in \mathcal{F}\}.$

The structure  $\mathcal{C}^+ = \langle \mathbf{Uf}(\mathcal{C}), \cup, -, \emptyset, C, \mathbf{C}_{i < \alpha}, \{D_{ij}\}_{i,j < \alpha} \rangle$  is called the canonical extension of  $\mathcal{C}$ .

Let us discuss the connection between representability and canonical extensions.

The following definitions and facts are due to Henkin, Monk, and Tarski [1].

Let  $\mathcal{A} \in \mathbf{CA}_\alpha$  and  $x \in \mathcal{A}$ . Recall that the *dimension of  $x$*  is the set of all ordinals  $\gamma < \alpha$  such that  $c_\gamma x \neq x$ . More formally,

$$\Delta x = \{\gamma \mid \gamma < \alpha \text{ \& } c_\gamma x \neq x\}$$

Let us discuss some metamathematical intuitions standing behind the notion of a dimension. Let  $\Theta$  be a first-order theory and  $\mathcal{C}/\equiv_\Theta$  its Lindenbaum-Tarski algebra. Let  $\varphi$  be a formula in the signature of  $\Theta$ . Then  $\Delta(\varphi/\Theta)$  consists of all  $\kappa < \alpha$  such that  $\exists x_\kappa \varphi \leftrightarrow \varphi$  is not valid in  $\Theta$ . That is,  $\Delta(\varphi/\Theta)$  contains ordinals  $\kappa$  for which  $x_\kappa$  is free in  $\varphi$ . Moreover,  $\Delta(\varphi/\Theta)$  consists only of those ordinals for which  $x_\kappa$  is free in every  $\psi \in \varphi/\Theta$ .

In particular, an element  $x$  is called *zero-dimensional* if  $\Delta x = 0$ . Zero-dimensional elements reflect equivalence classes of sentences in the Lindenbaum-Tarski algebra of a given first-order theory. Thus, the set of zero-dimensional elements form a Boolean algebras of sentences associated with  $\Theta$ .

**Definition 10.** Let  $\mathcal{A}$  be an  $\alpha$ -dimensional cylindric algebra. Let  $\alpha$  be an ordinal and  $\Gamma$  a subset  $\alpha$ , then an element  $x \in \mathcal{A}$  is  $\Gamma$ -closed if  $\Delta x \cdot \Gamma = \emptyset$ . Alternatively,  $x$  is a  $\Gamma$ -cylinder.

$\text{Cl}_\Gamma \mathcal{A}$  is the set of all  $\Gamma$ -closed elements.

Metamathematically,  $\Gamma$ -closed elements reflect universal closures (is it correct?).

Let  $\mathcal{C} = \langle C, +, -, 0, 1, \{d_{ij}\}_{i,j < \beta}, \{c_i\}_{i < \beta} \rangle$  be a  $\beta$ -dimensional cylindric algebra and  $\alpha \leq \beta$  an ordinal. The  $\alpha$ -th reduct of  $\mathcal{C}$ , denoted as  $\mathfrak{Rd}_\alpha \mathcal{C}$ , is an algebra having the form

$$\mathfrak{Rd}_\alpha \mathcal{C} = \langle C, +, -, 0, 1, \{d_{ij}\}_{i,j < \alpha}, \{c_i\}_{i < \alpha} \rangle$$

$\mathcal{B}$  is a subreduct of  $\mathcal{C}$ , denoted as  $\mathcal{B} \sqsubseteq^r \mathcal{C}$ , if  $\mathcal{B} \subseteq \mathfrak{Rd}_\gamma \mathcal{C}$  for some  $\gamma \leq \beta$ .

**Definition 11.** Let  $\mathcal{C}$  be a  $\beta$ -dimensional cylindric algebra and  $\alpha$  an ordinal such that  $\alpha \leq \beta$ . The neat  $\alpha$ -reduct of  $\mathcal{C}$ , denoted as  $\mathfrak{Nr}_\alpha \mathcal{C}$ , is the subalgebra  $\mathcal{A}$  of  $\mathfrak{Rd}_\alpha \mathcal{C}$  with  $\mathcal{A} = \text{Cl}_\kappa \mathcal{C}$  where  $\alpha + \kappa = \beta$ .

Let  $\mathbb{K}$  be a class of  $\beta$ -dimensional cylindric algebras, then we put

$$\mathbf{Nr}_\alpha \mathbb{K} = \{\mathfrak{Nr}_\alpha \mathcal{C} \mid \mathcal{C} \in \mathbb{K}\}$$

An algebra  $\mathcal{B}$  is a neat subreduct of  $\mathcal{C}$ , or  $\mathcal{B}$  is neatly embeddable to  $\mathcal{C}$  if there exists an ordinal  $\gamma \leq \alpha$  such that  $\mathcal{C} \subseteq \mathfrak{Rd}_\gamma \mathcal{B}$ .

One may define neat reducts alternatively as follows. Let  $\mathcal{C}$  be a  $\beta$ -dimensional cylindric algebra and  $\alpha$  an ordinal such that  $\alpha \leq \beta$ . The neat  $\alpha$ -reduct of  $\mathcal{C}$  is the  $\alpha$ -dimensional cylindric algebra having the form

$$\mathfrak{Nr}_\alpha \mathcal{C} = \langle \{a \in \mathcal{C} \mid \forall j(\alpha \leq j \ \& \ j < \beta \Rightarrow c_j a = a)\}, +, -, 0, 1, \{d_{ij}\}_{i,j < \alpha}, \{c_\gamma\}_\gamma \rangle$$

We will adapting the following proof of the fact that **RRA** is canonical<sup>1</sup> to our case. This proof is due to Monk, but that was describe in McKenzie's thesis [4].

1. A relation algebra  $\mathcal{A}$  is representable iff  $\mathcal{A}$  neatly embeds to some omega-dimensional cylindric algebra,
2. If  $\mathcal{A}$  neatly embeds in  $\mathcal{A}$  then  $\mathcal{A}^+$  neatly embeds in  $\mathcal{B}^+$ ,
3.  $\mathbf{CA}_\alpha$  is closed under canonical extensions,
4. Voilà.

**Definition 12.** Let  $\mathcal{C} \in \mathbf{CA}_\alpha$ , where  $\alpha \geq 3$ . The relation algebra reduct of  $\mathcal{C}$ , written as  $\mathfrak{Ra}(\mathcal{C})$ , is the algebra having the form

$$\langle \text{dom}(\mathfrak{Nr}_2(\mathcal{C})), 0, 1, +, -, \mathbf{1}', \smile, ; \rangle$$

where:

1.  $+$ ,  $-$ ,  $0$ , and  $1$  are defined as usual in  $\mathcal{C}$ ,
2.  $\mathbf{1}' = d_{01} \in \mathfrak{Nr}_2(\mathcal{C})$ ,
3.  $r \smile = s_0^2 s_1^0 s_2^1 r$  for  $r \in \mathfrak{Nr}_2(\mathcal{C})$ ,
4. Let  $r, s \in \mathfrak{Nr}_2(\mathcal{C})$ , then  $r; s = c_2(s_2^1 r \cdot s_2^0 s)$

Moreover,  $\mathfrak{Nr}_\beta(\mathcal{C})$  and  $\mathfrak{Ra}(\mathcal{C})$  are closed under these operations. There is also the following fact by due to Henkin, Monk, and Tarski [2]:

**Theorem 2.** Let  $\mathcal{C} \in \mathbf{CA}_\alpha$  for  $\alpha \geq 4$ , then  $\mathfrak{Ra}(\mathcal{C})$  is a relation algebra.

The following characterisation results are by Henkin, Monk, and Tarski [2] as well:

**Theorem 3.**

1.  $\mathbf{RA} = \mathbf{SRA} \mathbf{CA}_4$ ,
2.  $\mathbf{RRA} = \bigcap_{3 \leq n < \omega} \mathbf{SRA} \mathbf{CA}_n = \mathbf{SRA} \mathbf{CA}_\alpha$ , where  $\alpha$  is an infinite ordinal.

Let  $\mathcal{C} \in \mathbf{CA}_\alpha$ , then  $\mathcal{R} \in \mathbf{RA}$  neatly embeds to  $\mathcal{C}$ , if  $\mathcal{R}$  is isomorphic to some subalgebra of  $\mathfrak{Ra}(\mathcal{C})$ .

---

<sup>1</sup>This idea is by Ian Hodkinson

**Theorem 4.**  *$\mathbf{RRA}$  is closed under canonical extensions.*

*Proof.* Let  $\mathbf{R} \in \mathbf{RRA}$ . By the second item of 3, every representable relation algebra is isomorphic to some subalgebra of the relation algebra reduct  $\mathfrak{Ra}\mathcal{C}$  for some  $\mathcal{C} \in \mathbf{CA}_\omega$ . But neat embeddings respect canonical extensions, so if  $\mathbf{R} \hookrightarrow_n \mathcal{C}$ , so is  $\mathbf{R}^+ \hookrightarrow_n \mathcal{C}^+$ .  $\mathbf{CA}_\alpha$  is closed under canonical extensions, so is  $\mathbf{RRA}$ .  $\square$

**Lemma 1** (Henkin, Monk, Tarski).

*Let  $\mathcal{A}$  be a BAO of type  $\mathbf{CA}_\alpha$  and  $\mathcal{B}$  be a  $\beta$ -dimensional cylindric algebra such that  $\beta \leq \alpha$  and  $\mathcal{A}$  neatly embeds to  $\mathcal{B}$  by a complete embedding. Then  $\mathcal{A}^+$  neatly embeds to  $\mathcal{B}^+$  by a complete embedding.*

*Proof.*  $\square$

**Theorem 5** (This assumption is by Ian Hodkinson).

*Let  $\mathcal{A}$  be a BAO of type  $\mathbf{CA}_\omega$  such that  $\mathcal{A}$  neatly embeds into  $\mathbf{CA}_{\omega+\omega}$  by a complete embedding. Then  $\mathcal{A}$  is completely representable as  $\mathbf{CA}_\omega$ .*

*Proof.* Hmmmmm, I believe so.  $\square$

Lemma 1 and Theorem 5 imply the following theorem.

**Theorem 6.** *Let  $\mathcal{C} \in \mathbf{RCA}_\omega$ , then  $\mathcal{C}^+ \in \mathbf{RCA}_\omega$ . That is,  $\mathbf{RCA}_\omega$  is closed under canonical extensions.*

*Proof.* :monkahmm:  $\square$

## References

- [1] L Henkin, J D Monk, and A Tarski. *Cylindric Algebras Part I*. North-Holland, 1971.
- [2] Leon Henkin, J. Donald Monk, and Alfred Tarski. Cylindric algebras. part ii. *Journal of Symbolic Logic*, 53(2):651–653, 1988.
- [3] Robin Hirsch and Ian Hodkinson. *Relation algebras by games*. Elsevier, 2002.
- [4] Ralph McKenzie. The representation of relation algebras. *PhD Thesis, University of Colorado*, 1968.
- [5] Steven K Thomason. Categories of frames for modal logic. *The journal of symbolic logic*, 40(3):439–442, 1975.