

# Mechanics 1

## Session 3 – Newton's Laws of Motion: Rethinking Forces

---

DR BEN HANSON

1

MECHANICS 1 – NEWTON'S LAWS OF MOTION

## Last Lecture

Trajectories

---

### We learned:

- The physics of projectile trajectories
- How to represent a trajectory as a vector equation
- How to represent the components of the vector using kinematic equations

### You should be able to:

- Describe the physical path taken by a projectile in terms of  $(x, y)$  coordinates
- Describe the physical path taken by a projectile in terms of a position vector  $\underline{r}$
- Calculate the position of a projectile at some time  $t$ , given its initial position, velocity and acceleration

DR BEN HANSON

2

# This Lecture

---

Question: What causes acceleration (and therefore motion)?

Question: What is a force?



DR BEN HANSON

3

# This Lecture

---

Newton's Laws of Motion

**We will learn:**

- The conceptual importance of Newton's 3 Laws of Motion
- Real-world examples of motion for which each of these laws applies
- How to mathematically analyse multiple forces acting simultaneously

**You will be able to:**

- Describe each of Newton's 3 Laws of Motion, with examples to illustrate
- Draw force diagrams for various physical systems
- Mathematically "resolve" force vectors into perpendicular directions

DR BEN HANSON

4

# Newton's Laws of Motion

A Conceptual Journey

---

5

MECHANICS 1 – NEWTON'S LAWS OF MOTION

## Newton's Laws of Motion

A Conceptual Journey

---

**Newton's Law's of Motion take us from a kinematic perspective to a mechanical perspective**

The development of the concept of “force” as a physical, measurable quantity, consistent between different types of interaction (mechanical, electromagnetic, gravitation etc) is a profound moment in the history of physics. So let's walk the path again together...


Disclaimer: This is not going to be historically accurate. It's a conceptual story

DR BEN HANSON

6

# Newton's Laws of Motion

Kinematics




DR BEN HANSON

7

# Newton's Laws of Motion

Kinematics



We can see that things move. But why?  
What is the "cause" of that motion?

DR BEN HANSON

8

# Newton's Laws of Motion

Kinematics

Imagine an object, sliding across a frozen surface...



1. First observation
  - a) Objects move, and slow down
  - b) The thicker or rougher the external surface, the slower they slow down
2. Second Observation
  - a) Objects move, and change direction
  - b) The sharper the turn or larger the contact angle, the greater the change of direction

**Objects appear to keep moving unless there is "something" to stop them. Let's call that something a "force".**

<https://www.youtube.com/watch?v=Sic7CckO-js>

DR BEN HANSON

9

# Newton's Laws of Motion

Newton's First Law

**A body remains at rest, or in motion at a constant speed in a straight line, unless acted upon by a force.**

DR BEN HANSON

10

# Newton's Laws of Motion

## Newton's First Law

**A body remains at rest, or in motion at a constant speed in a straight line, unless acted upon by a force.**

Forces change an object's speed and direction? Force must be a vector!

DR BEN HANSON

11

## Task 1

Forces Acting on Objects

12

# Task 1

## Forces Acting on Objects

### Tasks:

For the following examples, describe with your neighbours in which direction the forces are acting:

1. A trailer is being pulled by a car along the M62 motorway.
  1. The car accelerates from 55mph to 60mph, the speed limit. In what direction is force acting on the trailer?
  2. The car and trailer are moving at a steady 60mph. In what direction is force acting on the trailer?
2. Imagine that you are in a car and a dog runs out into the road. You have to perform an emergency stop, bringing your car to a full stop as quickly as possible.
  1. In what direction would the force be acting on the car? What is the source of that force?
  2. In what direction would the force be acting on you, the driver? What is the source of that force?
3. A child is sliding down a slide. The slide starts off at an angle of  $45^\circ$  to the ground, but curves to be  $15^\circ$  at the bottom.
  1. What are the forces acting on the child at the top of the slide?
  2. What are the forces acting on the child at the bottom of the slide?
  3. What force was acting on the child to cause them to change from a direction of  $45^\circ$  to  $15^\circ$ ?

DR BEN HANSON

13

# Newton's Laws of Motion

## Mass & Impulse

14

# Newton's Laws of Motion

Mass

Imagine pushing a pram...



DR BEN HANSON

15

# Newton's Laws of Motion

Mass

But what about...a plane!?



**Heavier objects are harder to  
accelerate / decelerate!**

<https://www.youtube.com/watch?v=B-4NuNelk3Q>

DR BEN HANSON

16



# Newton's Laws of Motion

## Impulse

Imagine pushing for a short amount of time...



- From this video:  
Large force => Large momentum change
- From the previous videos:  
More contact time => More momentum change

Define Impulse,  $\underline{I} = \underline{F}_N \Delta t$

[https://www.youtube.com/watch?v=1-a4YJ\\_JZqo](https://www.youtube.com/watch?v=1-a4YJ_JZqo)

DR BEN HANSON

17

# Newton's Laws of Motion

## Newton's Second Law

**The impulse acting on an object is equal to its change in momentum**

$$\underline{I} = \Delta \underline{p}$$

$$\rightarrow \Delta \underline{p} = \underline{F}_N \Delta t$$

DR BEN HANSON

18

# Newton's Laws of Motion

## Newton's Second Law

**The impulse acting on an object is equal to its change in momentum**

Impulse equals change in momentum,

$$\underline{F}_N \Delta t = \Delta \underline{p}$$

Rearrange,

$$\underline{F}_N = \frac{\Delta \underline{p}}{\Delta t}$$

Take limit,

$$\underline{F}_N = \frac{d\underline{p}}{dt}$$

DR BEN HANSON

19

# Newton's Laws of Motion

## Newton's Second Law

**The impulse acting on an object is equal to the change in its momentum**

Take limit,

$$\underline{F}_N = \frac{d\underline{p}}{dt}$$

Substitute,

$$\underline{F}_N = \frac{d}{dt}(m\underline{v})$$

Constant mass,

$$\underline{F}_N = m \frac{d}{dt}(\underline{v})$$

Hence,

$$\underline{F}_N = m\underline{a}$$

DR BEN HANSON

20

# Newton's Laws of Motion

## Newton's Second Law

**If the mass is constant, the net force on an object is equal to its mass multiplied by its acceleration**

$$\underline{F}_N = m\underline{a}$$

A force can be defined simply as the cause of acceleration! Similarly, if an object is accelerating, there must be a non-zero force acting on it!

DR BEN HANSON

21

# Newton's Laws of Motion

## Net Force & Reaction Forces

22

# Newton's Laws of Motion


## Net Force & Reaction Forces

**Scenario:** A box is at rest on the floor. It is not accelerating.

**Question:** What forces are acting upon it, and why?

1. The box is exerting a force on the ground
2. In response, the ground is exerting a force (of the same magnitude) in the opposite direction on the box.

Interesting...



DR BEN HANSON

23

# Newton's Laws of Motion


## Net Force & Reaction Forces

**Scenario:** A box is at being pulled by someone. They are exerting a constant force  $F_p$  on the box via a rope. The box is moving at a constant speed in the horizontal direction.

**Question 1:** What forces are acting upon it, and why?

**Question 2:** Is  $F_p > F_f$ ,  $F_p < F_f$  or  $F_p = F_f$

**Question 3:** What forces are acting upon the person, and why?



DR BEN HANSON

24

# Newton's Laws of Motion

## Newton's Third Law

**If body A applies a force to body B, then body B applies an equal and opposite force to body A**

DR BEN HANSON

25

# Newton's Laws of Motion

## Newton's Laws

1. A body remains at rest, or in motion at a constant speed in a straight line, unless acted upon by a force.
2. The impulse acting on an object is equal to its change in momentum
  - a)  $\Rightarrow$  The net force on an object is equal to the rate of change of its momentum:  $\underline{F}_N = \frac{dp}{dt}$
  - b)  $\Rightarrow$  If the mass is constant, the net force on an object is equal to its mass multiplied by its acceleration:  $\underline{F}_N = m\underline{a}$
3. If body A applies a force to body B, then body B applies an equal and opposite force to body A

DR BEN HANSON

26

# Newton's Laws of Motion

## Gravitation

---

27

MECHANICS 1 – NEWTON'S LAWS OF MOTION

# Newton's Laws of Motion

## Newton's Gravitation

---

Wait a minute...

1. Newton's Second Law implies that, for the same force, lighter objects would accelerate faster, and heavier objects slower.
2. We can observe this experimentally when objects move parallel to the ground

**Let's do an  
experiment!**

So why does everything fall at the same rate  
(as originally shown by Galileo)?

DR BEN HANSON

28

# Newton's Laws of Motion

## Newton's Gravitation

Now, imagine being sat under an apple tree...



<https://www.youtube.com/watch?v=h48BWDeBLno>

DR BEN HANSON

29

# Newton's Laws of Motion

## Newton's Gravitation

**Scenario:** An apple falls on your head. You have recently discovered that  $\underline{F}_N = m\underline{a}$ . This equation says that the acceleration of an object always implies the existence of a force that is causing the acceleration. Further, this acceleration, for a given force, is inversely proportional to the mass ( $\underline{a} = \frac{\underline{F}_N}{m}$ ). However, for this ubiquitous downwards force of gravity ( $\underline{a} = \frac{\underline{F}_G}{m}$ ), acceleration is somehow independent of mass.

**Question:** How can this be true? *Hint: Think about the mathematics. How can an equation be independent of a variable?*

DR BEN HANSON

30

# Newton's Laws of Motion

## Newton's Gravitation

**Scenario:** An apple falls on your head. You have recently discovered that  $F_N = m\underline{a}$ . This equation says that the acceleration of an object always implies the existence of a force that is causing the acceleration. Further, this acceleration, for a given force, is inversely proportional to the mass ( $\underline{a} = \frac{F_N}{m}$ ). However, for this ubiquitous downwards force of gravity ( $\underline{a} = \frac{F_G}{m}$ ), acceleration is somehow independent of mass.

**Question:** How can this be true? *Hint: Think about the mathematics. How can an equation be independent of a variable?*

**Acceleration is inversely proportional to mass, but everything falls at the same rate. Gravitational force must therefore also be proportional to mass!**

We'll do Newtonian gravity properly next week 😊

DR BEN HANSON

31

## Task 2

Conceptually Understanding Newton's Laws

32



## Task 2

### Conceptually Understanding Newton's Laws

**Task:** With your neighbours, discuss how each of Newton's Laws apply to various every day situations.

**Examples:**

1. When a car is accelerating / decelerating, what are the forces acting on the car? In what direction must they be acting.
2. If a bus and a car are moving at the same speed, what can we say about the net force on each of them?
3. When the floor outside is icy, we sometimes slip (and maybe fall ☹). Using Newton's First Law, explain why this is.
4. When you throw a ball (say, a football), it accelerates from a speed of zero to a speed  $|u|$  before it leaves your hand. Newton's Third Law says the the ball exerts the same force on you as you exerted on it. Explain why you do not go flying backwards at the same speed as the ball.

DR BEN HANSON

33

## Mechanical Equilibrium

and also Resolving Forces

34

# Mechanical Equilibrium

## Definition

Mechanical equilibrium is when the net force vector on an object is zero (i.e. zero in every direction)

$$\underline{F}_{Net} = \sum_i \underline{F}_i = \underline{0}$$

As such, the object may either be moving at a constant velocity or not moving at all (i.e. a constant velocity  $\underline{v} = \underline{0}$ )

DR BEN HANSON


35

# Mechanical Equilibrium

## Force Diagrams

**Scenario:** A box is at being pulled by someone. They are exerting a constant force  $\underline{F}_p$  on the box via a rope. The box is moving at a constant speed in the horizontal direction.

**Question 1:** Is this object in mechanical equilibrium?



DR BEN HANSON

36


# Mechanical Equilibrium

## Force Diagrams

**Scenario:** The person pulling the box stumbles and releases the rope.

**Question 1:** Is this object in mechanical equilibrium?

**Question 2:** If the object has a mass of  $50\text{kg}$  and was initially moving at  $0.75\text{m/s}$ , and  $\underline{F}_f = 30\text{N}$ , how long will the box take to stop moving?



DR BEN HANSON

37

# Mechanical Equilibrium

## Force Diagrams

**Scenario:** The person pulling the box stumbles and releases the rope.

**Question 2:** If the object has a mass of  $50\text{kg}$  and was initially moving at  $0.75\text{m/s}$ , and  $\underline{F}_f = 30\text{N}$ , how long will the box take to stop moving?

$$\underline{F}_{Net} = m\underline{a}$$


$$\rightarrow F_x = ma_x, \quad a_x = \frac{F_x}{m} = -0.6\text{ms}^{-2}$$

$$\text{1D SUVAT,} \quad v_x(t) = u_x + a_x t$$

$$v_x(t) = 0, \quad 0 = 0.75 - 0.6t$$

Solve,

$$t = 1.25\text{s}$$



DR BEN HANSON

38

# Mechanical Equilibrium

## Resolving Forces


**Scenario:** A box is at being pulled uphill by someone. They are exerting a constant force  $\underline{F}_p$  on the box via a rope. The box is not moving at all.

**Question 1:** What forces are acting on this object?

**Question 2:** Is this object in mechanical equilibrium?

**Question 3:** If the ramp is at an angle  $\theta$  to the horizontal, derive an equation for the reaction force magnitude,  $|\underline{R}|$ .

**Question 4:** Using Newton's Second Law, show that  $|\underline{F}_p| = |\underline{F}_f| + |\underline{F}_G| \sin(\theta)$



DR BEN HANSON


39

# Mechanical Equilibrium

## Resolving Forces

**Scenario:** A box is at being pulled uphill by someone. They are exerting a constant force  $\underline{F}_p$  on the box via a rope. The box is not moving at all.

**Question 3:** If the ramp is at an angle  $\theta$  to the horizontal, derive an equation for the reaction force magnitude,  $|\underline{R}|$ .



DR BEN HANSON


40

# Mechanical Equilibrium

## Resolving Forces

**Scenario:** A box is at being pulled uphill by someone. They are exerting a constant force  $F_p$  on the box via a rope. The box is not moving at all.

**Question 3:** If the ramp is at an angle  $\theta$  to the horizontal, derive an equation for the reaction force magnitude,  $|R|$ .



DR BEN HANSON

41


# Mechanical Equilibrium

## Resolving Forces

**Scenario:** A box is at being pulled uphill by someone. They are exerting a constant force  $F_p$  on the box via a rope. The box is not moving at all.

**Question 3:** If the ramp is at an angle  $\theta$  to the horizontal, derive an equation for the reaction force magnitude,  $|R|$ .

$$|R| = |F_G| \cos(\theta)$$



DR BEN HANSON

42

# Mechanical Equilibrium

## Resolving Forces

**Scenario:** A box is at being pulled uphill by someone. They are exerting a constant force  $F_p$  on the box via a rope. The box is not moving at all.

**Question 4:** Using Newton's Second Law, show that  $|\underline{F}_p| = |\underline{F}_f| + |\underline{F}_G| \sin(\theta)$

No motion parallel to ramp,


$$\underline{F}_{j'} = m \underline{a}_{j'} = 0$$

Resolve all forces and sum,

$$|\underline{F}_f| + |\underline{F}_G| \sin(\theta) - |\underline{F}_p| = 0$$

Rearrange,

$$|\underline{F}_p| = |\underline{F}_f| + |\underline{F}_G| \sin(\theta)$$



DR BEN HANSON

43

## Forces

Just a Small Philosophical Thing

44

# Forces

And Philosophy (it's important)

1. A body remains at rest, or in motion at a constant speed in a straight line, unless acted upon by a force.
2. The impulse acting on an object is equal to its change in momentum
  - a) => The net force on an object is equal to the rate of change of its momentum:  $\underline{F}_N = \frac{dp}{dt}$
  - b) => If the mass is constant, the net force on an object is equal to its mass multiplied by its acceleration:  $\underline{F}_N = m\underline{a}$
3. If body A applies a force to body B, then body B applies an equal and opposite force to body A

**Question: Do forces “exist”?**

DR BEN HANSON