

# Mechanics 1

## Session 11 – Momentum and Its Conservation

---

DR BEN HANSON

1

MECHANICS 1 – MOMENTUM

## Last Lecture

Energy Conservation

---

### We learned:

- How relative motion works in classical physics
- That acceleration, and therefore force, is sometimes measured differently in different reference frames
- To derive the location of the centre of mass of a collection of particles

### You should be able to:

- Calculate the relative speed between two moving objects
- Calculate the centre of mass of a collection of particles
- Calculate the velocity (and potentially the acceleration) of a collection of particles using its centre of mass dynamics

DR BEN HANSON

2

# This Lecture

## Momentum

---

**We will:**

- Understand what momentum is conceptually
- See that momentum is a vector
- Understand why momentum is always conserved following collisions
- See that everything becomes mathematically easier in the centre of mass reference frame

**You will be able to:**

- Use the concept of momentum conservation to calculate the subsequent kinetic properties (velocities) following a collision

DR BEN HANSON

3

# Momentum

What is it?

---

4

# Momentum

What is it?

The momentum of an object is, simply, the mass multiplied by the velocity.


But because it is a conserved quantity, it has a wealth of uses in physics calculations.

DR BEN HANSON

5

# Momentum

Why is it Conserved?




DR BEN HANSON

6

# Momentum

## Why is it Conserved?



Impulse over small (collision) time,

$$I = \vec{F}\Delta t$$

Impulse is also momentum change,

$$I = \Delta \vec{p}$$

Combine,

$$\Delta \vec{p} = \vec{F}\Delta t$$

Object 1,

$$\Delta \vec{p}_1 = \vec{F}_{21}\Delta t$$

Object 2,

$$\Delta \vec{p}_2 = \vec{F}_{12}\Delta t$$

Total momentum change,

$$\Delta \vec{p}_T = \Delta \vec{p}_1 + \Delta \vec{p}_2$$


$$\Delta \vec{p}_T = \vec{F}_{21}\Delta t + \vec{F}_{12}\Delta t$$

DR BEN HANSON

7

# Momentum

## Why is it Conserved?



Total momentum change,

$$\Delta \vec{p}_T = \vec{F}_{21}\Delta t + \vec{F}_{12}\Delta t$$

Factorise,

$$\Delta \vec{p}_T = (\vec{F}_{21} + \vec{F}_{12})\Delta t$$

Newton's 3<sup>rd</sup> Law,

$$\vec{F}_{21} = -\vec{F}_{12}$$

Hence,

$$\Delta \vec{p}_T = 0$$


The change in the total momentum of all objects after a collision,  $\Delta \vec{p}_T$ , is zero. Hence, momentum is conserved throughout collisions!

DR BEN HANSON

8

# Momentum

## The Full Equations



Hence,

$$\Delta \vec{p}_T = 0$$

Initially,

$$\vec{p}_T = m_1 \vec{u}_1 + m_2 \vec{u}_2$$

After collision,

$$\vec{p}_T = m_1 \vec{v}_1 + m_2 \vec{v}_2$$

Hence,

$$(m_1 \vec{v}_1 + m_2 \vec{v}_2) - (m_1 \vec{u}_1 + m_2 \vec{u}_2) = 0$$

$$m_1 \vec{u}_1 + m_2 \vec{u}_2 = m_1 \vec{v}_1 + m_2 \vec{v}_2$$

In general, for  $N$  colliding objects,

$$\sum_{i=1}^N m_i \vec{u}_i = \sum_{i=1}^N m_i \vec{v}_i$$

DR BEN HANSON

9

## Task 1

### Momentum Calculations

10

## Task 1

### Momentum Calculations

**Scenario:** Two children are playing marbles. A circle with radius  $R = 5\text{cm}$  is drawn on the ground in chalk, and the first child's marble, mass  $m_1 = 100\text{g}$  and radius  $R_1 = 0.75\text{cm}$ , is placed at the centre of the circle. The second child rolls their larger marble, mass  $m_2 = 250\text{g}$  and radius  $R_2 = 1.25\text{cm}$  toward the first and hits it. Just before it hits, the second marble has velocity  $\vec{u}_2 = 30\text{cms}^{-1}\hat{j}$ . Following the collision, the second marble has a new velocity  $\vec{v}_2 = 2.8\text{cms}^{-1}\hat{i} + 16.6\text{cms}^{-1}\hat{j}$ .

**Tasks:**

1. Calculate the velocity of the first marble after the collision.
2. If the ground has co-efficient of kinetic friction  $\mu_k = 0.15$ , will the first marble still be in the circle when it comes to a stop? *Hint: You can treat a rolling marble as a simple sliding marble for this question (i.e. friction acts as normal).*

DR BEN HANSON

11

## Task 2

### An Interesting Derivation with Energy Conservation

12

## Task 2

### An Interesting Derivation with Energy Conservation

When (kinetic) energy is also conserved during a collision, it is called an “elastic collision”.

**Tasks:**

1. By considering both conserved kinetic energy and momentum during a 1D collision between two objects with initial speeds  $u_1$  and  $u_2$ , and final speeds  $v_1$  and  $v_2$ , show that:

$$\frac{v_2 - v_1}{u_1 - u_2} = 1$$

DR BEN HANSON

13

## Task 3

### Newton's Cradle

14

## Task 3

### Newton's Cradle

**Tasks:**

1. Explain how this device works from fundamental principles. Use concepts such as impulse and momentum conservation.
2. Prove to yourself mathematically how this device works
3. Would this device continue to move forever? If not, why not?



DR BEN HANSON

15

# Momentum & Centre of Mass

## Frames of Reference

16



# Momentum & Centre of Mass

## Frames of Reference

The laws of physics are fundamentally the same in all inertial reference frames. But some are easier to deal with mathematically than others


Let's switch to the centre of mass reference frame and see what happens

DR BEN HANSON

17

# Momentum & Centre of Mass

## Frames of Reference



$$M_T \vec{r}_{CM} = \sum_{i=1}^N m_i \vec{r}_i$$

$$\vec{r}_{CM} = \frac{1}{M_T} \sum_{i=1}^N m_i \vec{r}_i$$

$$M_T \vec{v}_{CM} = \sum_{i=1}^N m_i \vec{v}_i$$

$$\vec{v}_{CM} = \frac{1}{M_T} \sum_{i=1}^N m_i \vec{v}_i$$

Recall that the centre of mass is the location at which all internal forces cancel out, and the only forces that appear to be acting on the CoM are external forces (i.e. forces from outside the system)

I wonder if that has anything to do with open and closed systems with regards to energy and work done<sup>1</sup>

We now see that the momentum of the centre of mass is equal to the sum over all momenta in the system

<sup>1</sup>It absolutely does. Consider it in your own time ☺

DR BEN HANSON

18

## Momentum & Centre of Mass

### Frames of Reference




Diagram showing a system of particles (blue circles) with a center of mass (black dot) and a vector  $\vec{r}_{CM}$  pointing from an origin to the center of mass.

$$M_T \vec{r}_{CM} = \sum_{i=1}^N m_i \vec{r}_i$$

$$\vec{r}_{CM} = \frac{1}{M_T} \sum_{i=1}^N m_i \vec{r}_i$$

$$M_T \vec{v}_{CM} = \sum_{i=1}^N m_i \vec{v}_i$$

$$\vec{v}_{CM} = \frac{1}{M_T} \sum_{i=1}^N m_i \vec{v}_i$$

Total momentum  
(static reference frame),

Total momentum  
(CoM reference frame),

$$\vec{p}_T = \sum_{i=1}^N m_i \vec{v}_i$$

$$\vec{p}_T = \sum_{i=1}^N m_i (\vec{v}_i - \vec{v}_{CM})$$

Expand,

$$\vec{p}_T = \left( \sum_{i=1}^N m_i \vec{v}_i \right) - \left( \sum_{i=1}^N m_i \vec{v}_{CM} \right)$$

Factorise,

$$\vec{p}_T = \left( \sum_{i=1}^N m_i \vec{v}_i \right) - \vec{v}_{CM} \left( \sum_{i=1}^N m_i \right)$$

DR BEN HANSON

19

## Momentum & Centre of Mass

### Frames of Reference




Diagram showing a system of particles (blue circles) with a center of mass (black dot) and a vector  $\vec{r}_{CM}$  pointing from an origin to the center of mass.

$$M_T \vec{r}_{CM} = \sum_{i=1}^N m_i \vec{r}_i$$

$$\vec{r}_{CM} = \frac{1}{M_T} \sum_{i=1}^N m_i \vec{r}_i$$

$$M_T \vec{v}_{CM} = \sum_{i=1}^N m_i \vec{v}_i$$

$$\vec{v}_{CM} = \frac{1}{M_T} \sum_{i=1}^N m_i \vec{v}_i$$

Factorise,

$$\vec{p}_T' = \left( \sum_{i=1}^N m_i \vec{v}_i \right) - \vec{v}_{CM} \left( \sum_{i=1}^N m_i \right)$$

Sum,

$$\vec{p}_T' = \left( \sum_{i=1}^N m_i \vec{v}_i \right) - \vec{v}_{CM} M_T$$

Solve,

$$\boxed{\vec{p}_T' = 0}$$

The total momentum of a system as measured from the centre of mass reference frame is zero! Imagine how easy this makes calculations of, say, galaxies colliding...

DR BEN HANSON

20

# A Little History

## Conserved Quantities in General

21

# A Little History

## Conserved Quantities in General

These conserved quantities in physics are useful. Mathematically, they let us set derivatives over time equal to zero, and thus simplify equations. Someone noted this use...



Emmy Noether (1882-1935)

Euler-Lagrange Equations,

$$\frac{dL}{dq} - \frac{d}{dt} \left( \frac{dL}{d\dot{q}} \right) = 0$$

More fundamental than  
Newton's laws...

$$\frac{dL}{dq} = 0,$$

$$\frac{d}{dt} \left( \frac{dL}{d\dot{q}} \right) = 0$$

22

## A Little History

### Conserved Quantities in General

These conserved quantities in physics are useful. Mathematically, they let us set derivatives over time equal to zero, and thus simplify equations. Someone noted this use...



Emmy Noether (1882-1935)

$$\frac{dL}{dq} = 0, \quad \frac{d}{dt} \left( \frac{dL}{d\dot{q}} \right) = 0$$

Integrate,

$$\boxed{\frac{dL}{d\dot{q}} = c}$$

This thing,  $\frac{dL}{d\dot{q}}$ , is constant over time. Noether showed that this is due to symmetry in the underlying laws of physics! More on this in 3<sup>rd</sup> year. For now...

DR BEN HANSON

23

## A Little History

### Conserved Quantities in General

These conserved quantities in physics are useful. Mathematically, they let us set derivatives over time equal to zero, and thus simplify equations. Someone noted this use...



Emmy Noether (1882-1935)

Integrate,

$$\frac{dL}{d\dot{q}} = c$$

$$L = K(\dot{q}) - U(q), \quad \frac{d}{d\dot{q}} \left( \frac{1}{2} m \dot{q}^2 - U(q) \right) = c$$

Differentiate,

$$\boxed{m\dot{q} = c}$$

$q$  is just a coordinate, like  $x$ .  $\dot{q}$ , then, is a velocity. Therefore, momentum is constant over time. It is conserved!

DR BEN HANSON

24