

Comment on “The incomplete beta function law for parallel tempering sampling of classical canonical systems” [J. Chem. Phys. 120, 4119 (2004)]

David A. Kofke

Citation: [The Journal of Chemical Physics](#) **121**, 1167 (2004); doi: 10.1063/1.1758211

View online: <http://dx.doi.org/10.1063/1.1758211>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/121/2?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

Erratum: “Estimation of the Heat Capacities of Organic Liquids as a Function of Temperature Using Group Additivity. An Amendment.” [J. Phys. Chem. Ref. Data 33, 1071 (2004)]

J. Phys. Chem. Ref. Data **34**, 39 (2005); 10.1063/1.1889439

[The incomplete beta function law for parallel tempering sampling of classical canonical systems](#)

J. Chem. Phys. **120**, 4119 (2004); 10.1063/1.1644093

[Comment on “Surface diffusion near the points corresponding to continuous phase transitions” \[J. Chem. Phys. 109, 3197 \(1998\)\]](#)

J. Chem. Phys. **111**, 11232 (1999); 10.1063/1.480479

[Comment on: “The hindered rotor density-of-states interpolation function” \[J. Chem. Phys. 106, 6675 \(1997\)\] and “The hindered rotor density- of-states” \[J. Chem. Phys. 108, 2314 \(1998\)\]](#)

J. Chem. Phys. **111**, 7163 (1999); 10.1063/1.480272

[Comment on “Hindered rotor density-of-states interpolation function” \[J. Chem. Phys. 106, 6675 \(1997\)\] and “The hindered rotor density-of-states” \[J. Chem. Phys. 108, 1748 \(1998\)\]](#)

J. Chem. Phys. **111**, 7161 (1999); 10.1063/1.480268



NEW Special Topic Sections

NOW ONLINE
Lithium Niobate Properties and Applications:
Reviews of Emerging Trends

AIP Applied Physics Reviews

COMMENTS

Comment on "The incomplete beta function law for parallel tempering sampling of classical canonical systems" [J. Chem. Phys. 120, 4119 (2004)]

David A. Kofke^{a)}

Department of Chemical and Biological Engineering, University at Buffalo,
The State University of New York, Buffalo, New York 14260-4200

(Received 26 March 2004; accepted 9 April 2004)

[DOI: 10.1063/1.1758211]

Recently, Kofke¹ has derived the following expression for the acceptance probability of swaps between two parallel-tempering replicas of inverse temperatures $\beta' > \beta$ [see Eq. (10) of Ref. 1]:

$$\bar{p}_{\text{acc}}(C) = 4 \frac{\Gamma(2C)}{[\Gamma(C)]^2} \frac{2C+1}{C} \int_0^{\beta'/\beta} \frac{\kappa^C}{(1+\kappa)^{2(C+1)}} d\kappa. \quad (1)$$

Equation (1) is derived under the assumption that the heat capacity of the system is almost constant and equal to C (in units of the Boltzmann's constant) on an inverse-temperature interval containing the values β and β' . It is also assumed that the sampled space is the phase space, which contains both the momentum and the position coordinates. If only the (d -dimensional) configuration space is considered, the heat capacity C must be replaced by $C_V = C - d/2$, the potential contribution to the heat capacity.

Standard properties of the Euler's gamma and beta functions together with the substitution $\theta = \kappa/(1+\kappa)$ lead to the following equivalent formulation of Eq. (1):

$$\bar{p}_{\text{acc}}(C) = \frac{2}{B(C+1, C+1)} \int_0^{1/(1+R)} \theta^C (1-\theta)^C d\theta, \quad (2)$$

where $R = \beta'/\beta > 1$ and where $B(C+1, C+1)$ denotes the respective Euler's beta function. Replacing C with $C-1$ in Eq. (2), one obtains the so-called empirical incomplete beta function law, which is Eq. (22) of Ref. 2. Since the replacement $C \rightarrow C+1$ does not change the asymptotic properties of Eq. (2) in the limit $C \rightarrow \infty$, it follows that Eq. (10) of Ref. 1 correctly predicts the $C^{-1/2}$ loss of efficiency upon the increase of the heat capacity (or dimensionality) of the sampled system. More precisely, to obtain a nonzero and nonunity limiting probability $p \in (0,1)$ in the limit $C \rightarrow \infty$, one must decrease the ratio R upon increasing C according to the law $R = 1 + \alpha_C/C^{1/2}$, with α_C chosen such that²

$$\alpha \equiv \lim_{C \rightarrow \infty} \alpha_C = 2 \operatorname{erf}^{-1}(1-p). \quad (3)$$

Indeed, since

$$\frac{1}{1+R} = \frac{1}{2} \frac{1}{1 + \alpha_C/(2C^{1/2})} \approx \frac{1}{2} \left(1 - \frac{1}{2} \frac{\alpha_C}{C^{1/2}} \right) \quad (4)$$

as $C \rightarrow \infty$, Eq. (2) above and Theorem 2 of Ref. 2 give

$$\lim_{C \rightarrow \infty} \bar{p}_{\text{acc}}(C) = 1 - \operatorname{erf}\left(\frac{\alpha}{2}\right) = p. \quad (5)$$

In the Introduction of Ref. 2, Predescu, Predescu, and Ciobanu have asserted that Kofke's result does not accurately predict the loss of efficiency in parallel tempering simulations with the increase of the heat capacity or dimensionality. In view of the above-presented analysis, their assertion is incorrect. However, their conclusion has been based on analysis of Eq. (12) of Ref. 1, which, upon setting $B \equiv 1/R \approx 1 - \alpha_C/C^{1/2}$ and letting $C \rightarrow \infty$, indicates that $\bar{p}_{\text{acc}}(C)$ diverges to $+\infty$. This improper behavior is largely due to an unfortunate typographical error³ in the formula as given in Ref. 1. Although correction of this error eliminates the divergence of $\bar{p}_{\text{acc}}(C)$ for $C \rightarrow \infty$, the limiting behavior is still not correct. This is because the formula was developed to describe how \bar{p}_{acc} decreases with increasing dimensionality; accordingly, it is written for a limit taken at constant B . Moreover, it does not attempt to get the behavior correct for B near unity. Therefore it cannot be used to quantify the constant- p loss of efficiency with increasing dimensionality in parallel-tempering simulations. In this regard a more useful asymptotic formula is

$$1 - \bar{p}_{\text{acc}}(C) = \left[\operatorname{erf}\left(\frac{1-B}{1+B} C^{1/2}\right) \right] \times [1 + O(C^{-1})]. \quad (6)$$

This relation is applicable for $C \rightarrow \infty$ limits taken either at constant B or constant \bar{p}_{acc} . In the former case, asymptotic expansion of the error function yields Eq. (12) of Ref. 3, while in the latter case the behavior is consistent with Eq. (5) and the limiting law for $B(C)$ given above.

The author is grateful to Dr. C. Predescu for very helpful correspondence regarding this Comment.

^{a)}Electronic mail: kofke@buffalo.edu

¹D. A. Kofke, J. Chem. Phys. **117**, 6911 (2002).

²C. Predescu, M. Predescu, and C. V. Ciobanu, J. Chem. Phys. **120**, 4119 (2004).

³Erratum to Ref. 1: D. A. Kofke, J. Chem. Phys. **120**, 10852 (2004).