

Foundations of Machine Learning (ECE 5984)

- Neural Networks (1)-

Eunbyung Park

Assistant Professor


School of Electronic and Electrical Engineering

[Eunbyung Park \(silverbottlep.github.io\)](https://silverbottlep.github.io)

Multi-Layer Perceptron



Linear Models as Shallow Neural Networks

- It is a single layer neural network




Linear Models

- Is linear model a good for all?




Nonlinear Models

- nth-degree Polynomial regression



$$f(x) = w_0 + w_1x + w_2x^2 + w_3x^3 + \cdots + w_nx^n$$


Polynomials as Neural Network



$$f(x) = w_0 + w_1x + w_2x^2 + w_3x^3 + \dots + w_nx^n$$

- Feature engineering is hard
- Can we make it non-linear w/o feature engineering?

Feed-Forward Neural Network




$$f(x) = w_1 w_2 x$$

Is it non-linear in x ?

Feed-Forward Neural Network

- Using non-linear activation function




$$f(x) = w_2 a(w_1 x)$$

$$a(x) = \max(0, x) \quad (\text{Rectifier Linear Unit})$$

$$a(x) = \frac{1}{1 + e^{-x}} \quad (\text{Sigmoid})$$


Feed-Forward Neural Network

- AKA, Multi-Layer Perceptron




Feed-Forward Neural Network

- Learning feature representations




Feed-Forward Neural Network

- Biology Inspired



Scaling Laws



Feed-Forward Neural Network

- Regression with two layers MLP

$$D = \{(x^{(1)}, y^{(1)}), \dots, (x^{(N)}, y^{(N)})\}$$


$$x^{(i)} \in \mathbb{R}^d, y^{(i)} \in \mathbb{R}, X \in \mathbb{R}^{N \times d}, Y \in \mathbb{R}^N$$

$$\theta = \{W_1, W_2\}, W_1 \in \mathbb{R}^{h \times d}, W_2 \in \mathbb{R}^{1 \times h}$$

$$f_{\theta}(x) = W_2 \sigma(W_1 x)$$

$$f_{\theta}: \mathbb{R}^d \rightarrow \mathbb{R}$$

$$L(\theta) = \frac{1}{2} \sum_{i=1}^N (y^{(i)} - f_{\theta}(x^{(i)}))^2 = \frac{1}{2} (Y - \sigma(W_1 X^T)^T W_2^T)^T (Y - \sigma(W_1 X^T)^T W_2^T)$$



Feed-Forward Neural Network

- Regression with two layers MLP

$$D = \{(x^{(1)}, y^{(1)}), \dots, (x^{(N)}, y^{(N)})\}$$

$$x^{(i)} \in \mathbb{R}^d, y^{(i)} \in \mathbb{R}, X \in \mathbb{R}^{N \times d}, Y \in \mathbb{R}^N$$

$$\theta = \{W_1, W_2\}, W_1 \in \mathbb{R}^{h \times d}, W_2 \in \mathbb{R}^{1 \times h}$$

$$f_\theta(x) = W_2 \sigma(W_1 x)$$

$$f_\theta: \mathbb{R}^d \rightarrow \mathbb{R}$$

$$L(W_1, W_2) = \frac{1}{2} \sum_{i=1}^N (y^{(i)} - f_\theta(x^{(i)}))^2 = \frac{1}{2} (Y - \sigma(W_1 X^\top)^\top W_2^\top)^\top (Y - \sigma(W_1 X^\top)^\top W_2^\top)$$


1. Can you take the gradients w.r.t θ ?
2. Does it have a closed form solution?
3. Is it a convex function?

Gradient Descent

- We are using *gradient descent* for training deep neural networks

$$W := W - \alpha \left(\frac{\partial L}{\partial W} \right)$$


(descent) (step-size) (gradient)



Learning Representations


Demand Prediction

- Linear regression





Demand Prediction


- Linear regression




Face Recognition



Face Recognition



Car Classification



Exercise

Neural Networks for Boolean Logic Gate Functions

- Which one is linearly separable?
- How many layers required to model 'AND'? And 'XOR'?

Suppose we have binary inputs that only take on values of 0 or 1. Below are truth tables and plots for the Boolean logic gate functions **AND** and **XOR**.

x_1	x_2	AND
0	0	0
0	1	0
1	0	0
1	1	1



x_1	x_2	XOR
0	0	0
0	1	1
1	0	1
1	1	0



Neural Networks for Boolean Logic Gate Functions



Notice that **AND** appears linearly separable (you could draw a line through the figure separating the positive and negative examples) whereas **XOR** does not. Thus, a simple neural network to model the **AND** function might not have a hidden layer whereas a simple neural network to model the **XOR** function might have a hidden layer. Which of the following Boolean logic gate functions are linearly separable?

- NAND
- OR
- NOR
- XNOR

Neural Networks for Boolean Logic Gate Functions

- Which gate function does the neural network make?

$$x_1 \in \{0,1\}, x_2 \in \{0,1\}$$





Sigmoid

Neural Networks for Boolean Logic Gate Functions

- Could you make 'OR'?

$$x_1 \in \{0,1\}, x_2 \in \{0,1\}$$




Sigmoid

Neural Networks for Boolean Logic Gate Functions

- Could you make 'XOR'?

$$x_1 \in \{0,1\}, x_2 \in \{0,1\}$$

x_1	x_2	output
0	0	0
0	1	1
1	0	1
1	1	0




Sigmoid

The Universal Approximator

The Universal Approximation Theorem

- A single hidden layer neural network can approximate any continuous function arbitrarily well, given enough hidden units.
- This holds for many different activation functions, e.g. sigmoid, tanh, ReLU, etc.

The Universal Approximation Theorem



Cybenko Theorem


- Cybenko Approximation by Superposition of Sigmoidal Function

Let $C([0,1]^n)$ denote the set of all continuous function $[0,1]^n \rightarrow \mathbb{R}$, let σ be any sigmoidal activation function then the finite sum of the form $f(x) = \sum_{i=1}^N \alpha_i \sigma(w_i^\top x + b_i)$ is dense in $C([0,1]^n)$


For any $g \in C([0,1]^n)$ and any $\epsilon > 0$, there exists $f: x \rightarrow \sum_{i=1}^N \alpha_i \sigma(w_i^\top x + b_i)$, such that $|f(x) - g(x)| < \epsilon$ for all $x \in [0,1]^n$.

Cybenko Theorem


- Cybenko Approximation by Superposition of Sigmoidal Function



The Universal Approximation Theorem



The Universal Approximation Theorem




$$w = 5, b = 0$$




$$w = 5, b = 3$$




$$w = 10, b = -7$$




$$w = 100, b = -20$$




The Universal Approximation Theorem





$$\begin{aligned}w_1 &= 100, b_1 = -20 \\w_2 &= 100, b_2 = -40 \\w_3 &= 1, w_4 = -1\end{aligned}$$





$$\begin{aligned}w_1 &= 100, b_1 = -60 \\w_2 &= 100, b_2 = -80 \\w_3 &= 2, w_4 = -2\end{aligned}$$





The Universal Approximation Theorem





The Universal Approximation Theorem





The Universal Approximation Theorem





The Universal Approximation Theorem




The Universal Approximator in 2D





The Universal Approximator in 2D



The Universal Approximator in 2D



The Universal Approximator in 2D



The Universal Approximation Theorem


- Single layer might be enough, but it requires ‘enough’ neurons.
- Informally, ‘shallower and wider’ networks require exponentially more hidden units to compute ‘narrower and deeper’ neural networks
 - [Lecture 2 | The Universal Approximation Theorem - YouTube](#)

The Chain Rule

The Chain Rule

- A single variable chain rule

$$f, g, h: \mathbb{R} \rightarrow \mathbb{R}$$



$$f: h \circ g$$

$$f'(x) = h'(g(x))g'(x)$$

$$\frac{dz}{dx} = \frac{dz}{dy} \frac{dy}{dx}$$

$$y = g(x), z = h(y)$$


Simple Example

$$f(x, y, z) = (x + y)z$$

$$q = x + y, \quad f = qz$$

$$\frac{\partial q}{\partial x} = 1, \quad \frac{\partial q}{\partial y} = 1$$

$$\frac{\partial f}{\partial q} = z, \quad \frac{\partial f}{\partial z} = q$$



Simple Example


$$f(x, y, z) = (x + y)z$$

$$q = x + y, \quad f = qz$$

$$\frac{\partial q}{\partial x} = 1, \quad \frac{\partial q}{\partial y} = 1$$

$$\frac{\partial f}{\partial q} = z, \quad \frac{\partial f}{\partial z} = q$$

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$



Simple Example


$$f(x, y, z) = (x + y)z$$

$$q = x + y, \quad f = qz$$

$$\frac{\partial q}{\partial x} = 1, \quad \frac{\partial q}{\partial y} = 1$$

$$\frac{\partial f}{\partial q} = z, \quad \frac{\partial f}{\partial z} = q$$

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$



Simple Example


$$f(x, y, z) = (x + y)z$$

$$q = x + y, \quad f = qz$$

$$\frac{\partial q}{\partial x} = 1, \quad \frac{\partial q}{\partial y} = 1$$

$$\frac{\partial f}{\partial q} = z, \quad \frac{\partial f}{\partial z} = q$$

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$




Simple Example

$$f(x, y, z) = (x + y)z$$

$$q = x + y, \quad f = qz$$

$$\frac{\partial q}{\partial x} = 1, \quad \frac{\partial q}{\partial y} = 1$$
$$\frac{\partial f}{\partial q} = z, \quad \frac{\partial f}{\partial z} = q$$

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y} \frac{\partial q}{\partial y}$$



Simple Example


$$f(x, y, z) = (x + y)z$$

$$q = x + y, \quad f = qz$$

$$\frac{\partial q}{\partial x} = 1, \quad \frac{\partial q}{\partial y} = 1$$

$$\frac{\partial f}{\partial q} = z, \quad \frac{\partial f}{\partial z} = q$$

$$\frac{\partial f}{\partial z} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial z}$$



Sigmoid Example


$$\sigma(x, w) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$

$$f(x) = \frac{1}{x}, \quad g(x) = 1 + x, \quad h(x) = e^{-x}, \quad i(x) = w_0x_0 + w_1x_1 + w_2$$

Sigmoid Example

$$\sigma(x, w) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$

$$f(x) = \frac{1}{x}, \quad g(x) = 1 + x, \quad h(x) = e^{-x}, \quad i(x) = w_0x_0 + w_1x_1 + w_2$$



Backpropagation (Regression w/ MLP)

Gradient

- In vector calculus, the *gradient* of a *scalar-valued* differentiable function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ at the point x

$$\nabla f: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$\nabla f = \frac{\partial f}{\partial x} = \left[\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right]$$

Jacobian

- In vector calculus, the *Jacobian* of a *vector-valued* differentiable function is the matrix of all its first-order partial derivatives.

$$f: \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$J_{ij} = \frac{\partial f_i}{\partial x_j}$$

$$J = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

Matrix Calculus

$$X \in \mathbb{R}^{n \times m}, y \in \mathbb{R}$$

$$f: \mathbb{R}^{n \times m} \rightarrow \mathbb{R}$$

$$y = f(x)$$

$$\frac{\partial y}{\partial X} = \begin{bmatrix} \frac{\partial y}{\partial X_{11}} & \cdots & \frac{\partial y}{\partial X_{1m}} \\ \vdots & \ddots & \vdots \\ \frac{\partial y}{\partial X_{n1}} & \cdots & \frac{\partial y}{\partial X_{nm}} \end{bmatrix} \quad n \times m$$

Matrix Calculus

$$X \in \mathbb{R}^{n \times m}, y \in \mathbb{R}^l$$

$$f: \mathbb{R}^{n \times m} \rightarrow \mathbb{R}^l$$

$$y = f(x)$$

$$\frac{\partial y_1}{\partial X} = \begin{matrix} & & n \times m \\ \begin{bmatrix} \frac{\partial y_1}{\partial X_{11}} & \cdots & \frac{\partial y_1}{\partial X_{1m}} \\ \vdots & \ddots & \vdots \\ \frac{\partial y_1}{\partial X_{n1}} & \cdots & \frac{\partial y_1}{\partial X_{nm}} \end{bmatrix} \end{matrix}$$

$$\frac{\partial y}{\partial X} \quad l \times n \times m \quad (3 \text{ dim tensor})$$

Finite Difference

- Numerical method to compute the gradients based on the definition of gradients

$$\frac{df}{dx} \approx \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

Forward
difference

$$\frac{df}{dx} = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

$$\frac{df}{dx} \approx \frac{f(x) - f(x - \Delta x)}{\Delta x}$$

Backward
difference

$$\frac{df}{dx} \approx \frac{f(x + \Delta x) - f(x - \Delta x)}{2\Delta x}$$

Central
difference

Finite Difference

- Numerical method to compute the gradients based on the definition of gradients

$$\frac{df}{dx} \approx \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

Forward
difference

$$\frac{df}{dx} = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

$$\frac{df}{dx} \approx \frac{f(x) - f(x - \Delta x)}{\Delta x}$$

Backward
difference

What's wrong with this
approach?

$$\frac{df}{dx} \approx \frac{f(x + \Delta x) - f(x - \Delta x)}{2\Delta x}$$

Central
difference


Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial W} ? \quad \frac{\partial L}{\partial v} ?$$




Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial v_i}$$




Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial v_i} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial v_i} = (y - o) \frac{\partial o}{\partial v_i} = (y - o) a_i$$



Two Layers MLP


$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial v_i} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial v_i} = (y - o) \frac{\partial o}{\partial v_i} = (y - o) a_i$$

$$\frac{\partial L}{\partial v} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial v} = (y - o) \frac{\partial o}{\partial v} = (y - o) a$$



Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$


$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial v_i} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial v_i} = (y - o) \frac{\partial o}{\partial v_i} = (y - o) a_i$$

$$\frac{\partial L}{\partial v} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial v} = (y - o) \frac{\partial o}{\partial v} = (y - o) a^\top$$

$1 \times m$ $1 \times m$




Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial W_{ij}} =$$




Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial W_{ij}} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a} \frac{\partial a}{\partial h} \frac{\partial h}{\partial W_{ij}}$$



Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$


$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial W_{ij}} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a} \frac{\partial a}{\partial h} \frac{\partial h}{\partial W_{ij}}$$

$$h_i = \sum_{j=1}^n W_{ij} x_j$$

$$\frac{\partial L}{\partial W_{ij}} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a_i} \frac{\partial a_i}{\partial h_i} \frac{\partial h_i}{\partial W_{ij}}$$



Two Layers MLP


$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$

$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial W_{ij}} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a_i} \frac{\partial a_i}{\partial h_i} \frac{\partial h_i}{\partial W_{ij}} \quad h_i = \sum_{j=1}^n W_{ij} x_j$$

$$\frac{\partial L}{\partial W_{ij}} = (y - o) v_i \left(\sigma(h_i) (1 - \sigma(h_i)) \right) x_j$$




Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$
$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$\frac{\partial L}{\partial W_{ij}} = (y - o) v_i \left(\sigma(h_i)(1 - \sigma(h_i)) \right) x_j$$

$$\frac{\partial L}{\partial W} = (y - o) \begin{bmatrix} \sigma(h_1)(1 - \sigma(h_1)) v_1 x_1 & \cdots & \sigma(h_1)(1 - \sigma(h_1)) v_1 x_n \\ \vdots & \ddots & \vdots \\ \sigma(h_m)(1 - \sigma(h_m)) v_m x_1 & \cdots & \sigma(h_m)(1 - \sigma(h_m)) v_m x_n \end{bmatrix}$$



Backpropagation

(Regression w/ MLP) (2)

Two Layers MLP

$$x \in \mathbb{R}^n, y \in \mathbb{R}, h \in \mathbb{R}^m, a \in [0,1]^m, o \in \mathbb{R}$$

$$W \in \mathbb{R}^{m \times n}, v \in \mathbb{R}^m$$

$$h = Wx \quad a = \sigma(h) \quad o = v^\top a$$


$$L(W, v) = \frac{1}{2} (y - o)^2$$

$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a} \frac{\partial a}{\partial h} \frac{\partial h}{\partial W}$$

$m \times m$ $m \times m \times n$

$$1 \times m \times n$$

$$1 \times 1 \quad 1 \times m$$




Two Layers MLP

$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a} \frac{\partial a}{\partial h} \frac{\partial h}{\partial W}$$

$m \times m$ $m \times m \times n$

$1 \times m \times n$ 1×1 $1 \times m$




$$\frac{\partial L}{\partial o} = (y - o)$$

Two Layers MLP

$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a} \frac{\partial a}{\partial h} \frac{\partial h}{\partial W}$$

$$1 \times m \times n \quad 1 \times 1 \quad 1 \times m$$

$$\frac{\partial o}{\partial a} = v^T$$




Two Layers MLP

$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a} \frac{\partial a}{\partial h} \frac{\partial h}{\partial W}$$

$$1 \times m \times n \quad 1 \times 1 \quad 1 \times m$$

$$\frac{\partial a}{\partial h} = \begin{bmatrix} \sigma(h_1)(1 - \sigma(h_1)) & 0 & 0 \\ 0 & \sigma(h_2)(1 - \sigma(h_2)) & 0 \\ 0 & 0 & \sigma(h_m)(1 - \sigma(h_m)) \end{bmatrix}$$




Algebraic Magic

$$\frac{\partial L}{\partial W} = \overset{m \times m}{\frac{\partial L}{\partial o}} \overset{m \times m \times n}{\frac{\partial o}{\partial a}} \frac{\partial a}{\partial h} \frac{\partial h}{\partial W}$$

$$\overset{1 \times m \times n}{1 \times 1} \overset{1 \times m}{1 \times m}$$

$$\begin{aligned} \frac{\partial h_1}{\partial W} &= \begin{bmatrix} \frac{\partial h_1}{\partial W_{11}} & \cdots & \frac{\partial h_1}{\partial W_{1n}} \\ \vdots & \ddots & \vdots \\ \frac{\partial h_1}{\partial W_{m1}} & \cdots & \frac{\partial h_1}{\partial W_{mn}} \end{bmatrix} = \begin{bmatrix} \frac{\partial h_1}{\partial W_{11}} & \cdots & \frac{\partial h_1}{\partial W_{1n}} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ \frac{\partial h_2}{\partial W} &= \begin{bmatrix} \frac{\partial h_2}{\partial W_{11}} & \cdots & \frac{\partial h_2}{\partial W_{1n}} \\ \vdots & \ddots & \vdots \\ \frac{\partial h_2}{\partial W_{m1}} & \cdots & \frac{\partial h_2}{\partial W_{mn}} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ \frac{\partial h_2}{\partial W_{21}} & \cdots & \frac{\partial h_2}{\partial W_{2n}} \\ 0 & 0 & 0 \end{bmatrix} \end{aligned}$$



Tensor Product

- N-mode product

$$X \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N} \qquad Y \in \mathbb{R}^{J \times I_n}$$



$$(X \times_n Y)_{i_1, \dots, i_{n-1}, \textcolor{red}{j}, i_{n+1}, \dots, i_N} = \sum_{\textcolor{green}{i}_n=1}^{\textcolor{green}{I}_n} X_{i_1, \dots, i_{n-1}, \textcolor{green}{i}_n, i_{n+1}, \dots, i_N} Y_{\textcolor{red}{j}, \textcolor{green}{i}_n}$$

$$(X \times_n Y) \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times J \times \cdots \times I_N}$$

Tensor Product

- N-mode product
 - Matricization -> matrix multiplication


$$\overset{1 \times m \times n}{\frac{\partial L}{\partial W}} = \underset{1 \times m}{\frac{\partial L}{\partial h}} \overset{m \times m \times n}{\frac{\partial h}{\partial W}}$$



Tensor Product

$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial o} \frac{\partial o}{\partial a} \frac{\partial a}{\partial h} \frac{\partial h}{\partial W}$$

$$1 \times m \times n \quad 1 \times 1 \quad 1 \times m$$



$$\text{reshape} \left(\frac{\partial h}{\partial W} \right) = \begin{bmatrix} \frac{\partial h_1}{\partial W_{11}} & \dots & \frac{\partial h_1}{\partial W_{1n}} & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & \frac{\partial h}{\partial W_{21}} & \dots & \frac{\partial h_2}{\partial W_{2n}} & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\partial h_3}{\partial W_{31}} & \dots & \frac{\partial h_3}{\partial W_{3n}} & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \end{bmatrix}$$

$$m \times mn$$

Tensor Product


$m \times m \times n$

$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial h} \frac{\partial h}{\partial W}$$

$1 \times m$

$$\frac{\partial L}{\partial h} \text{reshape} \left(\frac{\partial h}{\partial W} \right) =$$

$$\begin{bmatrix} \frac{\partial L}{\partial h_1} & \frac{\partial L}{\partial h_2} & \dots & \frac{\partial L}{\partial h_m} \end{bmatrix} \begin{bmatrix} \frac{\partial h_1}{\partial W_{11}} & \dots & \frac{\partial h_1}{\partial W_{1n}} & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & \frac{\partial h_2}{\partial W_{21}} & \dots & \frac{\partial h_2}{\partial W_{2n}} & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\partial h_3}{\partial W_{31}} & \dots & \frac{\partial h_3}{\partial W_{3n}} & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \end{bmatrix}$$



Tensor Product

$$m \times m \times n$$


$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial h} \frac{\partial h}{\partial W}$$

$$1 \times m$$

$$\frac{\partial L}{\partial h} \text{reshape} \left(\frac{\partial h}{\partial W} \right) =$$

$$\begin{bmatrix} \frac{\partial L}{\partial h_1} & \frac{\partial L}{\partial h_2} & \dots & \frac{\partial L}{\partial h_m} \end{bmatrix} \begin{bmatrix} \frac{\partial h_1}{\partial W_{11}} & \dots & \frac{\partial h_1}{\partial W_{1n}} & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & \frac{\partial h_2}{\partial W_{21}} & \dots & \frac{\partial h_2}{\partial W_{2n}} & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\partial h_3}{\partial W_{31}} & \dots & \frac{\partial h_3}{\partial W_{3n}} & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \end{bmatrix}$$

$$= \begin{bmatrix} \frac{\partial L}{\partial h_1} \frac{\partial h}{\partial W_{11}} & \dots & \frac{\partial L}{\partial h_1} \frac{\partial h_1}{\partial W_{1n}} & \frac{\partial L}{\partial h_2} \frac{\partial h_2}{\partial W_{21}} & \dots & \frac{\partial L}{\partial h_2} \frac{\partial h_2}{\partial W_{2n}} & \dots \end{bmatrix}$$



Tensor Product

$m \times m \times n$


$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial h} \frac{\partial h}{\partial W}$$

$$\frac{\partial L}{\partial W} = \frac{\partial L}{\partial h} \frac{\partial h}{\partial W} \stackrel{1 \times m}{=} \text{reshape} \left(\frac{\partial L}{\partial h} \stackrel{1 \times m}{\text{reshape}} \left(\frac{\partial h}{\partial W} \right) \stackrel{m \times mn}{\text{reshape}} \right)$$

$$= \text{reshape} \left(\left[\frac{\partial L}{\partial h_1} \frac{\partial h}{\partial W_{11}} \quad \dots \quad \frac{\partial L}{\partial h_1} \frac{\partial h_1}{\partial W_{1n}} \quad \frac{\partial L}{\partial h_2} \frac{\partial h_2}{\partial W_{21}} \quad \dots \quad \frac{\partial L}{\partial h_2} \frac{\partial h_2}{\partial W_{2n}} \quad \dots \right] \right) \stackrel{1 \times mn}{\text{reshape}}$$

$$\stackrel{m \times n}{=} \begin{bmatrix} \frac{\partial L}{\partial h_1} \frac{\partial h_1}{\partial W_{11}} & \dots & \frac{\partial L}{\partial h_1} \frac{\partial h_1}{\partial W_{1n}} \\ \vdots & \ddots & \vdots \\ \frac{\partial L}{\partial h_m} \frac{\partial h_m}{\partial W_{m1}} & \dots & \frac{\partial L}{\partial h_m} \frac{\partial h_m}{\partial W_{mn}} \end{bmatrix} = \begin{bmatrix} \frac{\partial L}{\partial h_1} x_1 & \dots & \frac{\partial L}{\partial h_1} x_n \\ \vdots & \ddots & \vdots \\ \frac{\partial L}{\partial h_m} x_1 & \dots & \frac{\partial L}{\partial h_m} x_n \end{bmatrix} = \left(\frac{\partial L}{\partial h} \right)^\top x^\top$$

$(y - o)(\sigma(h)(1 - \sigma(h)) \odot v)$



Automatic Differentiation

Automatic Differentiation (AD)

- A procedure for automatic evaluation of derivatives of arbitrary algebraic functions
- Backpropagation == reverse-mode AD

Reverse-Mode AD (a.k.a Backpropagation)

$$f: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$c = g(b)$$

$$h: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$d = h(c)$$

$$e = i(d)$$

$$i: \mathbb{R}^{n_4} \rightarrow \mathbb{R}$$

$$\frac{\partial e}{\partial a}?$$

Loss function:
scalar function

Reverse-Mode AD (a.k.a Backpropagation)

$$\frac{\partial e}{\partial d} = \overset{1 \times 1}{\frac{\partial e}{\partial e}} \overset{1 \times n}{\frac{\partial e}{\partial d}} = \overset{1 \times 1}{1} \frac{\partial e}{\partial d}$$

$$f: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$c = g(b)$$

$$d = h(c)$$

$$h: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$e = i(d)$$

$$i: \mathbb{R}^{n_4} \rightarrow \mathbb{R}$$

Loss function:
scalar function

Reverse-Mode AD (a.k.a Backpropagation)

$$f: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$c = g(b)$$

$$d = h(c)$$

$$h: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$e = i(d)$$

$$i: \mathbb{R}^{n_4} \rightarrow \mathbb{R}$$

Loss function:
scalar function

$$\frac{\partial e}{\partial d} = \frac{\overset{1 \times 1}{\partial e}}{\partial e} \overset{1 \times n_4}{\frac{\partial e}{\partial d}} = 1 \frac{\partial e}{\partial d}$$

$$\frac{\partial e}{\partial c} = \frac{\overset{1 \times n_4}{\partial e}}{\partial e} \overset{n_4 \times n_3}{\frac{\partial e}{\partial d} \frac{\partial d}{\partial c}} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c}$$

Reverse-Mode AD (a.k.a Backpropagation)

$$f: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$c = g(b)$$

$$d = h(c)$$

$$h: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$e = i(d)$$

$$i: \mathbb{R}^{n_4} \rightarrow \mathbb{R}$$

Loss function:
scalar function

$$\frac{\partial e}{\partial d} = \frac{\overset{1 \times 1}{\partial e}}{\partial e} \overset{1 \times n_4}{\frac{\partial e}{\partial d}} = 1 \frac{\partial e}{\partial d}$$

$$\frac{\partial e}{\partial c} = \frac{\overset{1 \times n_4}{\partial e}}{\partial e} \overset{n_4 \times n_3}{\frac{\partial e}{\partial d} \frac{\partial d}{\partial c}} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c}$$

$$\frac{\partial e}{\partial b} = \frac{\overset{1 \times n_3}{\partial e}}{\partial e} \overset{n_3 \times n_2}{\frac{\partial e}{\partial d} \frac{\partial d}{\partial c} \frac{\partial c}{\partial b}} = \frac{\partial e}{\partial c} \frac{\partial c}{\partial b}$$

Reverse-Mode AD (a.k.a Backpropagation)

$$f: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$c = g(b)$$

$$h: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$d = h(c)$$

$$e = i(d)$$

$$i: \mathbb{R}^{n_4} \rightarrow \mathbb{R}$$

Loss function:
scalar function

$$\frac{\partial e}{\partial d} = \frac{\overset{1 \times 1}{\partial e}}{\partial e} \overset{1 \times n_4}{\frac{\partial e}{\partial d}} = 1 \frac{\partial e}{\partial d}$$

Vector-Jacobian
Product (VJP)

$$\frac{\partial e}{\partial c} = \frac{\overset{1 \times n_4}{\partial e}}{\partial e} \overset{n_4 \times n_3}{\frac{\partial e}{\partial d} \frac{\partial d}{\partial c}} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c}$$

$$\frac{\partial e}{\partial b} = \frac{\overset{1 \times n_3}{\partial e}}{\partial e} \overset{n_3 \times n_2}{\frac{\partial e}{\partial d} \frac{\partial d}{\partial c} \frac{\partial c}{\partial b}} = \frac{\partial e}{\partial c} \frac{\partial c}{\partial b}$$

$$\frac{\partial e}{\partial a} = \frac{\overset{1 \times n_2}{\partial e}}{\partial e} \overset{n_2 \times n_1}{\frac{\partial e}{\partial d} \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} \frac{\partial b}{\partial a}} = \frac{\partial e}{\partial b} \frac{\partial b}{\partial a}$$

Reverse-Mode AD (a.k.a Backpropagation)

$$f: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$c = g(b)$$

$$h: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$d = h(c)$$

$$e = i(d)$$

$$i: \mathbb{R}^{n_4} \rightarrow \mathbb{R}^3$$

What if i is
vector-valued function?

$$\frac{\partial e}{\partial d} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial d} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \frac{\partial e}{\partial d}$$

3 X
Computation

$$\frac{\partial e}{\partial c} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c}$$

$$\frac{\partial e}{\partial b} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} = \frac{\partial e}{\partial c} \frac{\partial c}{\partial b}$$

$$\frac{\partial e}{\partial a} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} = \frac{\partial e}{\partial b} \frac{\partial b}{\partial a}$$

Forward-Mode AD

single variable

$$f: \mathbb{R} \rightarrow \mathbb{R}^{n_1}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$c = g(b)$$

$$d = h(c)$$

$$h: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$e = L(d)$$

$$L: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$\frac{\partial b}{\partial a} = \frac{\overset{n_1 \times 1}{\partial b}}{\overset{1 \times 1}{\partial a}} \overset{1 \times 1}{\frac{\partial a}{\partial a}} = \frac{\partial b}{\partial a} \mathbf{1}$$

Forward-Mode AD

single variable

$$f: \mathbb{R} \rightarrow \mathbb{R}^{n_1}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$c = g(b)$$

$$h: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$d = h(c)$$

$$e = L(d)$$

$$L: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$\frac{\partial b}{\partial a} = \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial b}{\partial a} \mathbf{1}$$

$n_1 \times 1 \quad 1 \times 1$

$$\frac{\partial c}{\partial a} = \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial c}{\partial b} \frac{\partial b}{\partial a}$$

$n_2 \times n_1 \quad n_1 \times 1$

Forward-Mode AD

single variable

$$f: \mathbb{R} \rightarrow \mathbb{R}^{n_1}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$c = g(b)$$

$$h: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$d = h(c)$$

$$e = L(d)$$

$$L: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$\frac{\partial b}{\partial a} = \frac{\overset{n_1 \times 1}{\partial b}}{\overset{1 \times 1}{\partial a}} \overset{1 \times 1}{\frac{\partial a}{\partial a}} = \overset{n_1 \times 1}{\frac{\partial b}{\partial a}} \overset{1 \times 1}{1}$$

$$\frac{\partial c}{\partial a} = \frac{\overset{n_2 \times n_1}{\partial c}}{\overset{n_1 \times 1}{\partial b}} \overset{n_1 \times 1}{\frac{\partial b}{\partial a}} \overset{n_1 \times 1}{\frac{\partial a}{\partial a}} = \overset{n_2 \times n_1}{\frac{\partial c}{\partial b}} \overset{n_1 \times 1}{\frac{\partial b}{\partial a}}$$

$$\frac{\partial d}{\partial a} = \frac{\overset{n_3 \times n_2}{\partial d}}{\overset{n_2 \times 1}{\partial c}} \overset{n_2 \times 1}{\frac{\partial c}{\partial b}} \overset{n_2 \times 1}{\frac{\partial b}{\partial a}} \overset{n_2 \times 1}{\frac{\partial a}{\partial a}} = \overset{n_3 \times n_2}{\frac{\partial d}{\partial c}} \overset{n_2 \times 1}{\frac{\partial c}{\partial b}}$$

Forward-Mode AD

single variable

$$f: \mathbb{R} \rightarrow \mathbb{R}^{n_1}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$c = g(b)$$

$$h: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$d = h(c)$$

$$L: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

$$e = L(d)$$

$$\frac{\partial b}{\partial a} = \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial b}{\partial a} \overset{n_1 \times 1}{1} \overset{1 \times 1}{1}$$

Jacobian-Vector
Product (JVP)

$$\frac{\partial c}{\partial a} = \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \overset{n_2 \times n_1}{1} \overset{n_1 \times 1}{1}$$

$$\frac{\partial d}{\partial a} = \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \overset{n_3 \times n_2}{1} \overset{n_2 \times 1}{1}$$

$$\frac{\partial e}{\partial a} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \overset{n_4 \times n_3}{1} \overset{n_3 \times 1}{1}$$

Forward-Mode AD

$$f: \mathbb{R}^3 \rightarrow \mathbb{R}^{n_1}$$

$$b = f(a)$$

$$g: \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$$

$$c = g(b)$$

$$h: \mathbb{R}^{n_2} \rightarrow \mathbb{R}^{n_3}$$

$$d = h(c)$$

$$e = L(d)$$

$$L: \mathbb{R}^{n_3} \rightarrow \mathbb{R}^{n_4}$$

What if input is
Multi-variables?

$$\frac{\partial b}{\partial a} = \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial b}{\partial a} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

3 X
Computation

$$\frac{\partial c}{\partial a} = \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial c}{\partial b} \frac{\partial b}{\partial a}$$

$$\frac{\partial d}{\partial a} = \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial d}{\partial c} \frac{\partial c}{\partial a}$$


$$\frac{\partial e}{\partial a} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial c} \frac{\partial c}{\partial b} \frac{\partial b}{\partial a} \frac{\partial a}{\partial a} = \frac{\partial e}{\partial d} \frac{\partial d}{\partial a}$$

Automatic Differentiation (AD)




- For low dimensional outputs and high dimensional inputs
 - Objective function w/ deep neural networks
 - reverse-mode AD
- For high dimensional outputs and low dimensional inputs
 - Forward-mode AD

Computational Graph

```
t0 = x - m
t1 = t0 / s
t2 = np.log(s)
t3 = t1**2
t4 = t2 + np.log(2 * np.pi) / 2
t5 = t3 / 2
c = t4 + t5
```



Automatic Differentiation



References

- [mattjj/autodidact: A pedagogical implementation of Autograd \(github.com\)](#)
- [\[1502.05767\] Automatic differentiation in machine learning: a survey \(arxiv.org\)](#)
- [CSC321 Lecture 10: Automatic Differentiation \(toronto.edu\)](#)