

Quantium Virtual Internship - Retail Strategy and Analytics - Task

2

Daniel Ma

25/03/2022

```
#### Example code to install packages
#install.packages("data.table")
#### Load required libraries
library(data.table)
library(ggplot2)
library(ggmosaic)
library(readr)
library(readxl)
library(dplyr)
library(stringr)
library(tibbltime)
filePath <- "C:/Users/Signature/OneDrive/Documents/Careers/Virtual Experiences/From DXM YOGA/Quantium/"
data <- fread(paste0(filePath,"QVI_data.csv"))

#### Set themes for plots
theme_set(theme_bw())
theme_update(plot.title = element_text(hjust = 0.5))
```

Select control stores

The client has selected store numbers 77, 86 and 88 as trial stores and want control stores to be established stores that are operational for the entire observation period. We would want to match trial stores to control stores that are similar to the trial store prior to the trial period of Feb 2019 in terms of : - Monthly overall sales revenue - Monthly number of customers - Monthly number of transactions per customer Let's first create the metrics of interest and filter to stores that are present throughout the pre-trial period.

```
#### Calculate these measures over time for each store
data[, YEARMONTH := format(as.Date(Date), "%Y%m")]
#### Next, we define the measure calculations to use during the analysis.
measureOverTime <- data[, .(totSales = sum(TOT_SALES),
                           nCustomers = uniqueN(LYLT_CARD_NBR),
                           nTxnPerCust = uniqueN(TXN_ID)/uniqueN(LYLT_CARD_NBR),
                           nChipsPerTxn = sum(PROD_QTY)/uniqueN(TXN_ID),
                           avgPricePerUnit = sum(TOT_SALES)/sum(PROD_QTY))
                        , by = c("STORE_NBR", "YEARMONTH")][order(STORE_NBR, YEARMONTH)]
#### Filter to the pre-trial period and stores with full observation periods
storesWithFullObs <- unique(measureOverTime[, .N, STORE_NBR][N == 12, STORE_NBR])
preTrialMeasures <- measureOverTime[YEARMONTH < 201902 & STORE_NBR %in%
storesWithFullObs, ]
```

Now we need to work out a way of ranking how similar each potential control store is to the trial store. We can calculate how correlated the performance of each store is to the trial store. Let's write a function for this so that we don't have to calculate this for each trial store and control store pair.

```
#### Let's define inputTable as a metric table with potential comparison stores,
#metricCol as the store metric used to calculate correlation on, and storeComparison
#as the store number of the trial store.
calculateCorrelation <- function(inputTable, metricCol, storeComparison) {
  calcCorrTable = data.table(Store1 = numeric(), Store2 = numeric(), corr_measure =
  numeric())
  storeNumbers <- unique(inputTable[, STORE_NBR])
  for (i in storeNumbers) {
    calculatedMeasure = data.table("Store1" = storeComparison,
                                   "Store2" = i,
                                   "corr_measure" = cor(inputTable[STORE_NBR == storeComparison,
                                                             eval(metricCol)],
                                                             inputTable[STORE_NBR == i, eval(metricCol)]))
  }
  calcCorrTable <- rbind(calcCorrTable, calculatedMeasure)
}
return(calcCorrTable)
}
```

Apart from correlation, we can also calculate a standardised metric based on the absolute difference between the trial store's performance and each control store's performance. Let's write a function for this.

```
calculateMagnitudeDistance <- function(inputTable, metricCol, storeComparison) {
  calcDistTable = data.table(Store1 = numeric(), Store2 = numeric(), YEARMONTH =
  numeric(), measure = numeric())
  storeNumbers <- unique(inputTable[, STORE_NBR])
  for (i in storeNumbers) {
    calculatedMeasure = data.table("Store1" = storeComparison
    , "Store2" = i
    , "YEARMONTH" = inputTable[STORE_NBR == storeComparison, YEARMONTH]
    , "measure" = abs(inputTable[STORE_NBR == storeComparison, eval(metricCol)]
    - inputTable[STORE_NBR == i, eval(metricCol)])
  }
  calcDistTable <- rbind(calcDistTable, calculatedMeasure)
}

#### Standardise the magnitude distance so that the measure ranges from 0 to 1
minMaxDist <- calcDistTable[, .(minDist = min(measure), maxDist = max(measure)),
by = c("Store1", "YEARMONTH")]
distTable <- merge(calcDistTable, minMaxDist, by = c("Store1", "YEARMONTH"))
distTable[, magnitudeMeasure := 1 - (measure - minDist)/(maxDist - minDist)]
finalDistTable <- distTable[, .(mag_measure = mean(magnitudeMeasure)), by =
.(Store1, Store2)]
return(finalDistTable)
}
```

Now let's use the functions to find the control stores! We'll select control stores based on how similar monthly total sales in dollar amounts and monthly number of customers are to the trial stores. So we will need to use our functions to get four scores, two for each of total sales and total customers.

```

trial_store <- 77
corr_nSales <- calculateCorrelation(preTrialMeasures, quote(totSales), trial_store)
corr_nCustomers <- calculateCorrelation(preTrialMeasures, quote(nCustomers), trial_store)
#### Then, use the functions for calculating magnitude.
magnitude_nSales <- calculateMagnitudeDistance(preTrialMeasures, quote(totSales),
trial_store)
magnitude_nCustomers <- calculateMagnitudeDistance(preTrialMeasures,
quote(nCustomers), trial_store)

```

We'll need to combine the all the scores calculated using our function to create a composite score to rank on. Let's take a simple average of the correlation and magnitude scores for each driver. Note that if we consider it more important for the trend of the drivers to be similar, we can increase the weight of the correlation score (a simple average gives a weight of 0.5 to the corr_weight) or if we consider the absolute size of the drivers to be more important, we can lower the weight of the correlation score.

```

corr_weight <- 0.5
score_nSales <- merge(corr_nSales, magnitude_nSales, by = c("Store1", "Store2")
) [, scoreNSales := 0.5*(corr_measure + mag_measure)]
score_nCustomers <- merge(corr_nCustomers, magnitude_nCustomers, by = c("Store1", "Store2")
) [, scoreNCust := 0.5*(corr_measure + mag_measure)]

```

Now we have a score for each of total number of sales and number of customers. Let's combine the two via a simple average.

```

score_Control <- merge(score_nSales, score_nCustomers, by = c("Store1", "Store2"))
score_Control[, finalControlScore := scoreNSales * 0.5 + scoreNCust * 0.5]

```

The store with the highest score is then selected as the control store since it is most similar to the trial store.

```

#### Select control stores based on the highest matching store (closest to 1 but
#### not the store itself, i.e. the second ranked highest store)
#### Select the most appropriate control store for trial store 77 by finding the
# store with the highest final score.
control_store <- score_Control[Store1 == trial_store,][order(-finalControlScore)][2, Store2]
control_store

```

```
## [1] 233
```

Now that we have found a control store, let's check visually if the drivers are indeed similar in the period before the trial. We'll look at total sales first.

```


#### Visual checks on trends based on the drivers
measureOverTimeSales <- measureOverTime
pastSales <- measureOverTimeSales[, Store_type := ifelse(STORE_NBR == trial_store,
"Trial",
ifelse(STORE_NBR == control_store,
"Control", "Other stores"))
][, totSales := mean(totSales), by = c("YEARMONTH",
"Store_type")]
[, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %/% 100,
as.numeric(YEARMONTH) %% 100, 1, sep = "-"),

```

```

                                "%Y-%m-%d")
][as.numeric(YEARMONTH) < 201903 , ]
ggplot(pastSales, aes(TransactionMonth, totSales, color = Store_type)) +
geom_line() +
labs(x = "Month of operation", y = "Total sales", title = "Total sales by month")

```



Next, number of customers.

```

measureOverTimeCusts <- measureOverTime
pastCustomers <- measureOverTimeCusts[, Store_type := ifelse(STORE_NBR == trial_store,
"Trial",
ifelse(STORE_NBR == control_store,
"Control", "Other stores"))][, numCustomers := mean(nCustomers), by = c("YEARMONTH", "Store_type")
][, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %/% 100, as.numeric(YEARMONTH) %/% 10, as.numeric(YEARMONTH) % 10), format = "%Y-%m-%d")
][as.numeric(YEARMONTH) < 201903,]
ggplot(pastCustomers, aes(TransactionMonth, numCustomers, color = Store_type)) +
geom_line() +
labs(x = "Month", y = "Total number of Customers", title = "Total Customers by Month")

```



Assessment of trial The trial period goes from the start of February 2019 to April 2019. We now want to see if there has been an uplift in overall chip sales. We'll start with scaling the control store's sales to a level similar to control for any differences between the two stores outside of the trial period.

```
#### Scale pre-trial control sales to match pre-trial trial store sales
scalingFactorForControlSales <- preTrialMeasures[STORE_NBR == trial_store &
YEARMONTH < 201902, sum(totSales)]/preTrialMeasures[STORE_NBR == control_store &
YEARMONTH < 201902, sum(totSales)]
#### Apply the scaling factor
measureOverTimeSales <- measureOverTime
scaledControlSales <- measureOverTimeSales[STORE_NBR == control_store, ][ ,
controlSales := totSales * scalingFactorForControlSales]
```

Now that we have comparable sales figures for the control store, we can calculate the percentage difference between the scaled control sales and the trial store's sales during the trial period.

```
percentageDiff <- merge(scaledControlSales[, c("YEARMONTH", "controlSales")],
  measureOverTime[STORE_NBR == trial_store, c("totSales", "YEARMONTH")],
  by = "YEARMONTH")[, percentageDiff := abs(controlSales-totSales)/controlSales]
```

Let's see if the difference is significant!

```
#### As our null hypothesis is that the trial period is the same as the pre-trial
# period, let's take the standard deviation based on the scaled percentage difference
# in the pre-trial period
stdDev <- sd(percentageDiff[YEARMONTH < 201902 , percentageDiff])
```


```
#### Note that there are 8 months in the pre-trial period
#### hence 8 - 1 = 7 degrees of freedom
degreesOfFreedom <- 7
#### We will test with a null hypothesis of there being 0 difference between trial
# and control stores.

percentageDiff[, tValue := percentageDiff/stdDev
][, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %% 100, as.numeric(YEARMONTH) %% 100, 1,
                                     sep = "-"), "%Y-%m-%d")
][as.numeric(YEARMONTH) < 201905 & as.numeric(YEARMONTH) > 201901, .(TransactionMonth, tValue)]

##      TransactionMonth      tValue
## 1:      2019-02-01  1.223912
## 2:      2019-03-01  5.633494
## 3:      2019-04-01 11.336505
```

We can observe that the t-value is much larger than the 95th percentile value of the t-distribution for March and April - i.e. the increase in sales in the trial store in March and April is statistically greater than in the control store. Let's create a more visual version of this by plotting the sales of the control store, the sales of the trial stores and the 95th percentile value of sales of the control store.

```
measureOverTimeSales <- measureOverTime
#### Trial and control store total sales
pastSales <- measureOverTimeSales[, Store_type := ifelse(STORE_NBR == trial_store,
                                                         "Trial",
                                                         ifelse(STORE_NBR == control_store, "Control", "Other stores"))
][, totSales := mean(totSales), by = c("YEARMONTH", "Store_type")
][, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %% 100,
                                     as.numeric(YEARMONTH) %% 100, 1,
                                     sep = "-"), "%Y-%m-%d")
][Store_type %in% c("Trial", "Control"), ]
#### Control store 95th percentile
pastSales_Controls95 <- pastSales[Store_type == "Control",
                                 ][, totSales := totSales * (1 + stdDev * 2)
                                 ][, Store_type := "Control 95th % confidence interval"]
#### Control store 5th percentile
pastSales_Controls5 <- pastSales[Store_type == "Control",
                                 ][, totSales := totSales * (1 - stdDev * 2)
                                 ][, Store_type := "Control 5th % confidence interval"]
trialAssessment <- rbind(pastSales, pastSales_Controls95, pastSales_Controls5)
#### Plotting these in one nice graph
ggplot(trialAssessment, aes(TransactionMonth, totSales, color = Store_type)) +
  geom_rect(data = trialAssessment[ YEARMONTH < 201905 & YEARMONTH > 201901 ,],
  aes(xmin = min(TransactionMonth), xmax = max(TransactionMonth), ymin = 0 , ymax =
Inf, color = NULL), show.legend = FALSE) +
  geom_line() +
  labs(x = "Month of operation", y = "Total sales", title = "Total sales by month")
```



The results show that the trial in store 77 is significantly different to its control store in the trial period as the trial store performance lies outside the 5% to 95% confidence interval of the control store in two of the three trial months. Let's have a look at assessing this for number of customers as well.

```
#### Scale pre-trial control customers to match pre-trial trial store customers
scalingFactorForControlCust <- preTrialMeasures[STORE_NBR == trial_store &
YEARMONTH < 201902, sum(nCustomers)] / preTrialMeasures[STORE_NBR ==
control_store & YEARMONTH < 201902, sum(nCustomers)]
measureOverTimeCusts <- measureOverTime
scaledControlCustomers <- measureOverTimeCusts[STORE_NBR == control_store,
][, controlCustomers := nCustomers * scalingFactorForControlCust
][, Store_type := ifelse(STORE_NBR == trial_store, "Trial",
ifelse(STORE_NBR == control_store, "Control", "Other stores"))]
percentageDiff <- merge(scaledControlCustomers[, c("YEARMONTH", "controlCustomers")],
measureOverTimeCusts[STORE_NBR == trial_store, c("nCustomers", "YEARMONTH")],
by = "YEARMONTH"
)[, percentageDiff := abs(controlCustomers - nCustomers) / controlCustomers]
```

Let's again see if the difference is significant visually!


```
#### As our null hypothesis is that the trial period is the same as the pre-trial
# period, let's take the standard deviation based on the scaled percentage difference
# in the pre-trial period
stdDev <- sd(percentageDiff[as.numeric(YEARMONTH) < 201902, percentageDiff])
degreesOfFreedom <- 7
#### Trial and control store number of customers
```

```

pastCustomers <- measureOverTimeCusts[, nCusts := mean(nCustomers), by =
c("YEARMONTH", "Store_type")
][Store_type %in% c("Trial", "Control"), ]
#### Control store 95th percentile
pastCustomers_Controls95 <- pastCustomers[Store_type == "Control",
][, nCusts := nCusts * (1 + stdDev * 2)
][, Store_type := "Control 95th % confidence
interval"]
#### Control store 5th percentile
pastCustomers_Controls5 <- pastCustomers[Store_type == "Control",
][, nCusts := nCusts * (1 - stdDev * 2)
][, Store_type := "Control 5th % confidence
interval"]
trialAssessment <- rbind(pastCustomers, pastCustomers_Controls95,
pastCustomers_Controls5)

ggplot(trialAssessment, aes(TransactionMonth, nCusts, color = Store_type)) +
geom_rect(data = trialAssessment[ YEARMONTH < 201905 & YEARMONTH > 201901 ,]
, aes(xmin = min(TransactionMonth), xmax = max(TransactionMonth)
, ymin = 0, ymax = Inf, color = NULL),
show.legend = FALSE) +
geom_line(aes(linetype = Store_type)) +
labs(x = "Month of operation", y = "Total number of customers",
title = "Total number of customers by month")

```



Let's repeat finding the control store and assessing the impact of the trial for each of the other two trial

stores. ## Trial store 86

```
#### Calculate the metrics below as we did for the first trial store.
measureOverTime <- data[, .(totSales = sum(TOT_SALES),
                             nCustomers = uniqueN(LYLT_CARD_NBR),
                             nTxnPerCust = uniqueN(TXN_ID)/uniqueN(LYLT_CARD_NBR),
                             nChipsPerTxn = sum(PROD_QTY)/uniqueN(TXN_ID),
                             avgPricePerUnit = sum(TOT_SALES)/sum(PROD_QTY)
                           )
, by = c("STORE_NBR", "YEARMONTH")][order(STORE_NBR, YEARMONTH)]


trial_store <- 86
corr_nSales <- calculateCorrelation(preTrialMeasures, quote(totSales), trial_store)
corr_nCustomers <- calculateCorrelation(preTrialMeasures, quote(nCustomers), trial_store)
magnitude_nSales <- calculateMagnitudeDistance(preTrialMeasures, quote(totSales),
trial_store)
magnitude_nCustomers <- calculateMagnitudeDistance(preTrialMeasures,
quote(nCustomers), trial_store)
#### Now, create a combined score composed of correlation and magnitude
corr_weight <- 0.5
score_nSales <- merge(corr_nSales, magnitude_nSales, by = c("Store1", "Store2")
)[, scoreNSales := 0.5*(corr_measure + mag_measure)]
score_nCustomers <- merge(corr_nCustomers, magnitude_nCustomers, by = c("Store1", "Store2")
)[, scoreNCust := 0.5*(corr_measure + mag_measure)]
#### Finally, combine scores across the drivers using a simple average.
score_Control <- merge(score_nSales, score_nCustomers, by = c("Store1", "Store2"))
score_Control[, finalControlScore := 0.5*(scoreNSales+scoreNCust)]
#### Select control stores based on the highest matching store
#### (closest to 1 but not the store itself, i.e. the second ranked highest store)
#### Select control store for trial store 86
control_store <- score_Control[Store1 == trial_store,
][order(-finalControlScore)][2, Store2]
control_store
```

[1] 155

Looks like store 155 will be a control store for trial store 86. Again, let's check visually if the drivers are indeed similar in the period before the trial. We'll look at total sales first.


```
measureOverTimeSales <- measureOverTime
pastSales <- measureOverTimeSales[, Store_type := ifelse(STORE_NBR == trial_store,
                                                         "Trial",
                                                         ifelse(STORE_NBR == control_store, "Control", "Other stores"))
][, totSales := mean(totSales), by = c("YEARMONTH", "Store_type")]
[, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %/% 100,
                                     as.numeric(YEARMONTH) %% 100, 1,
                                     sep = "-"), "%Y-%m-%d")][YEARMONTH < 201903]

ggplot(pastSales, aes(TransactionMonth, totSales, color = Store_type)) +
  geom_line(aes(linetype = Store_type)) +
  labs(x = "Month", y = "Total sales", title = "Total sales by month")
```



Great, sales are trending in a similar way. Next, number of customers.

```
#### Conduct visual checks on trends based on the drivers
measureOverTimeCusts <- measureOverTime
pastCustomers <- measureOverTimeCusts[, Store_type := ifelse(STORE_NBR == trial_store,
"Trial", ifelse(STORE_NBR == control_store,
"Control", "Other stores"))
][, numCustomers := mean(nCustomers), by = c("YEARMONTH", "Store_type")]
][, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %/% 100,
as.numeric(YEARMONTH) %/% 100, 1, sep = "-"),
"%Y-%m-%d")][YEARMONTH < 201903, ]
ggplot(pastCustomers, aes(TransactionMonth, numCustomers, color = Store_type)) +
geom_line() +
labs(x = "Month", y = "Total number of Customers", title = "Total Customers by Month")
```



Good, the trend in number of customers is also similar. Let's now assess the impact of the trial on sales.


```
#### Scale pre-trial control sales to match pre-trial trial store sales
scalingFactorForControlSales <- preTrialMeasures[STORE_NBR == trial_store &
YEARMONTH < 201902, sum(totSales)]/preTrialMeasures[STORE_NBR == control_store &
YEARMONTH < 201902, sum(totSales)]
#### Apply the scaling factor
measureOverTimeSales <- measureOverTime
scaledControlSales <- measureOverTimeSales[STORE_NBR == control_store, ][ ,
controlSales := totSales * scalingFactorForControlSales]

percentageDiff <- merge(scaledControlSales[, c("YEARMONTH", "controlSales")],
  measureOverTime[STORE_NBR == trial_store, c("totSales", "YEARMONTH")],
  by = "YEARMONTH")[, percentageDiff := abs(controlSales-totSales)/controlSales]
#### As our null hypothesis is that the trial period is the same as the pre-trial
#period, let's take the standard deviation based on the scaled percentage difference
#in the pre-trial period
#### Over to you! Calculate the standard deviation of percentage differences during
#the pre-trial period
stdDev <- sd(percentageDiff[YEARMONTH < 201902 , percentageDiff])
degreesOfFreedom <- 7
#### Trial and control store total sales
measureOverTimeSales <- measureOverTime
pastSales <- measureOverTimeSales[, Store_type := ifelse(STORE_NBR == trial_store,
  "Trial",
  ifelse(STORE_NBR == control_store, "Control", "Other stores"))
][, totSales := mean(totSales), by = c("YEARMONTH", "Store_type")]
```

```

[, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %/% 100,
                                     as.numeric(YEARMONTH) %% 100, 1,
                                     sep = "-"), "%Y-%m-%d")
][Store_type %in% c("Trial", "Control"), ]
#### Calculate the 5th and 95th percentile for control store sales.
pastSales_Controls95 <- pastSales[Store_type == "Control",
][, totSales := totSales * (1 + stdDev * 2)
][, Store_type := "Control 95th % confidence interval"]
pastSales_Controls5 <- pastSales[Store_type == "Control",
][, totSales := totSales * (1 - stdDev * 2)
][, Store_type := "Control 5th % confidence interval"]
#### Then, create a combined table with columns from pastSales, pastSales_Controls95 and pastSales_Controls5
trialAssessment <- rbind(pastSales, pastSales_Controls95, pastSales_Controls5)
#### Plotting these in one nice graph
ggplot(trialAssessment, aes(TransactionMonth, totSales, color = Store_type)) +
  geom_rect(data = trialAssessment[ YEARMONTH < 201905 & YEARMONTH > 201901 ,],
  aes(xmin = min(TransactionMonth), xmax = max(TransactionMonth), ymin = 0 , ymax =
  Inf, color = NULL), show.legend = FALSE) +
  geom_line(aes(linetype = Store_type)) +
  labs(x = "Month of operation", y = "Total sales", title = "Total sales by month")

```




The results show that the trial in store 86 is not significantly different to its control store in the trial period as the trial store performance lies inside the 5% to 95% confidence interval of the control store in two of the three trial months. Let's have a look at assessing this for the number of customers as well.

```

#### Scale pre-trial control customers to match pre-trial trial store customers
scalingFactorForControlCust <- preTrialMeasures[STORE_NBR == trial_store &
YEARMONTH < 201902, sum(nCustomers)]/preTrialMeasures[STORE_NBR == control_store &
YEARMONTH < 201902, sum(nCustomers)]
#### Apply the scaling factor
measureOverTimeCusts <- measureOverTime
scaledControlCustomers <- measureOverTimeCusts[STORE_NBR == control_store,
][ , controlCustomers := nCustomers* scalingFactorForControlCust
][, Store_type := ifelse(STORE_NBR == trial_store, "Trial",
ifelse(STORE_NBR == control_store,
"Control", "Other stores"))
]
#### Calculate the percentage difference between scaled control sales and trial
# sales
percentageDiff <- merge(scaledControlCustomers[, c("YEARMONTH",
"controlCustomers")],
measureOverTime[STORE_NBR == trial_store, c("nCustomers",
"YEARMONTH")],
by = "YEARMONTH"
)[, percentageDiff :=
abs(controlCustomers-nCustomers)/controlCustomers]
#### As our null hypothesis is that the trial period is the same as the pre-trial
# period, let's take the standard deviation based on the scaled percentage difference
# in the pre-trial period
stdDev <- sd(percentDiff[YEARMONTH < 201902 , percentDiff])
degreesOfFreedom <- 7
#### Trial and control store number of customers
pastCustomers <- measureOverTimeCusts[, nCusts := mean(nCustomers), by =
c("YEARMONTH", "Store_type")
][Store_type %in% c("Trial", "Control"), ]
#### Control store 95th percentile
pastCustomers_Controls95 <- pastCustomers[Store_type == "Control",
][, nCusts := nCusts * (1 + stdDev * 2)
][, Store_type := "Control 95th % confidence interval"]
#### Control store 5th percentile
pastCustomers_Controls5 <- pastCustomers[Store_type == "Control",
][, nCusts := nCusts * (1 - stdDev * 2)
][, Store_type := "Control 5th % confidence interval"]

trialAssessment <- rbind(pastCustomers, pastCustomers_Controls95,
pastCustomers_Controls5)
#### Plotting these in one nice graph
ggplot(trialAssessment, aes(TransactionMonth, nCusts, color = Store_type)) +
geom_rect(data = trialAssessment[ YEARMONTH < 201905 & YEARMONTH > 201901 ,],
aes(xmin = min(TransactionMonth), xmax = max(TransactionMonth), ymin = 0 , ymax =
Inf, color = NULL), show.legend = FALSE) +
geom_line() +
labs(x = "Month of operation", y = "Total number of customers", title = "Total
Customers by month")

```



It looks like the number of customers is significantly higher in all of the three months. This seems to suggest that the trial had a significant impact on increasing the number of customers in trial store 86 but as we saw, sales were not significantly higher. We should check with the Category Manager if there were special deals in the trial store that were may have resulted in lower prices, impacting the results. ## Trial store 88

```
#### Conduct the analysis on trial store 88.
measureOverTime <- data[, .(totSales = sum(TOT_SALES),
                             nCustomers = uniqueN(LYLT_Y_CARD_NBR),
                             nTxnPerCust = uniqueN(TXN_ID)/uniqueN(LYLT_Y_CARD_NBR),
                             nChipsPerTxn = sum(PROD_QTY)/uniqueN(TXN_ID),
                             avgPricePerUnit = sum(TOT_SALES)/sum(PROD_QTY)
                             )
, by = c("STORE_NBR", "YEARMONTH")] [order(STORE_NBR, YEARMONTH)]
#### Use the functions from earlier to calculate the correlation of the sales and
# number of customers of each potential control store to the trial store
trial_store <- 88
corr_nSales <- calculateCorrelation(preTrialMeasures, quote(totSales), trial_store)
corr_nCustomers <- calculateCorrelation(preTrialMeasures, quote(nCustomers), trial_store)
#### Use the functions from earlier to calculate the magnitude distance of the
# sales and number of customers of each potential control store to the trial store
magnitude_nSales <- calculateMagnitudeDistance(preTrialMeasures, quote(totSales),
trial_store)
magnitude_nCustomers <- calculateMagnitudeDistance(preTrialMeasures,
quote(nCustomers), trial_store)
#### Create a combined score composed of correlation and magnitude by merging the
# correlations table and the magnitudes table, for each driver.
corr_weight <- 0.5
```

```

score_nSales <- merge(corr_nSales, magnitude_nSales, by = c("Store1", "Store2")
                    )[, scoreNSales := 0.5*(corr_measure + mag_measure)]
score_nCustomers <- merge(corr_nCustomers, magnitude_nCustomers, by = c("Store1", "Store2")
                        )[, scoreNCust := 0.5*(corr_measure + mag_measure)]
#### Combine scores across the drivers by merging sales scores and customer scores,
# and compute a final combined score.
score_Control <- merge(score_nSales, score_nCustomers, by = c("Store1", "Store2"))
score_Control[, finalControlScore := 0.5*(scoreNSales+scoreNCust)]
#### Select control stores based on the highest matching store
#### (closest to 1 but not the store itself, i.e. the second ranked highest store)
#### Select control store for trial store 88
control_store <- score_Control[Store1 == trial_store,
                              ][order(-finalControlScore)][2, Store2]
control_store

```

```
## [1] 237
```


We've now found store 237 to be a suitable control store for trial store 88. Again, let's check visually if the drivers are indeed similar in the period before the trial.

We'll look at total sales first.

```


#### Visual checks on trends based on the drivers
#### For the period before the trial, create a graph with total sales of the trial
# store for each month, compared to the control store and other stores.
measureOverTimeSales <- measureOverTime
pastSales <- measureOverTimeSales[, Store_type := ifelse(STORE_NBR == trial_store,
                                                         "Trial",
                                                         ifelse(STORE_NBR == control_store, "Control", "Other stores"))
][, totSales := mean(totSales), by = c("YEARMONTH", "Store_type")]
[, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %/% 100,
                                     as.numeric(YEARMONTH) % 100, 1,
                                     sep = "-"), "%Y-%m-%d")][YEARMONTH < 201903]
ggplot(pastSales, aes(TransactionMonth, totSales, color = Store_type)) +
  geom_line(aes(linetype = Store_type)) +
  labs(x = "Month", y = "Total sales", title = "Total sales by month")

```



Great, the trial and control stores have similar total sales. Next, number of customers.

```
#### Visual checks on trends based on the drivers
#### For the period before the trial, create a graph with customer counts of the
# trial store for each month, compared to the control store and other stores.
measureOverTimeCusts <- measureOverTime
pastCustomers <- measureOverTimeCusts[, Store_type := ifelse(STORE_NBR == trial_store,
"Trial", ifelse(STORE_NBR == control_store,
"Control", "Other stores"))
][, numCustomers := mean(nCustomers), by = c("YEARMONTH", "Store_type")]
][, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %/% 100,
as.numeric(YEARMONTH) %% 100, 1, sep = "-"),
"%Y-%m-%d")][YEARMONTH < 201903, ]
ggplot(pastCustomers, aes(TransactionMonth, numCustomers, color = Store_type)) +
geom_line() +
labs(x = "Month", y = "Total number of Customers", title = "Total Customers by Month")
```


Total number of customers of the control and trial stores are also similar. Let's now assess the impact of the trial on sales.

```
#### Scale pre-trial control store sales to match pre-trial trial store sales
scalingFactorForControlSales <- preTrialMeasures[STORE_NBR == trial_store &
YEARMONTH < 201902, sum(totSales)]/preTrialMeasures[STORE_NBR == control_store &
YEARMONTH < 201902, sum(totSales)]
#### Apply the scaling factor
measureOverTimeSales <- measureOverTime
scaledControlSales <- measureOverTimeSales[STORE_NBR == control_store, ][ ,
controlSales := totSales * scalingFactorForControlSales]
#### Calculate the absolute percentage difference between scaled control sales and
# trial sales
percentageDiff <- merge(scaledControlSales[, c("YEARMONTH", "controlSales")],
                        measureOverTime[STORE_NBR == trial_store, c("totSales", "YEARMONTH")],
                        by = "YEARMONTH")[, percentageDiff := abs(controlSales-totSales)/controlSales]
#### As our null hypothesis is that the trial period is the same as the pre-trial
#period, let's take the standard deviation based on the scaled percentage difference
#in the pre-trial period
stdDev <- sd(percentageDiff[YEARMONTH < 201902 , percentageDiff])
degreesOfFreedom <- 7
#### Trial and control store total sales
measureOverTimeSales <- measureOverTime
pastSales <- measureOverTimeSales[, Store_type := ifelse(STORE_NBR == trial_store,
"Trial",
ifelse(STORE_NBR == control_store, "Control", "Other stores"))
][, totSales := mean(totSales), by = c("YEARMONTH", "Store_type")]
```

```

[, TransactionMonth := as.Date(paste(as.numeric(YEARMONTH) %/% 100,
                                     as.numeric(YEARMONTH) %% 100, 1,
                                     sep = "-"), "%Y-%m-%d")
][Store_type %in% c("Trial", "Control"), ]
#### Control store 95th percentile
pastSales_Controls95 <- pastSales[Store_type == "Control",
][, totSales := totSales * (1 + stdDev * 2)
][, Store_type := "Control 95th % confidence interval"]
#### Control store 5th percentile
pastSales_Controls5 <- pastSales[Store_type == "Control",
][, totSales := totSales * (1 - stdDev * 2)
][, Store_type := "Control 5th % confidence interval"]
#### Combine the tables pastSales, pastSales_Controls95, pastSales_Controls5
trialAssessment <- rbind(pastSales, pastSales_Controls95, pastSales_Controls5)
#### Plot these in one nice graph
ggplot(trialAssessment, aes(TransactionMonth, totSales, color = Store_type)) +
  geom_rect(data = trialAssessment[ YEARMONTH < 201905 & YEARMONTH > 201901 ,],
  aes(xmin = min(TransactionMonth), xmax = max(TransactionMonth), ymin = 0 , ymax =
  Inf, color = NULL), show.legend = FALSE) +
  geom_line(aes(linetype = Store_type)) +
  labs(x = "Month of operation", y = "Total sales", title = "Total sales by month")

```




The results show that the trial in store 88 is significantly different to its control store in the trial period as the trial store performance lies outside of the 5% to 95% confidence interval of the control store in two of the three trial months. Let's have a look at assessing this for number of customers as well.

```

#### This would be a repeat of the steps before for total sales
#### Scale pre-trial control store customers to match pre-trial trial store
# customers
scalingFactorForControlCust <- preTrialMeasures[STORE_NBR == trial_store &
YEARMONTH < 201902, sum(nCustomers)]/preTrialMeasures[STORE_NBR == control_store &
YEARMONTH < 201902, sum(nCustomers)]
#### Apply the scaling factor
measureOverTimeCusts <- measureOverTime
scaledControlCustomers <- measureOverTimeCusts[STORE_NBR == control_store,
][, controlCustomers := nCustomers* scalingFactorForControlCust
][, Store_type := ifelse(STORE_NBR == trial_store, "Trial",
ifelse(STORE_NBR == control_store,
"Control", "Other stores"))
]
#### Calculate the absolute percentage difference between scaled control sales and
# trial sales
percentageDiff <- merge(scaledControlCustomers[, c("YEARMONTH",
"controlCustomers")],
measureOverTime[STORE_NBR == trial_store, c("nCustomers",
"YEARMONTH")],
by = "YEARMONTH"
)[, percentageDiff :=
abs(controlCustomers-nCustomers)/controlCustomers]
#### As our null hypothesis is that the trial period is the same as the pre-trial
# period, let's take the standard deviation based on the scaled percentage difference
# in the pre-trial period
stdDev <- sd(percentDiff[YEARMONTH < 201902 , percentDiff])
degreesOfFreedom <- 7 # note that there are 8 months in the pre-trial period hence
# 8 - 1 = 7 degrees of freedom
#### Trial and control store number of customers
pastCustomers <- measureOverTimeCusts[, nCusts := mean(nCustomers), by =
c("YEARMONTH", "Store_type")
][Store_type %in% c("Trial", "Control"), ]
#### Control store 95th percentile
pastCustomers_Controls95 <- pastCustomers[Store_type == "Control",
][, nCusts := nCusts * (1 + stdDev * 2)
][, Store_type := "Control 95th % confidence interval"]
#### Control store 5th percentile
pastCustomers_Controls5 <- pastCustomers[Store_type == "Control",
][, nCusts := nCusts * (1 - stdDev * 2)
][, Store_type := "Control 5th % confidence interval"]
#### Combine the tables pastSales, pastSales_Controls95, pastSales_Controls5
trialAssessment <- rbind(pastCustomers, pastCustomers_Controls95,
pastCustomers_Controls5)
#### Plotting these in one nice graph
ggplot(trialAssessment, aes(TransactionMonth, nCusts, color = Store_type)) +
geom_rect(data = trialAssessment[ YEARMONTH < 201905 & YEARMONTH > 201901 ],,
aes(xmin = min(TransactionMonth), xmax = max(TransactionMonth), ymin = 0 , ymax =
Inf, color = NULL), show.legend = FALSE) +
geom_line() +
labs(x = "Month of operation", y = "Total number of customers", title = "Total
number of customers by month")

```



Total number of customers in the trial period for the trial store is significantly higher than the control store for two out of three months, which indicates a positive trial effect. ## Conclusion We've found control stores 233, 155, 237 for trial stores 77, 86 and 88 respectively. The results for trial stores 77 and 88 during the trial period show a significant difference in at least two of the three trial months but this is not the case for trial store 86. We can check with the client if the implementation of the trial was different in trial store 86 but overall, the trial shows a significant increase in sales. Now that we have finished our analysis, we can prepare our presentation to the Category Manager.