

Problem E. Yet Another Two Integers Problem

Time limit 1000 ms
Mem limit 262144 kB

You are given two integers a and b .

In one move, you can choose some **integer** k from 1 to 10 and add it to a or subtract it from a . In other words, you choose an integer $k \in [1; 10]$ and perform $a := a + k$ or $a := a - k$. You may use **different** values of k in different moves.

Your task is to find the **minimum** number of moves required to obtain b from a .

You have to answer t independent test cases.

Input

The first line of the input contains one integer t ($1 \leq t \leq 2 \cdot 10^4$) — the number of test cases. Then t test cases follow.

The only line of the test case contains two integers a and b ($1 \leq a, b \leq 10^9$).

Output

For each test case, print the answer: the minimum number of moves required to obtain b from a .

Sample 1

Input	Output
6	0
5 5	3
13 42	2
18 4	92
1337 420	87654322
123456789 1000000000	9150
100500 9000	

Note

In the first test case of the example, you don't need to do anything.

In the second test case of the example, the following sequence of moves can be applied: $13 \rightarrow 23 \rightarrow 32 \rightarrow 42$ (add 10, add 9, add 10).

In the third test case of the example, the following sequence of moves can be applied: $18 \rightarrow 10 \rightarrow 4$ (subtract 8, subtract 6).