

Microbiology of Fermented Foods & Beverages - MFFB

Notes for the course NFOK14019U, at the University of Copenhagen September-November 2024

Lucas Daniel Paz Zuleta, TZS159

MSc student at University of Copenhagen

Last compiled: 09-10-2024

Contents

| | | |
|----------|--|----------|
| 1 | Course description | 1 |
| 1.1 | Course content | 1 |
| 1.1.1 | Learning objectives | 1 |
| | Knowledge | 2 |
| | Skills | 2 |
| | Competences | 2 |
| 1.1.2 | Teaching and learning methods | 2 |
| 1.1.3 | Exam | 2 |
| 2 | Lecture Notes | 3 |
| 2.1 | 04.09.24 - Microscope | 3 |
| 2.1.1 | Parts of the microscope | 3 |
| 2.1.2 | Contrast methods | 4 |
| 2.1.3 | How to use the microscope | 4 |
| 2.2 | 04.09.24 - Yeast: Cytology, taxonomy and physiology | 5 |
| 2.2.1 | Background | 5 |
| 2.2.2 | Eukaryotes | 5 |
| 2.2.3 | Organelles | 5 |
| 2.2.4 | Methodologies for species identification of yeast | 6 |
| 2.2.5 | Yeast taxonomy | 6 |
| 2.2.6 | Substrates for yeasts | 7 |
| 2.2.7 | Conventional methods for identification of yeast | 7 |
| 2.2.8 | Macro-morphological characteristics | 8 |
| 2.2.9 | Micro-morphological characteristics | 9 |
| 2.2.10 | Assimilation and fermentation | 9 |
| 2.2.11 | Vegetative reproduction (genera) | 10 |
| 2.2.12 | Lecture conclusion | 11 |
| 2.3 | 09.09.24 - Taxonomy and the species concept | 11 |
| 2.3.1 | Lecture content | 11 |
| 2.3.2 | Taxonomy and phylogeny | 11 |
| 2.3.3 | Gram-positive & -negative bacteria | 12 |
| 2.3.4 | Differentiating between bacterial species | 13 |
| 2.4 | 11.09.24 - Methodologies for identification and typing | 13 |
| 2.4.1 | Intended leaning outcomes | 13 |
| 2.4.2 | Background | 14 |
| 2.4.3 | Methodology overview | 14 |

| | | |
|----------|---|-----------|
| 2.5 | 16.09 - From 1 st to 3 rd generation DNA sequencing strategies | 15 |
| 2.5.1 | Sanger Sequencing of DNA 1 st generation sequencing | 15 |
| 2.5.2 | Next generation sequencing | 15 |
| | Sequencing by synthesis | 16 |
| | Bridge amplification | 16 |
| 2.5.3 | 3 rd generation sequencing | 16 |
| 3 | Laboratory Exercises | 17 |
| 4 | Lecture Exercises | 18 |
| 4.1 | 11.09.24 - Exercise 1 - Replication, Transcription and Translation in prokaryotes | 18 |
| 4.1.1 | Replication in prokaryotes | 18 |
| 4.1.2 | Transcription in prokaryotes | 19 |
| 4.1.3 | Translation in prokaryotes | 19 |
| 4.2 | 11.09.24 - Exercise 2 - PCR and Restriction Enzymes | 19 |
| 4.2.1 | What is the size (in base pairs) of the PCR-product? | 20 |
| 4.2.2 | How many copies of DNA do you have after 4 PCR cycles if you start with 3 copies of the DNA-fragment to be copied (considering ideal conditions)? | 20 |
| 4.2.3 | What is the size of 2 fragments after cutting with HpyCH4V? | 20 |
| 4.2.4 | Place the bands that you would observe after running your product on the an agarose gel (path 1 before cutting with enzyme, path 2 after cutting) | 20 |
| 4.3 | 11.09.24 - Exercise 3 - Primer3plus | 22 |
| 4.4 | 11.09.24 - Exercise 4 - Primers check and restriction enzymes | 23 |
| 4.4.1 | What <i>Leuconostoc</i> strain will the primers align to? | 23 |
| 4.4.2 | What would be the PCR product size (bp) if we use those primers against that species? | 23 |
| 4.4.3 | What will be the size of DNA products after digesting the amplicon with Hind III restriction enzyme? | 24 |
| 4.4.4 | What are the reaction conditions requires for ApeKI? | 24 |
| 4.4.5 | What is the digestion pattern of AclI? | 24 |
| 4.4.6 | Using ENDMEMO | 24 |
| 5 | Litterature resumees | 26 |
| 5.1 | 1. Lecture | 26 |
| 5.1.1 | Article 1 - Fermented Foods as Experimentally Tractable Microbial Ecosystems | 26 |
| 5.1.2 | Article 2 - Diversity of Microorganisms in Global Fermented Foods and Beverages | 29 |
| 5.1.3 | Article 3 - Molecular biology of the cell; chapter 6 | 35 |
| A | Appendix | 46 |

Chapter 1

Course description

1.1 Course content

The course has major focus on the microorganisms involved in the processing of various fermented foods and beverages. The course includes the taxonomy of important microorganisms especially lactic acid bacteria (LAB) and yeasts covering both phenotypic characteristics and molecular typing techniques for their identification. Fermentation is introduced as a sustainable green technology and innovative technologies to improve fermented foods and beverages are considered.

Methodologies for isolation will be covered, including both culture dependent and culture independent techniques. Techniques such as high-throughput sequencing and digital programmes for bioinformatics are applied.

Various fermentation techniques is introduced covering the use of starter cultures and other fermentation techniques such as back-slopping. Further, the role of fermentation in sustainable food production and in prevention of food waste will be discussed.

An introduction to various fermented foods and beverages will be given including products such as cheese, bread, wine and beer as well as a number of traditional indigenous fermented food and beverages. Focus will additionally be on microbial interactions including topics such as quorum sensing, bacteriocin formation, etc.

1.1.1 Learning objectives

The objective of the course is to give the students a thorough knowledge on the microbiology behind production of fermented food and beverages and to give the students skills within isolation and identification of microorganisms occurring in these products. Additionally the students will be able to evaluate the functionalities and applications of microbial starter cultures.

Knowledge

- Show overview of fermented food and beverages in general and the microorganisms involved in their production
- Describe important groups of microorganisms identified from fermented food and beverages
- Comprehend microbial taxonomic systems
- Describe microbial interactions and their importance in food systems
- Reflect on microbial cytology and physiology
- Define molecular techniques for identification and typing to species and strain level

Skills

- Apply procedures for isolation and identification of the predominant microorganisms in fermented food and beverages
- Explain at the molecular level the behaviour and interactions between various groups of microorganisms
- Assess the most important parameters leading to optimal product quality and food safety
- Apply food fermentation to develop innovative food products

Competences

- Predict the composition of the microbiota of specific fermented food and beverages
- Discuss presumed functionalities of microorganisms in fermented food or beverages related to product quality and food safety
- Communicate and work independently on own data and discuss the results in relation to existing literature

1.1.2 Teaching and learning methods

Lectures, theoretical and laboratory practicals. The lectures will introduce issues of importance for the understanding of microbial behaviour during production of fermented food and beverages. The theoretical and laboratory practicals will give the students practice on how to identify various microorganisms from fermented food and beverages including skills within various molecular techniques and digital tools for bioinformatic purposes. Knowledge on food innovation will be obtained throughout the course.

1.1.3 Exam

At the time of the oral examination, 20 min, one theoretical question is drawn, and the examination proceeds without preparation time. Two weeks before the exam, the questions will be given to the students. At the oral exam, the drawn question and the curriculum will account for 75% of the grade. The discussion of the laboratory work and the experimental results will account for 25% of the grade.

Chapter 2

Lecture Notes

2.1 04.09.24 - Microscope

2.1.1 Parts of the microscope

The Microscope is put together by different components, as shown in Figure 2.1.



Figure 2.1: The different components of a microscope.

The main components are:

- Ocular

This is the set of lenses that you peer into, when you use a microscope. The “only” purpose of this part is to relay the intermediate image created by the objective to the retina of our eyes, i.e. to make us see an image. The ocular also magnifies the intermediate image with the factor of the ocular (usually around 10x); however, increasing the magnification of the ocular does not increase the resolving power (finer details of the specimen will not be revealed)

- Objective

Resolves the structure of the specimen, creating an intermediate image inside the microscope. The image is magnified with the factor depicted on the objective (often 10X, 40X and 100X). The objectives can be changed by rotating the objective turret, but this must be done cautiously to avoid mechanical damage. Objectives fitted for phase contrast will also be labelled with the size of the phase ring (usually this is depicted on the objective as either PH1, PH2 or PH3)

- Microscope stage
The (translation) microscope stage holds the specimen firmly with a specimen holder, and allows the specimen to be moved in fine increments in two dimensions
- Condenser
 - * The condenser is located underneath the microscope stage. The purpose of this optical part is to illuminate our specimen optimally
 - * Our microscopes for microbiological work have a condenser with a rotating turret that enables the use of phase contrast imaging, as a specific annular ring in this turret must be matched with the size of the phase ring in the objective. There is a cut-out on the front of the turret that depicts the actual selection. In this picture the Ph2 ring has been selected
 - * Apart from the phase rings, the turret also includes a blank opening that allows for traditional brightfield observation depicted with a 0, and a setting for darkfield depicted with a DF

2.1.2 Contrast methods

Brightfield

This simply magnifies the specimen. However it is almost impossible to see cells, without a staining procedure, that selectively stains parts of the cells. Quite often, a stained specimen is also fixed, which means that the cells have been killed, so for live specimens (e.g. live bacteria), we need a different contrast method.

Darkfield

This is only useful for 10X objectives or smaller, so for observing virtually all living cells.

Phase contrast

This is the best method. The theory behind phase contrast is a little complicated (but hey - the guy won the Nobel Prize), but you can utilize phase contrast without knowing the theory. All you have to remember is:

- The number on the phase objective and the number on the condenser turret must match to obtain a proper image (e.g. Ph 2 for the 40 x objective)
- The Field diaphragm in the bottom of the microscope should be completely open

2.1.3 How to use the microscope

Preparing the microscope

Set the ocular distance to match YOUR interpupillary distance; The distance between the eyes of people differs a lot. Our microscopes are equipped with two eyepieces, in order for you to look down the microscope in the same way as you look through a binocular, and ultimately the same way as you look on the world as such. Therefore, the first thing to do when you approach a microscope is to adjust the ocular distance to your individual interpupillary distance. The ocular distance can be read on a mm-scale (commonly 50-75) between the oculars.

The interpupillary distance is extremely easy to obtain, because you simply ask a friend to measure the distance between your pupils with an ordinary ruler. Memorise this value (because it is the same for the rest of your life), and adjust the oculars to your distance before you do anything else. This increases the comfort of using a microscope tremendously.

2.2 04.09.24 - Yeast: Cytology, taxonomy and physiology

2.2.1 Background

Yeast are quite different from bacterias, first of all, they are bigger than bacterias. Yeast are also eukaryotic, which means that they have a nucleus, and they have a lot of organelles. The yeast *Saccharomyces cerevisiae* has a pair of 16 chromosomes, which is quite a lot for a yeast and thus making it a non-haploid. The many differences from bacteria will influence their identification, their growth characteristics as well as the interactions with the food matrix. Proper identification of yeast is therefore important for the correct use of starter cultures as well as for spoilage yeasts.

2.2.2 Eukaryotes

Since yeast are Eukaryotes, here are some key features of Eukaryotes:

- Eukaryotes contain far more DNA than prokaryotes (x 100-1000). Yeast genome sizes range from 10-15 Mb and encode 5000-10000 genes
- The DNA is organised in a number of chromosomes
- The chromosomes are organised in a nucleus
- The presence of mitochondria
- The presence of different organelles such as: the Golgi apparatus and the endoplasmic reticulum
- Besides partly being involved in killer toxin production, the function of plasmids in yeasts is not really known

2.2.3 Organelles

- **CW: Cell wall (physical protection, adhesion etc.)**
- **P: Periplasm**
- **CM: Plasma membrane (transport in/out of the cytoplasm)**
- **CMI: Invagination**
- **BS: Bud scar**
- **C: Cytoplasm**
- **N: Nucleus (DNA in chromosomes)**
- **M: Mitochondrion (ATP synthesis, enzymes)**
- **S: Secretory vesicles**
- **V: Vacuole (protein trafficking, storage)**
- **PER: Peroxisome**
- **ER: Endoplasmic reticulum**
- **G: Golgi apparatus**

In this course, we will focus on the cell wall, the plasma membrane, the nucleus, the mitochondrion and the vacuole. Though the other listed organelles are important, they are not as important for the physiology of yeast.

Cell wall

The cell wall gives the yeast physical protection against the environment. It also maintains the shape of the yeast cell, and it is important for the osmotic stability of the yeast cell. The cell wall also has permeability barriers, which means that solutes larger than 620-760 Da can not pass the cell wall. The cell wall also has enzyme support, cation binding, cell-cell recognition and adhesion to other surfaces.

Plasma membrane

The primary functions of the yeast plasma membrane are to dictate what enters and what leaves the cytoplasm i.e. to be a primary barrier for passage of hydrophilic molecules and prevent cytoplasmic contents to mix with the environment. The plasma membrane is also important for the uptake of nutrients, the excretion of waste products, the maintenance of the cell shape and the cell wall, and the cell division.

Nucleus

2.2.4 Methodologies for species identification of yeast

There are various methods for species identification of yeast, including:

- Micro- and macromorphological identification added phenotypic characterisation
- Sequencing of the D1/D2 domain of the 26S rRNA gene (> 99% homology = belong to the same species)
- Sequencing of multiple genes
- Sequencing of the ITS region (mostly moulds and yeasts identified by metagenomics)
- Identification of protein patterns by MALDI-TOF MS
- Analysis of biomass by Fourier Transform Infrared Spectroscopy (FTIR)
- Others...

2.2.5 Yeast taxonomy

The taxonomy of yeast is quite complex, but it is important to know the different levels of taxonomy. The taxonomy of yeast is as follows:

- Kingdom: Fungi (Both yeast and moulds are fungi)
- **Phylum:** e.g Ascomycota // Basidiomycota
- **Class:** e.g *Saccharomycetes*
- **Order:** e.g *Saccharomycetales*
- Family: e.g *Saccharomycetaceae*
- Genus: e.g *Saccharomyces*
- Species: e.g *Saccharomyces cerevisiae*

The following is not universally recognized, but are used in some contexts:

- Subspecies: e.g *Saccharomyces cerevisiae* subsp. *cerevisiae*
- Variety: e.g *Saccharomyces cerevisiae* var. *cerevisiae*
- Strain: e.g *Saccharomyces cerevisiae* subsp. *cerevisiae* strain S288C

2.2.6 Substrates for yeasts

Following are some examples of some substrates which are greatly used for the propagation of yeasts:

- Malt yeast extract Glucose Peptone agar (MYGP)
- Yeast extract Peptone Glucose (YPG) agar
- Malt Agar (MA)
- Lysine agar a.o. (for non-*Saccharomyces* yeasts)
- WLN medium (allows separation by coloration)
- Others...

There are some stress tolerant yeasts where high concentrations of substrates are needed in the media for the growth of specific yeast. Some examples of medias for stress tolerant yeasts can be seen in 2.2.

| Purpose | Media |
|--|---|
| Moderately xerotolerant yeasts | Yeast nitrogen base + sodium chloride (10% (w/v)) + glucose (5 (w/v)) |
| Xerotolerant yeast | 50% (w/v) glucose-yeast extract agar |
| Xerotolerant yeast from foods of high salt and sugar | Wort agar + sucrose (3.5% w/v) + glucose (1.0% w/v) |
| <i>Zygosaccharomyces rouxii</i> | Potato-dextrose agar + sucrose (60% w/v) |
| Sugar-tolerant yeast from concentrated orange juice | Glucose-citric acid-tryptone agar |
| Isolation and cultivation of xerotolerant yeast | Malt-extract agar + glucose (2, 20, 40, 50% w/v) |

Figure 2.2: Examples of medias for stress tolerant yeasts.

2.2.7 Conventional methods for identification of yeast

There are various methods for identification of yeast, here are a list of some of the most common methods:

- By their cultural characteristics
 - Solid media: colony morphology, color, size, shape, surface, margin, elevation, texture
 - Liquid media: turbidity, sediment, pellicle, ring, flocculation
- By their cell morphology and arrangement
 - Cell form and size
 - Vegetative reproduction by budding or fission
 - Pseudomycelium or true mycelium
- By their sexual characteristics
 - Perfect stage (*teleomorph*) or imperfect stage (*anamorph*)
 - By their formation and arrangement of spores
- Biochemical characteristics
 - Assimilation of carbon or nitrate sources (e.g. API20c, ID32c)
 - fermentation of carbohydrates
- Physical and chemical tolerance
 - Growth in the presence of 100 or 1000 ppm cycloheximide
 - Growth in the presence of 1% acetic acid
 - Growth on 50% glucose-yeast extract agar
 - Etc.

2.2.8 Macro-morphological characteristics

There are many different macro-morphological characteristics of yeast, some of the ones we will be using in this course laboratory exercises can be seen in figure 2.3.



Figure 2.3: Examples of macro-morphological characteristics of yeast that will be evaluated in the laboratory exercises.

Furthermore, the lecture included a list of some key aspects to keep in mind while looking at the macro-morphological characteristics colonies. These are:

- Tecture (mucoid, viscous, matted, coherent etc.)
- Color
- Size
- Surface (glistening, smooth, verrucose, rough etc.)
- Elevation
- Margin (entire, rhizoid etc.)
- Mycelium (x100 magnification)

2.2.9 Micro-morphological characteristics

Micro-morphological characteristics are also important when identifying yeast. Some of the key aspects are listed below:

- Cell form (spheroidal, ellipsoidal, ovoid, lemon-shaped, elongated, triangular etc.)
- Cell size (length, width)
- Cell arrangement (pairs, aggregates)
- Vegetative reproduction
 - Budding (monopolar, bipolar, multilateral)
 - Fission
- Mycelium
 - Pseudomycelium (not distinct septa)
 - True mycelium (distinct septa)

2.2.10 Assimilation and fermentation

Assimilation refers to the process by which an organisms takes in and incorporates nutrients from the environment. The assimilation of carbon compounds is important for the growth of yeast. Yeast assimilates carbon compounds by various pathways, like Krebs cycle (TCA pathway) or the glycolysis.

Fermentation is the process by which yeast converts sugars into alcohol and carbon dioxide. The fermentation of carbohydrates typically happens under anaerobic conditions. The fermentation process is a less efficient energy source than respiration. Nevertheless, without it, wine and beer could not be made. Brewers therefore exploit the yeasts fermentative capability by introducing yeast into an anaerobic environment.

Assimilation of carbon compounds

To propagate yeast while having assimilation in mind, the growth medium Yeast Nitrogen Base from "Difco" can be used. This medium contains:

- | | | |
|------------------|-------------------|--------------|
| • Hexoses | • Organic acids | • Alcohols |
| • Trisaccharides | • Disaccharides | • Glycosides |
| • Pentoses | • Polysaccharides | |

Fermentation of carbon compounds

Here are a list of some of the carbon compounds that yeast can ferment:

- Glucose
- Maltose
- Trehalose
- Galactose
- Raffinose
- Melibiose
- Saccharose
- Lactose
- Inulin

Assimilation of nitrogen compounds

To propagate yeast while having assimilation in mind, the growth medium Yeast Carbon Base from "Difco" can be used. This medium contains:

- Nitrate
- Ethylamine hydrochloride
- L-lysine
- Creatine
- Nitrite
- Cadaverine dihydrochloride
- Creatinine

2.2.11 Vegetative reproduction (genera)

Vegetative reproduction is the process by which a plant or fungus reproduces asexually. Yeast can reproduce vegetatively by budding or fission. Some examples of yeast vegetative reproduction is:

Bipolar

- *Hanseniaspora (Kloeckera)*
- *Rhodotorula**
- *Saccharomycodes*
- *Trichosporon*

Fission

- *Schizosaccharomyces*

Arthroconidia

- *Galactomyces (Geotrichum)*
- *Saccharomycopsis**

Enteroblastic Budding

- *Phaffia*

Multilateral

- *Candida*
- *Debaryomyces*
- *Dekkera (Brettanomyces)*
- *Kazachstania*
- *Kluyveromyces*
- *Pichia*
- *Rhodotorula**
- *Saccharomyces*
- *Saccharomycopsis**
- *Torulaspora*
- *Yarrowia*
- *Zygosaccharomyces*

*Two types of vegetative reproduction may be present

2.2.12 Lecture conclusion

- Yeasts are predominantly unicellular eukaryotes
- Yeasts might have both imperfect and perfect names
- Yeast taxonomy is changing continuously especially due to new molecular techniques
- Yeasts are currently mostly identified by sequencing of the D1/D2 region of the 26S rRNA gene – however, never forget to verify by examining morphology and include phenotypic tests
- Remember to consult the newest scientific literature in order to use up dated taxonomic names

2.3 09.09.24 - Taxonomy and the species concept

2.3.1 Lecture content

During this lecture the following topics will be the focus point.

- Definitions of taxonomy and phylogeny
- Phylogeny
- Microbial taxonomy: from domain to species
 - Species concept in Bacteria
 - Species concept in Yeast and Moulds
- Subspecies, variant and strains

2.3.2 Taxonomy and phylogeny

From figure 2.4 we can see that the common ancestor of Bacteria, Archaea and Eukaryota is the root of the tree, and the branches of the three domains is what sums up the tree.



Figure 2.4: The common ancestor of Bacteria, Archaea and Eukaryota and the sum of their respective branches

A definition **Taxonomy**, this is the naming and classification of organisms based on shared properties, which can be based on

- Genotypic properties
- phenotypic properties
- Phylogeny - evolutionary relationships

While working with taxonomy and phylogeny in a broader sense, some of the following terms are important to know:

- **Isolation:** The separation of a pure culture from a mixed culture
Sampling: The collection of a representative part of a population
Colony purification: The isolation of a single colony from a mixed culture
DNA extraction: The isolation of DNA from a pure culture
- **Identification:** The determination of the taxonomic position of an organism
Microscopy: The determination of the morphology of an organism through a microscope
Sequencing: The determination of the sequence of a gene or a genome

While looking at the phylogeny we can look deeper into the classification, in table 2.1 we can see an example of the classification of a Bacteria and Eukarya where we stop at species. This can be further divided into subspecies, variant and strains.

Table 2.1: An example of the classification of a Bacteria and Eukarya

| | | |
|---------|-----------------------|----------------------|
| Domain | Bacteria | Eukarya |
| Kingdom | None assigned | Fungi |
| Phylum | Firmicutes | Ascomycota |
| Class | Bacilli | Saccharomycetes |
| Order | Lactobacillales | Saccharomycetales |
| Family | Lactobacillaceae | Saccharomycetaceae |
| Genus | <i>Lactobacillus</i> | <i>Saccharomyces</i> |
| Species | <i>L. acidophilus</i> | <i>S. cerevisiae</i> |

In the following table, there will be listed some differences between prokaryotes and eukaryotes.

Table 2.2: Differences between prokaryotes and eukaryotes

| | Prokaryotes | Eukaryotes |
|---------------------|---|---|
| DNA | DNA is naked DNA is circular Usually no introns | DNA is bound to protein DNA is linear Usually has introns |
| Organelles | No nucleus No membrane-bound 70S ribosomes | Has a nucleus Membrane-bound 80S ribosomes |
| Reproduction | Binary fission Single chromosomes (haploid) | Mitosis & meiosis Chromosomes paired (diploid or more) |
| Average size | Smaller (1-5 μm) | Larger (10-100 μm) |

2.3.3 Gram-positive & -negative bacteria

(Almost) all bacteria can be divided into two groups, Gram-positive and Gram-negative bacteria. The difference between the two groups is the cell wall. The cell wall protects the bacteria e.g. against osmotic stress. The cell wall differs where the Gram-positive bacteria have a thick cell wall, while the Gram-negative bacteria

have a thin cell wall. The difference in the cell wall is due to the peptidoglycan layer, which is thicker in the Gram-positive bacteria than in the Gram-negative bacteria. The peptidoglycan layer is also the reason why the Gram-positive bacteria are more resistant to antibiotics than the Gram-negative bacteria. Here are some bulletpoints of the two groups:

- **Gram-positive bacteria**
 - Lactic Acid Bacteria (LAB)
 - Bacillus
- **Gram-negative bacteria**
 - Most Acetic Acid Bacteria (AAB)

2.3.4 Differentiating between bacterial species

There are several methods to differentiate between bacterial species, some of the most common methods are:

- **DNA:DNA hybridization (DDH):**
 - The DNA of two organisms is mixed and the degree of hybridization is measured.
 - If DDH > 70% the organisms are considered to belong to the same species
- **16S rRNA gene sequencing:**
 - The 16S rRNA gene is sequenced and compared to a database, if the sequence is > 97.5-98.5% identical to a known species, the organism is considered to belong to the same species
- **Whole genome sequencing:**
 - Average nucleotide identity (ANI) > 95-96% is considered to belong to the same species
 - Digital DNA-DNA hybridization (dDDH) > 70% is considered to belong to the same species

2.4 11.09.24 - Methodologies for identification and typing

2.4.1 Intended learning outcomes

After this lecture, the students should be able to:

- Get to know how different microbial identification and typing methods work
- Apply this knowledge to comprehend microbial identification and taxonomy
- Reflect on the application possibilities of the different methods for microbial identification and typing

2.4.2 Background

- When you want to know what microorganisms are in a fermented food
IDENTIFICATION
- Often identification is coupled with quantification (predominant/minor microorganisms)
- Many different methods are available
 - Make an overview of the most used methods
 - Elaborate the steps of the methods → insights into the structure/mechanistics
 - Show how the methods are applied
- All methods have strengths and limitations

Note: typing is identification of isolates below species level

2.4.3 Methodology overview

Culture dependent methods

This is the most common method for identification of microorganisms. The method is based on the growth of the microorganisms on a solid or liquid medium. Here are an example of a method which is based on this:

- Plating
- Isolation
- Purification (The next 4 steps are methods for identification)
- Finger print based¹
 - rep-PCR, RAPD, RFLP or PFGE (All these are based on PCR)
- Sequence based²
 - rRNA gene, MLST or MLSA
- Biochemical identification³
 - MALDI-TOF or FTIR
- Classical⁴
 - Macro/micro morphology, phenotypic tests (fermentation, assimilation or growth conditions)

Culture independent methods

- Direct extraction of DNA or RNA
- Molecular methods (the next two steps are both molecular methods methods)
- PCR¹
 - Species specific PCR
- qRT-PCR²
 - Next generation sequencing

2.5 16.09 - From 1st to 3rd generation DNA sequencing strategies

DNA is quite small, 2 nm thick. Though, if folded out, the DNA would be around 1.8 m long.

2.5.1 Sanger Sequencing of DNA 1st generation sequencing

5'- TACAAC**TGAGCG**ACT -3'

3'- ATGTTGACTCGCTGA -5'

Here we have a blue sequence and a black complementing sequence. If we want to sequence the blue sequence, we can use the black sequence as a primer. The primer will bind to the blue sequence, and the DNA polymerase will start to build a new strand of DNA. The DNA polymerase will build the new strand by adding nucleotides to the 3' end of the primer. The nucleotides are added in a 5' to 3' direction. The DNA polymerase will continue to add nucleotides until it reaches the end of the DNA strand. The DNA polymerase will then stop, and we will have a new strand of DNA.

Though with the Sanger sequencing, we will have a mix of DNA strands, where each strand will end with a different nucleotide. The DNA strands will then be separated by size, and the sequence can be read by the size of the DNA strands.

With Sanger sequencing, the -A, -C, -G and -T nucleotides will cut the DNA strand at different lengths, because it cuts at the respective letter. An example of how the Sanger sequencing works can be seen in figure 2.5.



Figure 2.5: An example of how the method of Sanger sequencing of DNA works with the given blue sequence

2.5.2 Next generation sequencing

Next generation sequencing (NGS), has been evolving a lot since extensive research started in the 90's. Today we have a lot of different NGS methods, one of the most common methods is the Illumina sequencing. The

Illumina sequencing is based on the Sanger sequencing, but with a few modifications. Their HiSeqX can sequence around 50 human genomes in a day with a cost sub 1000 USD/genome.

Sequencing by synthesis

The Illumina sequencing is based on the sequencing by synthesis. It needs a primer, but these are fairly simple to make. After the primer has been added, the DNA polymerase will start to build a new strand of DNA. The DNA polymerase will add nucleotides from -A, -T, -G, and -C, to the 3' end of the primer, and the nucleotides will be added in a 5' to 3' direction. The DNA polymerase will continue the flashing with nucleotides until it reaches the end of the DNA strand. The DNA polymerase will then stop, and we will have a new strand of DNA.

The maximum read out length of the Illumina is between 75 bp to 350 bp, but the average read length is around 150 bp.

Bridge amplification

A piece of DNA is attached to a glass slide, and the DNA is then amplified. The DNA is then denatured, and the DNA is then amplified again. The DNA is then denatured again, and the DNA is then amplified again. This process is repeated until the DNA is amplified enough. The DNA is then sequenced, and the sequence is read by the computer. This method grows exponentially like PCR, though the DNA is amplified in a half circle, and not in a straight line.

An example of how the Illumina bridge amplification works can be seen in figure 2.6.



Figure 2.6: An example of how the Illumina bridge amplification works.

2.5.3 3rd generation sequencing

Chapter 3

Laboratory Exercises

Chapter 4

Lecture Exercises

4.1 11.09.24 - Exercise 1 - Replication, Transcription and Translation in prokaryotes

For an automated version of answering this exercise, please see the file w2e1.py in the Python folder. This file will automatically replicate, transcribe and translate the given DNA sequence. Use the file as a template for future exercises if needed.

Figure 4.1 shows a DNA sequence which will be used in Exercise 1 and its sub tasks. The given DNA sequence can be seen in figure 4.1

5'-_____ -3'
3'- GATTAGGTA ACTGTGATT CGTACGTAACGTGACGATATTAGCATCCACGCATACAGACGATATGCATAGCTGATCATATCGCC-5'

Figure 4.1: Sequence of DNA which is used in Exercise 1

4.1.1 Replication in prokaryotes

Replicate the DNA sequence based on the anti-sense (template) strand written below using the following primer 5'- AUCCAUU -3' (This RNA complementary to the DNA sequence and is the defined -35 region, which will be used in the transcription subsection).

- While replicating the DNA, identify the leading and lagging strands. Furthermore remember that G and C are the most stable base pairs, A and T are the least stable base pairs
- The DNA sequence is replicated from the 3' to 5' direction
- Use the primer as to define where to start from

5'- XXXATCCA **TTGACA** CTAAGCATGCATTGCACTGC **TATAAT** CGTAGGT
GGCGT **ATG** TCTGCTATACGTATCGACTAGTA **TAG** CGG -3' (This is the replicated)

3'- GAT **TAGGTAA** CTGTGATTTCGTACGTAACGTGACGATATTAGCATCCACCGCATACAGACGATATGCATAGCTGATCATATCGCC-5' (This is the given)

So, to begin with, we will look at the given RNA primer, and convert it to a DNA primer. 5'-AUCCAUU-3' will be converted to 5'-ATCCATT-3' (since RNA has Uracil instead of Thymine).

Afterwards we will look at the given DNA sequence and find the complementary strand to it. The complementary strand will be the anti-sense strand. 5'-ATCCATT-3' will be converted to 3'-TAGGTA-5' (since A pairs with T, T pairs with A, C pairs with G, and G pairs with C).

Now we will highlight the given sequence in yellow and start translating the DNA sequence from the 3' to 5' direction. We will start from the primer and move towards the end of the sequence.

This concludes this part of the exercise.

4.1.2 Transcription in prokaryotes

Identify the Pribnow (-10 region) and (-35) regions in the above DNA double-helix (sense strand). Transcribe DNA into a Messenger-RNA (mRNA).

In this course the Pribnow box (-10 region) is defined as the region 5'-TATAAT-3' and the -35 region is defined as 5'-TTGACA-3'. We will therefore

We found the Pribnow box sequence and highlighted it in green. The mRNA strand will be transcribed from the anti-sense strand, so we will use the antisense strand to transcribe the mRNA until the -35 region which we highlighted in turquoise.

We can now go to the right end (10-ish letters) of the pribnow box and start transcribing the mRNA. In this course, the start codon is defined as the first AUG codon after the Pribnow box. The mRNA strand will be transcribed from the 3' to 5' direction.

The start codon is highlighted in apricot starting from the 13th letter to the right. The stop codon is highlighted in bittersweet.

The Pribnow 10 region is 5'- ATCCATT -3' and the -35 region is 5'- CGATAGG -3'.

Now that we have identified both the start- and stop codon, we can start to transcribe the mRNA strand. The first codon will start from number 10 after the the Pribnow box. The mRNA strand will therefore be:

5'- GUA UGU CUG CUA UAC GUA UCG ACU AGU AUA GCG -3'

4.1.3 Translation in prokaryotes

Translate the mRNA strand from transcription step into amino acids.

The mRNA strand is written in the last subtask, and have already been translated into codons. It's now possible to translate the codons to amino acids.

H₂N- Val - Cys - Leu - Leu - Tyr - Val - Ser - Thr - Ser - Iso - Ala -COOH

4.2 11.09.24 - Exercise 2 - PCR and Restriction Enzymes

A part of DNA fragment shown below is to be amplified using PCR and the primers F and R. Subsequently the PCR product is cut with restriction enzyme HpyCH4V and the resultant products are run on a gel. The

```

      20      40      60      80
5'- TGTCGGTGTG CCATAACCTA AGTCATAACG ATCCGTGAAT GCATACCCGT TTTGAATAAT TGAATCCAAG AAACCTTGAT CTGTACTTGA-3'
3'- ACAGCCACAC GGTATTGGAT TCAGTATTGC TAGGCACCTA CGTATGGGCA AAACCTATTA ACTTAGGTTC TTTGAACATA GACATGAACT-5'

```

Figure 4.2: Sequence of DNA which is used in Exercise 2

The two given primers are:

- Primer F: 5'- AACCT-3'
- Primer R: 5'-TGGAT-3' (3'-TAGGT-5')

4.2.1 What is the size (in base pairs) of the PCR-product?

The Primer F starts from the 3' end of the DNA sequence, and the Primer R starts from the 5' end of the DNA sequence. The PCR product will therefore be the DNA sequence between the two primers.

Primer F has to fit to the upper strand, and Primer R has to fit on the lower strand. The primers sequence are therefore:

- Primer F: 5'-AACCT-3'
- Primer R: 5'-TGGAT-3' → 3'-TAGGT-5'

Primer F starts from the 15th to the right and ends at the 19th base pair. Primer R starts from the 64th from the right and ends at the 68th base pair. The PCR product will therefore be the DNA sequence between the 19th and 64th base pair.

The size of the PCR-product is therefore $68 - 14 = 54$ base pairs.

4.2.2 How many copies of DNA do you have after 4 PCR cycles if you start with 3 copies of the DNA- fragment to be copied (considering ideal conditions)?

After each PCR cycle, the amount of DNA doubles. After 4 PCR cycles, the amount of DNA will be $2^4 = 16$ times the original amount of DNA. Though, we are starting with 3 copies of the DNA fragment, so the amount of DNA after 4 PCR cycles will be $3 \cdot 2^4 = 48$ copies of the DNA fragment.

4.2.3 What is the size of 2 fragments after cutting with HpyCH4V?

The restriction enzyme HpyCH4V recognizes the sequence 3'-ACGT-5'. The DNA sequence is cut at the recognition site, and the two fragments will be the DNA sequence before the recognition site and the DNA sequence after the recognition site.

5'- TGTCGGTGTG CCATAACCTA AGTCATAACG ATCCGTGAAT GCATACCCGT TTTGAATAAT TGAATC-
CAAG AAAGTTGTAT CTGTACTTGA-3'

3'- ACAGCCACAC GGTATTGGAT TCAGTATTGC TAGGCACTTA CGTATGGGCA AAAGTTATTA ACT-
TAGGTTT TTTGAACATA GACATGAAGT-5'

HpyCH4V will therefore cut the DNA sequence at number 41, and the two fragments will be:

5'- AACCTA AGTCATAACG ATCCGTGAAT GCATACCCGT TTTGAATAAT TGAATCCA-3'

3'- TTGGAT TCAGTATTGC TAGGCACTTA CGTATGGGCA AAAGTTATTA ACTTAGGT-5'

There will therefore be 27 base pairs in the first fragment and 27 base pairs in the second fragment.

4.2.4 Place the bands that you would observe after running your product on the an agarose gel (path 1 before cutting with enzyme, path 2 after cutting)

When you run the PCR product on a gel electrophoresis, you will see the DNA fragments separated based on their size. The smaller fragments will move faster through the gel than the larger fragments. This means that both the DNA fragment before and after cutting with the restriction enzyme will be visible on the gel since they doesn't have the same fragment size .

Therefore you will see two bands on the gel, one at 54 base pairs (before cutting) and one at 27 base pairs (after cutting). This can be seen in Figure 4.3.




Figure 4.3: Gel electrophoresis of the DNA fragments and the PCR products' bands

4.3 11.09.24 - Exercise 3 - Primer3plus

Based on the sequence below design primers that fulfill following conditions:

The following information is given:

- size (22-24bp)
- T_m (55-57)
- GC% (35-70)
- amplicon size (300-400bp)

The following sequence is given:

```
TTGCAAGCCCTGATTTAAGCATTACACCAGCATCATCAAAGTAATACGCATTTCCATTTTGATCAGTTAT
ATATTGGTCAAAAACCTTGATGACCTTTCTTATCGAAATAACAATTGTACCATCTTCGTTTTGTAAGAAG
CTCTTGACCAGTTCGATACCATTAGGTAAGAAGAAGTAGGTATGGTTATTAATCTTTTGCAAACCTGTTA
CTAAATGACCTGTTTTATCAAAAATAGTAATAGTTATTATCATCATCTTGAATAAATGCATTTTGGGCACG
ATAACCACTTAATGTATAATAGATGATTCCTTTATCATCGTGTGTAAAGCCAGTTTCTGACAGATCATT
GTTAACTGTTTTGGTAAGTAGTCACCATCCTCAGTGTTTCGAAACGACCTTAAAGTACTTATCAGAACCCA
TATCTTTCAATACGTATCCAGCACCTTTACCTTGGATGTTAGAGCCATTGAAGTACTTAGCCGCCCCTC
AGTAATCTTTACACTGCCATCAATTGGAACGCCAGTTGAGATTTGATTACCTTTAAATAGGGATGGATAG
AGTGCCCTGTAACCTCTTCTAAGAAGGCACCACCATAACATCTCTTGATATTGACCACCCCCACGACTTTGTA
CAACATATAAGGCATTGTCAATATCAGAATCTGTATCGTCATCTCCAAATGAATTTGTTCTTGTGACAGT
AGCTAATTCTTGCTCTGGCAAATTATAAATTTGGTCCGGCACCCAATCGGCAATGGCTTGAATACCGCTA
GCATGTAAGGCTTTAATAGCATCGCGCAACTGATCAGCAGTTCCATATTTTGTGCGTGTGCCATAACCTA
AGTCATAACGATCCGTGAATGCATACCCGTTTTGAATAATTGAATCCAAGAACTTGTATCTGTACTTGA
ACGATATTGTGGTGCCAATTGGAAGCTTGTACACCCCATTGCTTAAATTGGTCCGCATTCTGAGCGATG
```

ACTACGTTTGTATATTCACCTGCTGTCTGTAGCAAATGCTTGGAAGTTTGAGAAACCTTCGTAGATGACTT
 GAGAATCAAGAGCAGCGTTTGAATGGAACACTTTATCACTCGTGTGTTGTTGTATCAGAGGCCGTTTCG
 TGCATCTTGATCTTGTGCGCACCTACCGGAACCCAAACTGCCAAGTAACCAGAACTTGTGGATTTTGT
 ACACCATAAATTGATTCATTCGTAAAAATCAAATCGCCGTTAGCATCTGTGTACGCCACAGGTGCATTTT
 CATCAGTATCATAATAAGCTAATCCATCTGCAGTTGTTGATAACAAAGCACGATAAGCTTGGTTCTTATG
 AGCTGCCCCCATATGCAATGTGACAGTATGCCATCCTCTAATTGTAGCTCCGCGTTATTGCTGACGATG
 ACTCCAATACCTTCAGTACGCGTCTCAGATGTTCCAGTGTCAGAAGCCGTCATGGCATCTTTACCATAGC
 GAACACTTGTTAACACGTCATTACTATCAACGGACATCGATTGGCCACCAGCAACATATTGGACTCTAGC
 CTTACAGCAAAGTGTTAATCGCATCATAGTATGGTGACTTTGTTGCCATATATTGACCATCATCTGTATAT
 AAATCGCCATAATAGACACGAGGAACAGTATCCTTGTGGTTAGCAACATCGCATAAGCACTAGCCATAT
 TATATTGTGTGTACTTTTTGTCTGCTAATTTTTTCATCTTCATTGTATACTTTGAAAGCAGCTGCCAATTG

Use Primer3Plus (<http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi/>)

While setting the conditions, it's important to remember that the amplicon size is the size of the DNA fragment that will be amplified by the primers. This is named "Product range size" on the web-side under general settings.

Write answers in the table below:

Table 4.1: The given table which should be filled out with the information from the Primer3Plus web-side

| Primer | Sequence (5'-3') | Primer size (bp) | T _m (°C) | GC content% |
|---------|-------------------------|------------------|---------------------|-------------|
| Forward | TCAGTAATCTTTACACTGCCATC | 23 | 56.1 | 39.1 |
| Reverse | TTATGACTTAGGTTATGGCACAC | 23 | 56.1 | 39.1 |

The second part of this exercise was to use this information to deduct which bacteria this sequence belongs to. The sequence is from the bacteria *Leuconostoc mesenteroides*.

4.4 11.09.24 - Exercise 4 - Primers check and restriction enzymes

The following information is given:

- Primer F: 5'- GGAACCCAAACTGCCAAGT-3'
- Primer R: 5'-CCTCGTGTCTATTATGGCGATT-3'

Using "In silico PCR amplification" (<http://insilico.ehu.es/PCR/>) answer following questions:

4.4.1 What *Leuconostoc* strain will the primers align to?

We start by going to the web-side. Then we choose *Leuconostoc* and press "Next step". Afterwards we paste the primer sequences into the "Forward primer" and "Reverse primer" fields. We then choose "APPLY TO ALL *Leuconostoc*" and press "Amplify".

Then we can see 2 bands where the primers has traveled at 485 and 485 base pairs. The primers will therefore align to the 2 *Leuconostoc mesenteroides* strains.

4.4.2 What would be the PCR product size (bp) if we use those primers against that species?

Based on the bands that was produced the PCR product size will be 485 base pairs.

For the next subsections, the following information is given:

Using NEBcutter V2.0 <http://tools.neb.com/NEBcutter2/> retrieve the amplicon from "In silico PCR amplification" and answer following questions:

4.4.3 What will be the size of DNA products after digesting the amplicon with Hind III restriction enzyme?

To use the web-side you need to copy the DNA sequence from the "In silico PCR amplification" web-side and paste it into the "Enter DNA sequence" field and press submit. Then you look at the summary and find the restriction enzyme HindIII and hold the mouse over it to see where it cuts.

For the two *Leuconostoc* strains the HINDIII cuts at:

- Cuts at bp 167 (*Leuconostoc mesenteroides* subsp. *mesenteroides* ATCC 8293)
- Cuts at bp 167 (*Leuconostoc mesenteroides* subsp. *mesenteroides* J18)

This means that the DNA products will be 167- and 318 base pairs long after digesting the amplicon with HindIII restriction enzyme. This makes sense since the total sequence, for both, is 485 base pairs long. Thus $167+318 = 485$ base pairs.

4.4.4 What are the reaction conditions requires for ApeKI?

To answer this, the following table has to be filled out. The information can be found on the NEBcutter web-side by pressing "Enzyme list" and then "ApeKI". Then in the bottom right corner of the first table, there is a button called "NEV Restriction Enzyme - Activity/Performance Chart". Then find the enzyme ApeKI and press on the buffer, there all the needed information is found.

Table 4.2: The given table which should be filled out with the information from the NEBcutter web-side

| Buffer | NEBuffer™ r3.1 |
|---------------------|------------------------------------|
| Salt | 100 mM NaCl |
| Main | 50 mM Tris-HCl |
| pH | 7.9 at 25°C |
| Mg | 10 mM MgCl ₂ |
| rAlbumin | 100 µg/ml Recombinant Albumin |
| Reaction temp. (°C) | 75°C (Obtained from the solutions) |

4.4.5 What is the digestion pattern of AciI?

The AciI enzyme cuts at the following base pairs:

Primer F 5'-...AA↓CGTT...-3'

Primer R 3'-...TTGC↑AA...-5'

This information can be found on the page before you press "NEV Restriction Enzyme - Activity/Performance Chart".

4.4.6 Using ENDMEMO

For this exercise we need to use this web-side (<http://www.endmemo.com/bio/egel.php>) In the graph on the right place bands you would receive after digesting the amplicon with HindIII, ApeKI and AciI in one reaction.




Figure 4.4: Gel electrophoresis of the DNA fragments and the PCR products' bands

Here we simply need to insert a sequence and choose the right restriction enzymes. The bands will then be shown on the gel.

- HindIII cuts at 167 bp; bands will be at 167- and 318 bp
Blue color
- ApeKI cuts at 184 bp; bands will be at 184- and 301 bp
Green color
- AclI cuts at 233 bp; bands will be at 233- and 252 bp
Red color

Chapter 5

Literature resumees

5.1 1. Lecture

5.1.1 Article 1 - Fermented Foods as Experimentally Tractable Microbial Ecosystems

Introduction

The study of microbial communities faces significant challenges, largely due to their complexity and the difficulty in understanding how these multi-species communities are assembled, organized, and function. As a result, many aspects of microbial ecology remain unexplored.

The article highlights that the vast diversity and complexity of microbial communities are key factors contributing to our limited understanding. Additionally, the difficulty in isolating the majority of species from natural environments and cultivating them in vitro presents another notable challenge [3].

One approach to addressing these challenges is the development of simplified and controlled environments to study microbial communities. These controlled systems allow for reproducible results through measurable substrates, microbial growth, and community formation [3].

Several models with these properties have already been developed. These range from synthetic mixtures of microbial strains to more complex models composed of culture-able strains from free-living or host-associated communities, as well as naturally occurring communities that have been extensively sampled or perturbed in situ [3].

Fermented Foods as Experimentally Tractable Ecosystems

Fermented foods have been around for thousands of years and are an excellent example of how humans have optimized conditions to promote various properties of foods. Here are some of the main reasons why fermented foods have been produced for so long [3]:

- Good preserving effects
- Flavour can be improved
- Aroma can be enhanced
- Texture can be changed

Some factors for manipulating microbial growth are listed below [3]:

- Temperature
- Salinity
- Moisture

The article describes some microbial communities of fermented foods (MCoFFs) and some examples of these, which will be listed below [3]:

- Multi-species biofilms associated with surfaces
e.g. cheese rinds
- Suspended biofilms in liquid
e.g. kombucha, kefir, and vinegar
- Dispersed growth in liquid
e.g. lambic beers, natural wines, and yogurt
- Semi-solid substrates
e.g. kimchi and miso

Many MCoFFs have been studied in detail, and the article highlights that some of these communities have been shown to have bacterial- and fungi species that have co-evolved over time. The microbial diversity within and across fermented foods is vast, and the article emphasizes that these communities are excellent models for studying microbial ecology. In the article, cheese rind (the MCoFFs on the surface of the cheese) was used as an example of the characterization and development of MCoFFs, since on the surface of cheese the microbial communities form as it ages [3].

Using Fermented Foods to Link Patterns, Processes, and Mechanisms of Microbial Community Assembly

In order to understand how species come together to form a community within a given environment microbiologists have developed several approaches for both plant- and animal communities. The ones named in the article is listed below [3]:

- **Dispersal**
Contributions from the amount and timing of microbial propagules colonizing a habitat
- **Biotic selection**
Interactions between species
- **Abiotic selection**
Interactions between a species and the environment
- **Drift**
Stochastic (random) changes in the relative abundances of species within communities
- **Diversification**
Evolution of new species within communities

MCoFFs are ideal for linking microbial patterns with ecological processes. For instance, many fermented

foods exhibit clear microbial successions where early colonizers are replaced by other species. This process can be studied to understand how environmental changes and species interactions drive these transitions. New techniques like RNA sequencing and transposon mutagenesis have proven valuable in uncovering the molecular mechanisms behind these interactions. Additionally, tools like imaging mass spectrometry are enhancing our understanding of how microbial species communicate chemically [3].

Taste of Place? Microbial Biogeography of Fermented Foods

Though there are many different types of MCoFFs, many share some similarities. If a similar MCoFFs is used in two different cheese factories in two different locations, the microbial communities will be different, and with some distinct properties which can affect the texture, taste or aroma composition of the final product. This also enables the study of microbial biogeography, which is the study of how microbial communities are distributed across different locations. MCoFFs can therefore help linking patterns in microbial diversity from place to place. This is possible because each community has clear species abundance distributions [3].

Microbial Evolution in a Community Context

MCoFFs offer valuable systems for studying both ecological processes and microbial evolution. Microorganisms used in beer, wine, dairy, and sake production have been useful in understanding how microbes adapt to fermentation environments. Research suggests that these organisms often lose genes not needed in nutrient-rich settings, gain new traits through gene transfer, and adjust their metabolism to fit the fermentation niche [3].

Looking ahead, linking ecological and evolutionary processes could provide further insights. For example, studying long-term microbial communities could help explore how species interactions influence evolution. Experimentally testing coevolution within these communities could offer useful data on microbial adaptation over time [3].

Translation to Other Microbial Communities and Potential Applications

The study of microbial community assembly in traditional fermented foods (MCoFFs) can significantly enhance food quality and safety. Insights from these ecosystems can be applied to other microbial communities, such as the human microbiome. For example, the competitive mechanisms of *Lactobacillus reuteri*, found in both sourdough and human gut microbiomes, illustrate how patterns observed in MCoFFs can be relevant to other systems [3].

Research on cheese rind microbial diversity has revealed similarities with the human skin microbiome, suggesting that moisture is a major driver of surface biofilms in both environments. This indicates that ecological selection processes may be conserved across different microbial communities [3].

MCoFFs can directly impact human health as they are consumed and can interact with the human gut microbiome. While the probiotic effects of simplified MCoFFs like yogurt have been extensively studied, the broader microbial diversity in MCoFFs could introduce additional beneficial microbes and metabolites to the human digestive system. Studies have shown that MCoFFs can remain viable during passage through the human digestive tract [3].

Furthermore, MCoFFs offer valuable opportunities for designing synthetic microbial communities for applications in medicine, industry, and agriculture. Understanding the interactions within these communities can inform the construction of synthetic microbial ecosystems. By studying pre-existing MCoFFs, researchers can

learn about design principles and explore the potential for combining microbes from different MCoFFs into new, functional compositions [3].

Ultimately, research on food microbial communities could result in safer, more delicious foods and contribute to the development of essential principles for microbial community design [3].

5.1.2 Article 2 - Diversity of Microorganisms in Global Fermented Foods and Beverages

Introduction

The article describes how rice with fermented and non-fermented legumes is a staple diet in many counties in Asia. Wheat/barley-based breads/loaves followed by milk and fermented milk products, meat and fermented meats are more common in the western world (West Asia, Europe and North America). The staple diet in Africa and South America comprise of sorghum and maize with wild legume seeds, meat and milk products [2]. Consortia of microorganisms are found naturally in uncooked plant and animal materials, utensils, food containers, earthen pots and in general in the environment. When producing fermented foods, it is normal to introduce a starter culture, which contains functional microorganisms [2]. The microorganisms in fermented food has converted the chemical composition of the raw food, which results in a change in the nutritional value of the food, making it more enriched [2].

Microorganisms in fermented foods

Lactic acid bacteria (LAB) are one of the most important groups of microorganisms in fermented foods. They are used in the production of fermented foods and beverages. Some of the major genera of the LAB are listed below [2]:

- *Alkalibacterium*
- *Carnobacterium*
- *Enterococcus*
- *Lactobacillus*
- *Lactococcus*
- *Leuconostoc*
- *Oenococcus*
- *Pediococcus*
- *Streptococcus*
- *Tetragenococcus*
- *Vagococcus*
- *Weissella*

Bacillus has been found in alkaline-fermented foods in Asia and Africa. The species of *Bacillus* that are found in abundance in legume-based fermented foods are listed below [2]:

- *Bacillus amyloliquefaciens*
- *Bacillus circulans*
- *Bacillus coagulans*
- *Bacillus firmus*
- *Bacillus licheniformis*
- *Bacillus megaterium*
- *Bacillus pumilus*
- *Bacillus subtilis*
- *Bacillus subtilis variety natto*
- *Bacillus thuringiensis*

There has been reported several species of *Kocuria*, *Micrococcus* and *Staphylococcus* in fermented foods. The species of *Kocuria* are found in abundance in fermented milk products, fermented sausages, meat-, and fish products [2].

Yeasts are also associated with fermentation of foods and alcoholic beverages. The yeasts named in the article are listed below [2]:

- *Brettanomyces*
- *Candida*
- *Cryptococcus*
- *Debaryomyces*
- *Dekkera*
- *Galactomyces*
- *Geotrichum*
- *Hansenula*
- *Hanseniaspora*
- *Hyphopichia*
- *Issatchenkia*
- *Metschnikowia*
- *Saccharomyces*
- *Pichia*
- *Kazachstania*
- *Rhodotorula*
- *Saccharomyces*
- *Kluyveromyces*
- *Rhodospiridium*
- *Saccharomycopsis*
- *Schizosaccharomyces*
- *Sporobolomyces*
- *Torulaspora*
- *Torulopsis*
- *Trichosporon*
- *Yarrowia*
- *Zygosaccharomyces*

Filamentous molds are also found in fermented foods. They hold a major role in various fermented products in respect of enzyme production and in the degradation of anti-nutritive factors. The listed filamentous molds are listed below [2].

- *Actinomucor*
- *Amylomyces*
- *Aspergillus*
- *Monascus*
- *Mucor*
- *Neurospora*
- *Paracilomyces*
- *Penicillium*
- *Rhizopus*
- *Ustilago*

Taxonomic tools for identification of microorganisms from fermented foods

A commonly used method for profiling both culturable and non-culturable microbial populations from fermented foods is DNA extraction. The technique is based on the separation of PCR-amplified DNA fragments by denaturing gradient gel electrophoresis (DGGE). This method is both used for profiling bacterial populations and yeast populations in fermented foods. Also, in fermented foods a combination of Propidium MonoAzide (PMA) treatment, which is done before DNA-extraction, and molecular quantifying methods can be used to enumerate the viable microorganisms accurately [2]. Random amplification of polymorphic DNA (RAPD) is a typing method based on the genomic DNA fragment profiles amplified by rep-PCR, and is commonly used for differentiation of LAB strains from fermented foods and other bacteria or yeast in the consortia. This technique is able to differentiate between strains of the same species [2]. Amplified fragment length polymorphism (AFLP) is another technique which is used to identify and discriminate LAB strains. The technique is based on the selective amplification and separation of genomic restriction fragments [2].

DGGE techniques and temperature gradient gel electrophoresis (TGGE) are used to study the microbial diversity in fermented foods. The techniques are based on the separation of PCR-amplified 16s rDNA and 26s rDNA gene fragments by denaturing gradient gel electrophoresis. The techniques are used to study the microbial diversity in fermented foods. DGGE is quite time consuming and is not able to determine relative abundance of the dominant species, nor is it able to distinguish between viable and non-viable cells [2].

Next generation sequencing (NGS) are effective tools for studying the microbial diversity in fermented foods. The techniques are able to sequence all the genetic material present in a sample, providing a comprehensive view of the microbial community's diversity and functional potential. These techniques require extensive training and a well-equipped molecular laboratory [2].

In table 5.1 some of the NGS which is listed in the text are described. These NGS are used to study the microbial diversity in fermented foods [2].

Global fermented foods

A study reported approximately 3500 fermented food- and beverages, which were divided into 250 categories. Globally, fermented foods are classified into 9 groups based of the substrates used from the plant- or animal

Table 5.1: Some NGS techniques with their respective key-points

| Metagenomics | Phylobiomics | Metatranscriptomics |
|---|---|---|
| <ul style="list-style-type: none"> Sequences all the genetic material present in a sample, providing a comprehensive view of the microbial community's diversity and functional potential. Allows the study of microbial communities without the need to culture individual species, overcoming limitations of traditional microbiology methods. By sequencing entire genomes, metagenomics can identify functional genes involved in processes like metabolism, antibiotic resistance, or nutrient cycling. | <ul style="list-style-type: none"> Is primarily concerned with analyzing the evolutionary relationships between microbial species, often using 16S rRNA or other conserved genes. It helps characterize the diversity and phylogenetic structure of microbial communities, determining how different species are related within an ecosystem. This technique is useful for identifying species and classifying them taxonomically based on their evolutionary lineage. | <ul style="list-style-type: none"> Focuses on sequencing the RNA transcripts present in a microbial community, providing a snapshot of active genes and gene expression levels. Reveals what genes are actively being expressed in a community at a given time. It helps assess how microbial communities respond to environmental changes, stress, or disease by analyzing shifts in gene expression. |

source. These groups are listed in table 5.2 [2].

Table 5.2: The 9 groups of fermented foods and beverages

| Number | Group |
|--------|---|
| 1 | Fermented cereals |
| 2 | Fermented vegetables and bamboo shoots |
| 3 | Fermented legumes |
| 4 | Fermented roots/tubers |
| 5 | Fermented milk products |
| 6 | Fermented and preserved meat products |
| 7 | Fermented, dried and smoked fish products |
| 8 | Miscellaneous fermented products |
| 9 | Alcoholic beverages |

Fermented milk products

Fermented milk products can be classified into two main groups based on the microorganisms involved; Lactic fermentation and Fungal-lactic fermentations. The lactic fermentation is dominated by Lactic Acid Bacteria (LAB), including thermophilic- (e.g., yogurt, Bulgarian buttermilk), probiotic- (e.g., acidophilus milk, bifidus milk), and mesophilic types (e.g., natural fermented milk, cultured milk, cultured cream, cultured buttermilk). The Fungal-lactic fermentations involves cooperation between LAB and yeasts to produce products like alcoholic milks and moldy milks [2].

Natural fermentation is an ancient method of milk processing that uses raw or boiled milk to ferment spontaneously. Starter cultures are used to initiate fermentation and can be either primary or secondary. The primary cultures mostly consist of *Lactococcus lactis*, *Lactobacillus delbrueckii*, and *Streptococcus thermophilus*, which participate in acidification. The secondary cultures are used in cheese-making and include *Brevibacterium linens*, *Propionibacterium freudenreichii*, and *Penicillium camemberti*, which develop flavor and texture during ripening [2].

Non-Starter Lactic Acid Bacteria (NSLAB) are present in high numbers in fermented milk and include *Enterococcus durans*, *Lactobacillus casei*, and *Staphylococcus* spp [2].

Fermented cereal foods

Fermented cereal foods are prepared differently in various regions of the world. In Asia, rice is fermented to produce alcoholic beverages or food beverages using mixed cultures. In Europe, America, and Australia, cereals like wheat, rye, barley, and maize are fermented using natural fermentation or commercial baker's yeast. In Africa, fermented cereal foods are used as staples, complementary, and weaning foods for infants and young children [2]. In Europe, traditional methods of bread preparation are still practiced without using commercial baker's yeast. Yeasts and LAB conduct dough fermentation, resulting in sourdough breads with higher lactic acid and acetic acid content [2]. Some LAB as *Enterococcus*, *Lactococcus*, *Lactobacillus*, *Leuconostoc*, *Pediococcus*, *Streptococcus*, and *Weissella* are deeply involved in cereal fermentation. Also some yeasts have a significant role such as the strains of *Saccharomyces cerevisiae* are the principal yeast, but other non-*Saccharomyces* yeasts like *Candida*, *Debaryomyces*, and *Hansenula* is also found in cereal fermentation [2].

Fermented vegetable foods

Many vegetables such as leafy greens, radish, cucumbers, and young edible bamboo shoots, are perishable and seasonal, therefore they are traditionally fermented into edible products to ensure longer shelflife and safe consumption [2]. The main microorganisms involved in the fermentation of vegetables are LAB, which are responsible for the acidification of the product. The most common LAB in fermented vegetables are *Lactobacillus* and *Pediococcus*, followed by *Leuconostoc*, *Weissella*, *Tetragenococcus*, and *Lactococcus* [2]. Some examples of fermented vegetable products are listed below in table 5.3 [2]:

Table 5.3: Examples of fermented vegetable products

| Food | Description |
|------------|---|
| Kimchi | A Korean fermented vegetable product, with a characterized microbial profile of LAB |
| Sauerkraut | A German fermented cabbage product, with a reported species of LAB |
| Gundurk | have a native population of LAB. Made by fermenting leafy greens which then sun dried to create a sour and slightly bitter flavor |

Fermented soybeans and other legumes

There are two types of fermented soybean foods, bacillus-fermented and mould-fermented. The bacillus-fermented foods are fermented by *Bacillus* spp. (mostly *B. subtilis*), and are characterized by stickiness. The mould-fermented foods are fermented by filamentous molds, mostly *Aspergillus*, *Mucor*, and *Rhizopus*. These types of foods are concentrated in the "Kinema-Natto-Thua Nao (KNT)-triangle" region, which includes Japan, east Nepal, north-east India, and northern Thailand. Examples of bacillus-, mould-, and other fermented foods are listed in table 5.4 [2].

Table 5.4: Examples of Bacillus-, mould-, and other fermented foods

| Bacillus-fermented | | Mould-fermented | | Fermented foods | |
|--------------------|----------|-----------------|-----------|-----------------|--------|
| Foodstuff | Origin | Foodstuff | Origin | Foodstuff | Origin |
| Natto | Japan | Miso and shoyu | Japan | Locust beans | Africa |
| Kinema | India | Tempe | Indonesia | Black-grams | India |
| Chungkokjang | Korea | Douchi and sufu | China | Maseura | Nepal |
| Thua nao | Thailand | Doenjang | Korea | | |
| Pekok | Myanmar | Pekok | Myanmar | | |
| Sieng | Cambodia | Sieng | Cambodia | | |

Fermented meat products

Overall, fermented meat products are divided into two categories, whole meat pieces (e.g. jamón serrano) and chopped meat pieces (e.g. chorizo). The main microbial groups involved in fermenting meats are LAB, coagulase-negative cocci, and yeasts. The LAB are responsible for the acidification of the meat, while the coagulase-negative cocci are responsible for the ripening of the meat. The yeasts are responsible for the development of the flavor and aroma of the meat [2].

Fermented fish products

Traditionally fish is preserved by either fermentation, sun/smoke drying and salting. Both bacteria and yeasts have been reported from these preservation methods.

Fermented miscellaneous products

The article discusses various fermented foods and the microorganisms involved in their production. These foods include [2]:

- Vinegar
Produced from sugar or ethanol-containing substrates and hydrolyzed starchy materials by aerobic conversion to acetic acid.
Dominant microorganisms include *Acetobacter* species and yeast species such as *Candida* and *Saccharomyces*.
- Fermented Teas
Such as miang, puer tea, fuzhuan brick, and kombucha.
Microorganisms involved include *Aspergillus*, *Penicillium*, and *Saccharomyces* species.
- Nata
A delicacy from the Philippines, produced by *Acetobacter xylinum*.
Two types of nata are known: nata de piña and nata de coco.
- Chocolate
Produced from cocoa bean fermentation.
The predominant microorganisms are *Acetobacter pasteurianus*, *Lactobacillus plantarum*, and *Saccharomyces cerevisiae*.

Continues on the next page.

- **Pidan**

A preserved egg prepared from alkali-treated fresh duck eggs, with a strong hydrogen sulfide and ammonia smell.

Dominant microorganisms include *Bacillus cereus*, *Staphylococcus cohnii*, and *Staphylococcus epidermidis*.

Some common microorganisms found in miscellaneous fermented foods include [2]:

- LAB; such as *Lb. fermentum*, *Lb. plantarum*, and *Weissella* species.
- Acetobacter; such as *Acetobacter pasteurianus* and *Acetobacter xylinum*.
- Yeast; such as *Saccharomyces*, *Candida*, and *Hanseniaspora* species.
- Mold; such as *Aspergillus* and *Penicillium* species.

Amylolytic starters

Amylolytic starters are traditional cultures of microorganisms used to produce alcoholic beverages and fermented foods from starchy materials. These starters are composed of consortia of filamentous molds, amylolytic, and alcohol-producing yeasts and LAB, and are typically made with rice or wheat as the base [2].

The article described some of the different types of amylolytic starters which are used in various countries, including [2]:

- **Marcha** (India and Nepal)
- **Hamei, humao, phab** (India)
- **Men** (Vietnam)
- **Ragi** (Indonesia)
- **Bubod** (Philippines)
- **Chiu/chu** (China and Taiwan)
- **Loogpang** (Thailand)
- **Mae/dombae/buh/puh** (Cambodia)
- **Nuruk** (Korea)

Furthermore the article named some of the common microorganisms involved in amylolytic starters which included [2]:

- Filamentous molds; such as *Mucor circinelloides*, *Rhizopus chinensis*, and *Aspergillus oryzae*.
- Yeasts; such as *Saccharomyces cerevisiae*, *Pichia anomala*, and *Candida glabrata*.
- LAB; such as *Pediococcus pentosaceus*, *Lactobacillus bifermentans*, and *Lactobacillus brevis*.
- Amylase-producing bacilli; such as *Bacillus subtilis* and *Bacillus amyloliquefaciens*.
- Acetic acid bacteria; such as *Acetobacter orientalis* and *Acetobacter pasteurianus*.

Alcoholic beverages

Alcoholic beverages can be classified into 10 types based on their production methods and ingredients. These types include [2]:

- **Non-distilled and unfiltered beverages produced by amylolytic starters;** such as kodo ko jaanr (fermented finger millets) and bhaati jaanr (fermented rice) of India and Nepal.
- **Non-distilled and filtered beverages produced by amylolytic starters;** such as saké of Japan.
- **Distilled beverages produced by amylolytic starters;** such as shochu of Japan and soju of Korea.
- **Beverages produced by human saliva:** Such as chicha of Peru, which involves the use of amylase in human saliva.
- **Mono-fermentation beverages:** Produced by a single-strain fermentation, such as beer.
- **Honey-based beverages:** Such as tej of Ethiopia.
- **Plant-based beverages:** Such as pulque of Mexico, toddy of India, and kanji of India.
- **Malted beverages:** Produced by malting (germination), such as sorghum (“Bantu”) beer of South Africa, pito of Nigeria and Ghana, and tchoukoutou of Benin.
- **Fruit-based beverages without distillation:** Such as wine and cider.
- **Distilled fruit and cereal beverages:** Such as whisky and brandy.

5.1.3 Article 3 - Molecular biology of the cell; chapter 6

In this chapter

In this chapter of the book, the following topics will be further described:

The central dogma of molecular biology; The discovery of the structure of DNA in the 1950s has led to significant progress in cell and molecular biology. We now know the complete genome sequences for thousands of organisms, including humans and our extinct relatives, such as the Neanderthals. The flow of genetic information in cells is from DNA to RNA to protein, a principle known as the central dogma of molecular biology [1].

Variations in the central dogma; While the central dogma is universal, there are important variations between organisms in the way information flows from DNA to protein. In eukaryotic cells, RNA transcripts are subject to processing steps in the nucleus, including RNA splicing, before they are translated into protein [1].

The role of RNA; RNA plays a crucial role in the cell, not only as a template for protein synthesis but also as a structural and catalytic molecule. Some RNAs fold into precise three-dimensional structures, while others act as regulators of gene expression [1].

The complexity of genomes; The genomes of most multicellular organisms are surprisingly disorderly, reflecting their chaotic evolutionary histories. Genes are often arranged in a long string of alternating short exons and long introns, with small bits of DNA sequence that code for protein interspersed with large blocks of seemingly meaningless DNA [1].

Decoding genomes; Decoding genomes is a complex task, even with the aid of powerful computers. Cells in our body can automatically locate the beginning and end of genes and decipher when and where each gene is expressed, but researchers still face significant challenges in understanding how genomes are decoded and used [1].

The challenges of cell biology; Despite significant progress in cell and molecular biology, there is still much to be discovered about how the information stored in an organism’s genome produces even the simplest unicellular bacterium, let alone how it directs the development of a human with approximately 30,000 genes.

The next generation of cell biologists will face many fascinating challenges in understanding the complexities of genomes and how they are decoded and used [1].

From DNA to RNA

Transcription and translation are the means by which cells read out, or express, the genetic instructions in their genes [1].

Regulation of gene expression is when the cells can synthesize a large amount of protein from a gene when necessary, and genes can be transcribed and translated with different efficiencies, allowing the cell to make vast quantities of some proteins and tiny amounts of others [1].

RNA molecules are single-stranded

Transcription is the first step a cell takes in reading out a needed part of its genetic instructions is to copy a particular portion of its DNA nucleotide sequence—a gene—into an RNA nucleotide sequence [1].

RNA structure is a linear polymer made of four different types of nucleotide subunits linked together by phosphodiester bonds. It differs from DNA chemically in two respects: it contains the sugar ribose and the base uracil (U) instead of thymine (T) [1].

RNA folding is single-stranded and can fold up into a particular shape, allowing some RNA molecules to have precise structural and catalytic functions [1].

Transcription produces RNA complementary to one strand of DNA

Transcription process begins with the opening and unwinding of a small portion of the DNA double helix to expose the bases on each DNA strand. One of the two strands of the DNA double helix then acts as a template for the synthesis of an RNA molecule [1].

RNA synthesis happens when nucleotide sequence of the RNA chain is determined by the complementary base-pairing between incoming nucleotides and the DNA template. The RNA chain is elongated one nucleotide at a time, and it has a nucleotide sequence that is exactly complementary to the strand of DNA used as the template [1].

Differs from DNA replication in several crucial ways. Unlike a newly formed DNA strand, the RNA strand does not remain hydrogen-bonded to the DNA template strand. Instead, the RNA chain is displaced and the DNA helix re-forms, releasing the RNA molecules as single strands [1].

RNA polymerase carry out transcription

RNA polymerases are enzymes that perform transcription. They catalyze the formation of the phosphodiester bonds that link the nucleotides together to form a linear chain [1].

Mechanism of RNA polymerase involves the RNA polymerase moving stepwise along the DNA, unwinding the DNA helix just ahead of the active site for polymerization to expose a new region of the template strand for complementary base-pairing, and extending the growing RNA chain by one nucleotide at a time in the 5'-to-3' direction [1].

Characteristics of RNA polymerase include the ability to start an RNA chain without a primer, being absolutely processive, and having a modest proofreading mechanism to correct errors in the growing RNA chain [1].

Comparison with DNA polymerase reveals that RNA polymerase is less accurate, with an error rate of about one mistake for every 10^4 nucleotides copied into RNA, despite catalyzing essentially the same chemical reaction as DNA polymerase [1].

RNA polymerase binds to DNA at specific sites

Types of RNA include messenger RNA (mRNA), which directs the synthesis of proteins, and noncoding RNAs, which do not code for protein and serve as enzymatic, structural, and regulatory components [1].

Functions of noncoding RNAs include directing the splicing of pre-mRNA to form mRNA, forming the core of ribosomes, and serving as adaptors that select amino acids and hold them in place on a ribosome for incorporation into protein [1].

Transcription units are segments of DNA that are transcribed into RNA. In eukaryotes, a transcription unit typically carries the information of just one gene, while in bacteria, a set of adjacent genes is often transcribed as a unit [1].

RNA composition in cells reveals that RNA makes up a few percent of a cell's dry weight, with most of the RNA being rRNA, and mRNA comprising only 3–5% of the total RNA in a typical mammalian cell [1].

Signals encoded in DNA tell RNA polymerase where to start and stop

Initiation of transcription is a crucial step in gene expression, where RNA polymerase recognizes where to start and stop transcribing a gene. The bacterial RNA polymerase core enzyme is a multisubunit complex that synthesizes RNA using the DNA template as a guide, with the help of the sigma (σ) factor [1].

During **transcription bubble formation**, the RNA polymerase holoenzyme opens up the double helix to expose a short stretch of nucleotides on each strand. The first ten or so nucleotides of RNA are synthesized using a "scrunching" mechanism, and the polymerase then moves down the DNA, synthesizing RNA in a stepwise fashion [1].

Chain elongation continues until the enzyme encounters the **Termination** signal, which consists of a string of A-T nucleotide pairs preceded by a twofold symmetric DNA sequence. The formation of a hairpin structure through Watson-Crick base-pairing helps to disengage the RNA transcript from the active site, marking the end of transcription [1].

Transcription Start and Stop Signals Are Heterogeneous in Nucleotide Sequence

Transcription initiation and termination involve a series of structural transitions in protein, DNA, and RNA molecules, and the signals encoded in DNA that specify these transitions are often difficult to recognize. Despite this, researchers have identified common features among bacterial promoters, which are often summarized in the form of a **consensus sequence**. A consensus sequence is derived by comparing many sequences with the same basic function and tallying up the most common nucleotides found at each position [1].

Promoter Strength and Sequence Variation The DNA sequences of individual bacterial promoters differ in ways that determine their strength, with promoters for genes that code for abundant proteins being much stronger than those associated with genes that encode rare proteins. The nucleotide sequences of their promoters are responsible for these differences [1].

Terminator Sequences Like bacterial promoters, transcription terminators also have a wide range of sequences, with the potential to form a simple hairpin RNA structure being the most important common feature. Terminator sequences are even more heterogeneous than promoter sequences [1].

Challenges in Locating Promoters and Terminators Despite our knowledge of bacterial promoters and terminators, their variation in nucleotide sequence makes it difficult to definitively locate them simply by analysis of the nucleotide sequence of a genome. Additional information, some of it from direct experimentation, is often needed to locate and accurately interpret the short DNA signals in genomes [1].

Promoter Orientation and Template Strand Promoter sequences are asymmetric, ensuring that RNA polymerase can bind in only one orientation. The promoter orientation specifies the strand to be used as a template, and genome sequences reveal that the DNA strand used as the template for RNA synthesis varies from gene to gene, depending on the orientation of the promoter [1].

Transcription Initiation in Eukaryotes Requires Many Proteins Eukaryotic nuclei have three types of RNA polymerase, with RNA polymerase II transcribing most genes, including those that encode proteins [1].

Key Differences from Bacterial RNA Polymerase

Eukaryotic RNA polymerase II differs from bacterial RNA polymerase in two main ways [1]:

- It requires many transcription-initiation factors, collectively called the general transcription factors.
- It must initiate transcription on DNA packaged into *nucleosomes* and higher-order chromatin structures.

RNA Polymerase II Requires a Set of General Transcription Factors

Eukaryotic RNA polymerase II requires a set of general transcription factors, denoted as TFIIA, TFIIB, TFIIC, TFIID, and so on, to position correctly at the promoter and initiate transcription [1].

Assembly of General Transcription Factors The assembly process begins when TFIID binds to the TATA sequence, a short double-helical DNA sequence primarily composed of T and A nucleotides. The binding of TFIID causes a large distortion in the DNA, which serves as a physical landmark for the location of an active promoter [1].

Role of TFIIF TFIIF, the most complicated of the general transcription factors, performs several enzymatic steps needed for the initiation of transcription, including unwinding the DNA and exposing the template strand [1].

Phosphorylation of RNA Polymerase II The phosphorylation of the C-terminal domain (CTD) of RNA polymerase II, specifically the serine located at the fifth position in the repeat sequence (Ser5), is a key step in the transition from initiation to elongation. This phosphorylation allows the polymerase to disengage from the cluster of general transcription factors and undergo conformational changes that tighten its interaction with DNA [1].

Release of General Transcription Factors Once the polymerase II has begun elongating the RNA transcript, most of the general transcription factors are released from the DNA, making them available to initiate another round of transcription with a new RNA polymerase molecule [1].

Polymerase II Also Requires Activator, Mediator, and Chromatin-Modifying Proteins

In addition to general transcription factors, RNA polymerase II requires several other proteins to initiate transcription in a eukaryotic cell, including [1]:

- **Transcriptional activators**, which bind to specific sequences in DNA (called enhancers) and help to attract RNA polymerase II to the start point of transcription.
- **Mediator**, a large protein complex that allows activator proteins to communicate properly with the polymerase II and with the general transcription factors.

Continues on the next page.

- **Chromatin-modifying enzymes**, including chromatin remodeling complexes and histone-modifying enzymes, which increase access to the DNA in chromatin and facilitate the assembly of the transcription initiation machinery onto DNA.

Assembly of Transcription Initiation Machinery The assembly of the transcription initiation machinery onto DNA involves the recruitment of many proteins (well over 100 individual subunits) and does not follow a prescribed pathway. The order of assembly differs from gene to gene, and some protein complexes may be brought to DNA as preformed subassemblies [1].

Release of RNA Polymerase II To begin transcribing, RNA polymerase II must be released from the large complex of proteins. This release often requires the in situ proteolysis of the activator protein [1].

Transcription Elongation in Eukaryotes Requires Accessory Proteins

Once RNA polymerase has initiated transcription, it requires accessory proteins to facilitate elongation. These proteins include [1]:

- **Elongation factors**, which decrease the likelihood that RNA polymerase will dissociate before it reaches the end of a gene.
- **ATP-dependent chromatin remodeling complexes**, which help RNA polymerase move through chromatin structure by either moving with the polymerase or rescuing stalled polymerases.
- **Histone chaperones**, which partially disassemble nucleosomes in front of a moving RNA polymerase and assemble them behind.

Modification of Histones During Transcription Elongation As RNA polymerase moves along a gene, enzymes bound to it modify the histones, leaving behind a record of where the polymerase has been. This information may aid in transcribing a gene over and over again once it has become active for the first time, and may also be used to coordinate transcription elongation with RNA processing [1].

Transcription Creates Superhelical Tension

DNA Supercoiling is a conformation that DNA adopts in response to superhelical tension, which can be created by opening or unwinding a double-helical DNA molecule [1].

Effect of RNA Polymerase on DNA Supercoiling is to create superhelical tension as it moves along a stretch of DNA that is anchored at its ends, generating positive superhelical tension in the DNA in front of it and negative helical tension behind it [1].

Consequences of DNA Supercoiling include making the DNA helix more difficult to open, but also facilitating the partial unwrapping of the DNA in nucleosomes in eukaryotes [1].

Removal of Superhelical Tension is achieved by DNA topoisomerase enzymes in eukaryotes, while in bacteria, DNA gyrase uses the energy of ATP hydrolysis to pump supercoils continuously into the DNA, maintaining the DNA under constant tension [1].

Transcription Elongation in Eukaryotes Is Tightly Coupled to RNA Processing

Eukaryotic mRNA Synthesis involves several steps beyond transcription, including covalent modification of the ends of the RNA and removal of intron sequences by RNA splicing [1].

Modification of Eukaryotic mRNAs includes capping on the 5' end and polyadenylation of the 3' end, allowing the cell to assess the integrity of the mRNA molecule before exporting it from the nucleus and translating it into protein [1].

RNA Splicing joins together the different portions of a protein-coding sequence, enabling eukaryotes to synthesize several different proteins from the same gene [1].

Coupling of RNA Processing to Transcription Elongation is achieved through the phosphorylation of the RNA polymerase II tail, which allows a new set of proteins to associate with the RNA polymerase and function in transcription elongation and RNA processing [1].

RNA Capping Is the First Modification of Eukaryotic Pre-mRNAs

5' Capping of Eukaryotic mRNAs involves the addition of a modified guanine nucleotide to the 5' end of the new RNA molecule, which is performed by three enzymes acting in succession [1].

Mechanism of 5' Capping includes the removal of a phosphate from the 5' end of the nascent RNA, the addition of a GMP in a reverse linkage, and the addition of a methyl group to the guanosine [1].

Function of the 5'-Methyl Cap is to signify the 5' end of eukaryotic mRNAs, helping the cell to distinguish mRNAs from other types of RNA molecules, and to facilitate further processing and export of the mRNA [1].

Binding of the 5'-Methyl Cap to a protein complex called CBC (cap-binding complex) in the nucleus, which helps the mRNA to be further processed and exported [1].

RNA Splicing Removes Intron Sequences from Newly Transcribed Pre-mRNAs

Eukaryotic Gene Structure is characterized by the presence of noncoding intervening sequences (introns) that interrupt the protein-coding sequences (exons) [1].

RNA Splicing is the process by which intron sequences are removed from the newly synthesized RNA, and the exons are joined together through two sequential phosphoryl-transfer reactions known as transesterifications [1].

Mechanism of RNA Splicing involves a complex machinery consisting of five additional RNA molecules and several hundred proteins, which hydrolyzes many ATP molecules per splicing event to ensure accurate and flexible splicing [1].

Advantages of RNA Splicing include facilitating the emergence of new and useful proteins over evolutionary time scales by allowing genetic recombination to combine the exons of different genes, and enabling eukaryotes to increase the coding potential of their genomes by producing a corresponding set of different proteins from the same gene through alternative splicing [1].

Nucleotide Sequences Signal Where Splicing Occurs

Recognition of Splicing Sites requires the splicing machinery to identify three portions of the precursor RNA molecule: the 5' splice site, the 3' splice site, and the branch point in the intron sequence [1].

Consensus Sequences for Splicing are short nucleotide sequences that are similar from intron to intron and provide cues for where splicing is to take place, but can accommodate extensive sequence variability [1].

Challenges in Deciphering Genome Sequences arise from the high variability of splicing consensus sequences and the possibility of alternative splicing, making it difficult to predict protein sequences solely from a genome sequence and identify all of the genes in a complete genome sequence [1].

Nucleotide Sequences Signal Where Splicing Occurs

Recognition of Splicing Sites requires the splicing machinery to identify three portions of the precursor RNA molecule: the 5' splice site, the 3' splice site, and the branch point in the intron sequence [1].

Consensus Sequences for Splicing are short nucleotide sequences that are similar from intron to intron and provide cues for where splicing is to take place, but can accommodate extensive sequence variability [1].

Challenges in Deciphering Genome Sequences arise from the high variability of splicing consensus sequences and the possibility of alternative splicing, making it difficult to predict protein sequences solely from a genome sequence and identify all of the genes in a complete genome sequence [1].

RNA Splicing Is Performed by the Spliceosome

Role of RNA Molecules in RNA Splicing involves specialized RNA molecules, known as snRNAs (small nuclear RNAs), that recognize the nucleotide sequences that specify where splicing is to occur and catalyze the chemistry of splicing [1].

Composition of the Spliceosome includes five snRNAs (U1, U2, U4, U5, and U6) complexed with at least seven protein subunits to form an snRNP (small nuclear ribonucleoprotein), which forms the core of the spliceosome [1].

Mechanism of Spliceosome Assembly involves the recognition of the 5' splice junction, the branch-point site, and the 3' splice junction through base-pairing between the snRNAs and the consensus RNA sequences in the pre-mRNA substrate, and the dynamic assembly and rearrangement of the spliceosome components during the splicing reaction [1].

Spliceosome Structure and Function is complex and dynamic, with some scientists believing that the spliceosome is a preexisting, loose assembly of all the components that captures, splices, and releases RNA as a coordinated unit, undergoing extensive rearrangements each time a splice is made [1].

The Spliceosome Uses ATP Hydrolysis to Produce a Complex Series of RNA-RNA Rearrangements

Energy Requirements for RNA Splicing reveals that ATP hydrolysis is not required for the chemistry of RNA splicing itself, but is necessary for the assembly and rearrangements of the spliceosome [1].

Role of ATP Hydrolysis in Spliceosome Assembly involves the use of energy from ATP hydrolysis to break existing RNA-RNA interactions and allow the formation of new ones, enabling the splicing signals on the pre-RNA to be examined by snRNPs several times during the course of splicing [1].

Purpose of Spliceosome Rearrangements is to increase the overall accuracy of splicing by allowing the spliceosomes to check and recheck the splicing signals, and to create the active sites for the two transesterification reactions [1].

Catalytic Sites of the Spliceosome are formed by both protein and RNA molecules, with the RNA molecules catalyzing the actual chemistry of splicing [1].

Disassembly of the Spliceosome requires another series of RNA-RNA rearrangements that require ATP hydrolysis, returning the snRNAs to their original configuration so that they can be used again in a new reaction [1].

Exon Junction Complex (EJC) is a set of proteins that bind to the mRNA near the position formerly occupied by the intron, marking the site of a successful splicing event and influencing the subsequent fate of the mRNA [1].

RNA Can Be Processed by Dicer to Produce siRNAs

Energy Requirements for RNA Splicing reveals that ATP hydrolysis is not required for the chemistry of RNA splicing itself, but is necessary for the assembly and rearrangements of the spliceosome [1].

Role of ATP Hydrolysis in Spliceosome Assembly involves the use of energy from ATP hydrolysis to break existing RNA-RNA interactions and allow the formation of new ones, enabling the splicing signals on the pre-RNA to be examined by snRNPs several times during the course of splicing [1].

Purpose of Spliceosome Rearrangements is to increase the overall accuracy of splicing by allowing the spliceosomes to check and recheck the splicing signals, and to create the active sites for the two transesterification reactions [1].

Catalytic Sites of the Spliceosome are formed by both protein and RNA molecules, with the RNA molecules catalyzing the actual chemistry of splicing [1].

Disassembly of the Spliceosome requires another series of RNA-RNA rearrangements that require ATP hydrolysis, returning the snRNAs to their original configuration so that they can be used again in a new reaction [1].

Exon Junction Complex (EJC) is a set of proteins that bind to the mRNA near the position formerly occupied by the intron, marking the site of a successful splicing event and influencing the subsequent fate of the mRNA [1].

Other Properties of Pre-mRNA and Its Synthesis Help to Explain the Choice of Proper Splice Sites

Challenges in Splice-Site Selection arise from the enormous size variation of intron sequences, which could lead to frequent splicing mistakes if not addressed by fidelity mechanisms [1].

Strategies to Increase Splicing Accuracy include coupling splicing to transcription, which helps the cell keep track of introns and exons, and "exon definition", which involves marking off exon sequences with SR proteins to increase the accuracy of splicing component deposition [1].

Mechanism of Exon Definition involves the assembly of SR proteins on exon sequences, which recruit U1 and U2 snRNAs to mark the downstream and upstream exon boundaries, respectively, and increase the accuracy of splicing.

Role of SR Proteins is to bind preferentially to specific RNA sequences in exons, termed splicing enhancers, and help to discriminate exon sequences from intron sequences [1].

Timing of Splicing Events reveals that the marking of exon and intron boundaries and the assembly of the spliceosome begin on an RNA molecule while it is still being elongated by RNA polymerase, but the actual chemistry of splicing can take place later, allowing for the removal of intron sequences in a non-linear order [1].

Chromatin Structure Affects RNA Splicing

Influence of Chromatin Structure on RNA Splicing reveals that the way a gene is packaged into chromatin can affect how the RNA transcript of that gene is ultimately spliced [1].

Nucleosome Positioning and RNA Splicing suggests that nucleosomes tend to be positioned over exons, acting as "speed bumps" that allow proteins responsible for exon definition to assemble on the RNA as it emerges from the polymerase [1].

Effect of Chromatin Structure on Transcription Rate shows that changes in chromatin structure can alter splicing patterns by affecting the rate at which RNA polymerase moves along DNA [1].

Direct Influence of Chromatin Structure on RNA Splicing involves specific histone modifications attracting components of the spliceosome, which can be transferred to the emerging RNA, affecting the final pattern of splicing [1].

RNA Splicing Shows Remarkable Plasticity

Factors Influencing Splice Site Choice include the strength of the three signals on the RNA (the 5' and 3' splice junctions and the branch point), co-transcriptional assembly of the spliceosome, chromatin structure, and exon definition [1].

Flexibility of Splicing allows for the splicing machinery to pick out the best possible pattern of splice junctions, even if the optimal one is damaged by mutation, and can lead to changes in splicing patterns caused by random mutations [1].

Consequences of Mutations Affecting Splicing can be severely detrimental to the organism, and it is estimated that 10% of point mutations that cause inherited human diseases produce aberrant splicing of the gene containing the mutation [1].

Regulation of RNA Splicing allows the cell to easily regulate the pattern of RNA splicing, and alternative splicing can give rise to different proteins from the same gene, enhancing the coding potential of genomes [1].

Regulated RNA Splicing involves the cell regulating the splicing patterns so that different forms of the protein are produced at different times and in different tissues, and is a common strategy to enhance the coding potential of genomes [1].

Spliceosome-Catalyzed RNA Splicing Probably Evolved from Self-splicing Mechanisms

Evolution of the Spliceosome suggests that RNA molecules were used as major catalysts in early cells, and RNA-catalyzed splicing reactions played critical roles in these cells [1].

Evidence for RNA-Catalyzed Splicing includes the existence of self-splicing RNA introns, which can catalyze their own splicing in the absence of proteins or other RNA molecules [1].

Mechanism of Self-Splicing Introns involves the RNA molecule folding into a specific three-dimensional structure that brings the intron/exon junctions together and catalyzes the two transesterification reactions [1].

Evolution of Pre-mRNA Splicing is thought to have evolved from a simpler, ancestral form of RNA self-splicing, with the basic chemistry of some self-splicing reactions being similar to pre-mRNA splicing [1].

RNA-Processing Enzymes Generate the 3' End of Eukaryotic mRNAs

Processing of the 3' End of Eukaryotic mRNAs involves the recognition of signals encoded in the genome by RNA-binding proteins and RNA-processing enzymes [1].

Role of CstF and CPSF is to bind to their recognition sequences on the emerging RNA molecule and assemble with additional proteins to create the 3' end of the mRNA [1].

Mechanism of 3'-End Processing involves the cleavage of the RNA from the polymerase, followed by the addition of a poly-A tail by poly-A polymerase, and the assembly of poly-A-binding proteins onto the tail [1].

Synthesis of the Poly-A Tail is a non-template-dependent process, where poly-A polymerase adds approximately 200 A nucleotides to the 3' end of the mRNA, using ATP as the nucleotide precursor [1].

Termination of Transcription is thought to be caused by the continued degradation of the newly synthesized RNA by a 5' → 3' exonuclease carried along on the polymerase tail, which eventually causes the RNA polymerase to release its grip on the template[1].

Bibliography

- [1] B. Alberts. *Molecular Biology of the Cell*. W.W. Norton, 2017. ISBN: 9781317563747. URL: <https://books.google.dk/books?id=2xIwDwAAQBAJ>.
- [2] Jyoti P. Tamang, Koichi Watanabe, and Wilhelm H. Holzapfel. "Review: Diversity of Microorganisms in Global Fermented Foods and Beverages". eng. In: *Frontiers in microbiology* 7 (2016), pp. 377–377. ISSN: 1664-302X.
- [3] Benjamin E. Wolfe and Rachel J. Dutton. "Fermented Foods as Experimentally Tractable Microbial Ecosystems". In: *Cell* 161.1 (2015), pp. 49–55. ISSN: 0092-8674. DOI: <https://doi.org/10.1016/j.cell.2015.02.034>. URL: <https://www.sciencedirect.com/science/article/pii/S0092867415002007>.

Chapter A
Appendix