



Molecular and Functional Properties of Milk

Exam Question 09: Increased Protein Degradation

Lucas Daniel Paz Zuleta, TZS159

MSc student at the University of Copenhagen

Last compiled: 27-03-2025

Link to Git repo.: https://github.com/DanishUnicorn/fp_exam_2025



Question 09

Situation

You are the manager of a medium-sized Danish co-operative dairy plant in Northern Jutland and suspect an increased protein degradation occurring in your milk.

Your task

Map out the reasons for this and determine which methods to use to prove that this is happening.

Advice

Explain the possible reasons for protein degradation occurring in milk and which proteins are susceptible, and the consequences for this proteolysis in selected products. List the methods which you would use to determine this and what actions you would take to prevent it occurring in the dairy plant and at the farm level.

Contents

1	Introduction	1
2	Milk Composition	1
3	Protein Degradation	1
3.1	Reasons for Protein Degradation	1
3.2	Susceptible Proteins	2
4	Consequences of Proteolysis in Milk Products	2
5	Methods for Analysis	3
5.1	Determining Protein Degradation	3
5.2	Preventing Protein Degradation	3
5.2.1	Dairy Plant Level	3
5.2.2	Farm Level	3
6	Conclusion	3

Synopsis for Question 09

1 Introduction

Milk is an important nutritional source for humans, and serves as the basis for a variety of dairy products. As the main ingredient in dairy products, the microbiological and compositional quality of the raw milk is of great importance and is directly correlated with the quality of the final dairy product [1].

Proteins play an essential role in determining texture, flavour, and functional properties. If the proteins present in milk are degraded, either enzymatically or due to microbial activity, product quality can suffer significantly [2]. Upholding a high standard for milk is therefore not only of importance with respect to consumer acceptance, but also of economic relevance [1].

2 Milk Composition

Milk is a complex liquid whose composition includes a variety of dilute salts, the simple sugar, mainly lactose, and vitamins where fat is emulsified as globules [3]. Proteins in milk is mostly present in the form of casein micelles, which are colloidal aggregates of thousands of molecules [4].

Though the protein content (approximately 3.2% [5]) of milk is relatively low, compared to that of fat globules [4], this synopsis will focus on proteins and their degradation in milk.

The milk protein content and composition is influenced by various factors, such as breed, lactation stage, genetic variants, and cell count [6]. Furthermore, the protein content of milk is divided into two major groups: caseins and whey proteins. Caseins make up approximately 80% of the total protein content, while whey proteins make up the remaining 20% [6]. The composition of milk casein proteins consists of α_{s1} -casein, α_{s2} -casein, β -casein, and κ -casein, in respective order [6]. The casein proteins are present in structures of micelles and are relatively hydrophobic fibrous proteins [4]. Of the existing whey proteins in milk, the major constituents are α -lactalbumin, β -lactoglobulin, bovine serum albumin, immunoglobulins, and enzymes, in respective order [6].

3 Protein Degradation

While the somatic cell count (SCC) in milk is an important indicator of udder health, it is used to monitor the presence of mastitis in dairy cows [7]. Mastitis is an inflammation of the mammary gland, and is a major problem for dairy producers, as an increase in SCC corresponds to an increase in the proteolytic potential in milk [2]. This protein degradation in milk can have negative effects on yield and quality of dairy products, such as cheese [2].

3.1 Reasons for Protein Degradation

There are various reasons for the degradation of proteins in milk, e.g. microbial proteases, endogenous proteases, and heat treatment [2].

Microbial Proteases Some psychrotrophic bacteria can survive pasteurization, as they can produce heat-stable proteases [8]. These proteases will be active, even after heat treatment, and can initiate the degradation of the

casein- and whey proteins in milk by proteolysis [9]. This is predominantly the case for psychrotrophic gram-negative bacteria [8]. If *Pseudomonas fluorescens* is present, normal pasteurization will be insufficient and the bacteria's extracellular proteinases will degrade the proteins by proteolysis [8].

Endogenous Proteases There are many enzymes with milk at its natural habitat, such as plasmin, cathepsin D, and cathepsin B, which can degrade the proteins in milk [2]. Plasmin is the most important and predominant protease in milk, and is largely responsible for the degradation of casein proteins. The quantity of plasmins is correlated with mastitis, analysis of the raw milk is therefore imperative [2].

Heat Treatment Though the primary objective of heat treatments as pasteurization and ultra high temperature pasteurization (UHT-pasteurization) is to kill pathogenic and spoilage bacteria, the temperature does not differentiate by microorganisms, it kills/inactivates all microorganisms which is not sufficiently heat resistant [9].

3.2 Susceptible Proteins

Microbial Proteases The proteins most susceptible to microbial derived enzymes, resulting in proteolysis are the caseins, as their high proline content makes them more susceptible to proteolysis [10].

Endogenous Proteases The most susceptible proteins to endogenous proteases are the caseins α – $casein_{s1}$, β -casein and the whey protein α -lactalbumin. The decrease for these three proteins are percentagely the same (26-75%), but the content of the caseins is approximately 10 times higher, therefore, the overall quantitative loss is greatest for caseins [7].

Heat treatment Caseins have a high content of proline, for α_{s1} -, α_{s2} -, β -, and κ -casein, the proline content is 17, 10, 35, and 20 residues per mole, respectively [10]. The high proline content in the caseins results in a low content of α -helix or β -sheet structures makes them structurally more prone to denaturation and aggregation under heat treatment [10].

4 Consequences of Proteolysis in Milk Products

Severely heat treating milk can result in degradation of proteins in milk which in turn can have negative effects on the quality of the final dairy product, e.g. cheese not ripening properly due to high moisture and excessive syneresis in yogurt [9]. Another consequence of heat treating milk is maillard reaction which can result in a brownish color and a cooked taste [11]. Enzymatic activity from plasmin has also been shown to have a direct impact on both bitterness, gellation and shortens shelf-life [11]. A combination of high enzymatic activity, resulting in more available amino acids, and high temperature treatments can result in a more pronounced maillard reaction [11]. If there is a reduced amount of casein proteins in milk, leading to reduced cheese yield, since caseins are the primary proteins responsible for curd formation [2]. The reduction in casein proteins will also affect the cheese ripening process, as the endogeneous proteinases contribute to the cheese ripening process through casein hydrolysis [2].

5 Methods for Analysis

5.1 Determining Protein Degradation

5.2 Preventing Protein Degradation

5.2.1 Dairy Plant Level

5.2.2 Farm Level

6 Conclusion

Bibliography

- [1] Maria Åkerstedt et al. “Protein degradation in bovine milk caused by *Streptococcus agalactiae*”. In: *Journal of Dairy Research* 79.3 (2012), pp. 297–303. DOI: [10.1017/S0022029912000301](https://doi.org/10.1017/S0022029912000301).
- [2] Lotte B Larsen et al. “Proteases and protein degradation in milk from cows infected with *Streptococcus uberis*”. In: *International Dairy Journal* 14.10 (2004), pp. 899–907. ISSN: 0958-6946. DOI: <https://doi.org/10.1016/j.idairyj.2004.03.006>. URL: <https://www.sciencedirect.com/science/article/pii/S0958694604000792>.
- [3] Anna Haug, Arne T. Høstmark, and Odd M. Harstad. “Bovine milk in human nutrition – a review”. In: *Lipids in Health and Disease* 6.25 (2007). DOI: [10.1186/1476-511X-6-25](https://doi.org/10.1186/1476-511X-6-25). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC2039733/>.
- [4] A.L. Kelly and L. Bach Larsen. “1 - Milk biochemistry”. In: *Improving the Safety and Quality of Milk*. Ed. by Mansel W. Griffiths. Woodhead Publishing Series in Food Science, Technology and Nutrition. Woodhead Publishing, 2010, pp. 3–26. ISBN: 978-1-84569-438-8. DOI: <https://doi.org/10.1533/9781845699420.1.3>. URL: <https://www.sciencedirect.com/science/article/pii/B9781845694388500015>.
- [5] Morten Dam Rasmussen. *Milk Secretion and Ejection*. Lecture slides from the course Molecular and Functional Properties of Milk, University of Copenhagen. 2025. URL: <https://absalon.ku.dk/courses/80667/files/folder/Lecture%5C%20Slides>.
- [6] Lotte Bach Larsen. *Protein Fraction in Milk – Factors Affecting Composition and Quality*. Lecture slides from the course Molecular and Functional Properties of Milk, University of Copenhagen. 2025. URL: <https://absalon.ku.dk/courses/80667/files/folder/Lecture%5C%20Slides>.
- [7] K. Krogh. *The mastitis complex – Causes and possibilities for prevention*. Lecture slides from the course Molecular and Functional Properties of Milk, University of Copenhagen. 2025. URL: <https://absalon.ku.dk/courses/80667/files/folder/Lecture%5C%20Slides>.
- [8] Valentin Rauh. *Heat Treatment 1 – Introduction and Categories*. Lecture slides from the course Molecular and Functional Properties of Milk, University of Copenhagen. 2025. URL: <https://absalon.ku.dk/courses/80667/files/folder/Lecture%5C%20Slides>.
- [9] P. F. Fox et al. “Chapter 9: Heat-Induced Changes in Milk”. In: *Dairy Chemistry and Biochemistry*. Cham: Springer International Publishing, 2015, pp. 345–375. ISBN: 978-3-319-14892-2. DOI: [10.1007/978-3-319-14892-2_9](https://doi.org/10.1007/978-3-319-14892-2_9). URL: https://doi.org/10.1007/978-3-319-14892-2_9.
- [10] P. F. Fox et al. “Chapter 4: Milk Proteins”. In: *Dairy Chemistry and Biochemistry*. Switzerland: Springer International Publishing, 2015, pp. 141–186. DOI: [10.1007/978-3-319-14892-2_4](https://doi.org/10.1007/978-3-319-14892-2_4).
- [11] Valentin Rauh and Yinghua Xiao. “The shelf life of heat-treated dairy products”. In: *International Dairy Journal* 125 (2022), p. 105235. DOI: [10.1016/j.idairyj.2021.105235](https://doi.org/10.1016/j.idairyj.2021.105235). URL: <https://doi.org/10.1016/j.idairyj.2021.105235>.