



Fruit and Berry Crop Physiology and Quality

NPLK14014U

Notes taken during the course, including lectures, exercises, curriculum, and practicals

Lucas Daniel Paz Zuleta, TZS159

MSc student at the University of Copenhagen

Last compiled: 01-11-2025

[Link to GitHub repository](#)



Preface

These course notes have been prepared as part of the NPLK14014U course Fruit and Berry Crop Physiology and Quality at the University of Copenhagen, covering the period from September to November 2025.

The notes compile material and reflections relevant to the course and are intended as a resource to enhance the learning experience for students. The content is shared freely and may be used as study material or as a template for structuring individual notes.

All information is provided without responsibility for its correctness, and users are encouraged to verify data, formulas, and interpretations with the original sources and course materials.

Please enjoy reading these notes, and feel free to reach out if you have any questions.

Course Description

Education

MSc Programme in Agriculture

Content

The focus is on fruit growth and fruit quality in relation to the use as fresh fruits or for processing. How is fruit growth and quality affected by the plants' physiological and genetic basis and how can it be influenced by different growing techniques and environmental factors? Similarities and differences among the fruit crop types (pit fruits, stone fruits, berries and nuts), with regard to demands in growing conditions are discussed. Furthermore, we analyze which physiological parameters are important in the different fruit species for determining yield and important quality components. Emphasis is on temperate fruits, nuts, berries and fruit vegetables, grown mainly in open field or in tunnel systems. The reference growing systems are the common commercial systems, including organic growing. The course also addresses examples of the genetic and quality variation among cultivars and the importance of different quality attributes in relation to postharvest use (fresh consumption, cooking, juice processing or fruit wine making). In general the crop specific aspects of the following main topics will be covered:

- Yield and quality components (organ development and interactions) and determinant factors
- Allocation of dry matter and nutrient among sources and sinks in fruiting plants
- Control of vigour and plant structure by pruning and management of nutrients and irrigation
- Effects of preharvest factors (climate, a-biotic or biotic stresses) on internal and external quality of fruits
- Content and development of secondary and bioactive compounds in fruits.
- Maturation, ripening and assessment of optimal harvest and quality aspects of fruits and berries.
- Post harvest usability and sensory aspects of different cultivars and fruit types.

In addition to fresh use, special attention is given to production and quality of fruit juices. Biotechnological aspects are addressed at a limited level.

Learning Outcome

The course is targeted to students interested in plant science (Horticulture and Agriculture) and food science students who are particularly interested in fruit and berry crops and the quality and use of the raw materials/food products these crops provide.

Knowledge

- The physiological basis for production of fruiting crops (including fruit vegetables such as tomato and cucumber).
- Overview of development of the major plant organs with focus on the fruit and its quality and understand how and why it varies with genotype and preharvest growing conditions
- Describe the variation among the major cultivars used of fruits and berries in terms of development and quality parameters.
- Reflect on the importance of fruit and berries for human health

Skills

- Apply basic knowledge of physiology and biochemistry from plant and food science at the whole plant and organ level.
- Analyse a fruiting crop based on the crop specific yield and quality components.
- Explain how and why different techniques are used in the fruit industry and how it affects plant growth and product/fruit quality.

Competences

- Analyse the methods used to obtain optimal productivity and product quality.
- Discuss trade offs in management, such as between optimal sensory quality and storability, between yield and quality or pesticide use vs organic growing

Literature

Literature lists will be available from the course responsible.

Recommended Academic Qualifications

Academic qualifications equivalent to a BSc degree is recommended.

Teaching and Learning Methods

Besides lectures the course will include practicals, where the students are working with cultivar evaluation, quality analysis or aspects of fruit growing physiology (plant and organ development etc). Part of the hands on teaching will be field based in the experimental fruit collections at the Pomatum.

The practicals will be made in groups, while the individual student is given the opportunity, in a major report written throughout the course, to focus on an area of special interests. Thus individual competences with emphasis on either fruit growing physiology or fruit quality aspects of fruits as raw materials for industry processing or fresh consumption can be developed. The topic of the major report are to be presented to the class in a short lecture based on a selected journal paper.

2 or 3 excursions will be arranged in connection with the different course subjects.

Workload

Table 1: A table with an overview over the workload for the course.

Category	Hours
Lectures	35
Class Instruction	5
Preparation	40
Practical exercises	30
Excursions	21
Project work	75
Total	206

Exam

Table 2: A table with an overview over the elaborated description of the course

Credit	7.5 ECTS
Type of assessment	<ul style="list-style-type: none">• Oral examination, 20 minuter• Written assignment, ca. 3 uger
Type of assessment details	The portfolio includes a major report and 2 out of 4 additional products (e.g. exercise reports or presentation) Weight of exam components: Evaluation of major report 50 %, oral examination in portfolio contents and curriculum 50%.
Examination prerequisites	Submitted and approval of the reports for theoretical and practical exercises
Aid	All aids allowed Read about how to use Generative AI on KuNet
Marking scale	7-point grading scale
Censorship form	<ul style="list-style-type: none">• No external censorship• One internal examiner
Re-exam	The exam is an oral exam, as for the ordinary exam. Submission of an individual major report 1 week before the oral re-exam is required. The topic may be as for the ordinary exam but in a revised version.

Contents

1	Lecture Notes	1
1	Introduction	1
2	Lecture 01: Introduction - 01/09-2025	1
3	Lecture 02: Yield and quality components - 01/09-2025	2
4	Lecture 03: Bud and shoot development - 01/09-2025	3
5	Lecture 04: Flowers and flowerbud development - 05/09-2025	4
6	Lecture 05: Bud dormancy - 05/09-2025	5
7	Lecture 06: Flowering, pollination and fruit set - 08/09-2025	6
8	Lecture 07: Fruit Thinning - 08/09-2025	7
9	Lecture 08+09: Source-sink and carbon allocation - 08/09-2025	8
10	Lecture 10: Table Grapes - growing techniques and cultivars - 12/09-2025	9
11	Lecture 11+12: Fruit development of large fruited spe- cies - 15/09-2025	10
12	Lecture 13+14: Fruit development of small fruited spe- cies / berries - 15/09-2025	11
13	Lecture 15+16: Aromas in fruits - 22/09-2025	12
14	Lecture 17+18: Organic growing of fruit and berries 22/09-2025	13
15	Lecture 19: Light utilisation and canopy management 26/09-2025	14
16	Lecture 20: Canopy management/General techniques - 26/09-2025	15
17	Lecture 21: Juice processing and Quality - 29/09-2025	16
18	Lecture 22: Canopy management/ special techniques - 29/09-2025	18
19	Lecture 24: Nutrients, use and effects on development - 06/10-2025	18
20	Lecture 25: Nutrients use and effects on quality - 06/10-2025	19
21	Lecture 23: Stresses and effects on quality - 10/10-2025	20
22	Lecture 28: Potentials in fruit and berries for fruit wine production in Denmark - 10/10-2025	21
23	Lecture 30: Growing of wild berries - 20/10-2025	22
24	Lecture 31: Bioactive compounds in fruit and berries - 20/10-2025	23
25	Lecture 32: Water management and water stress - 24/10-2025	24
26	Lecture 33: Vitamins and colours in fruit and berries - 24/10-2025	26
27	Lecture 34: Fruit harvest and postharvest - 27/10-2025	27
2	Literature résumés	29
1	RM for L02 - Fruit Yield and Quality. Components and Determinants	29
2	RM for L03 - Buds and Bud Development	30
3	RM for L04 - Chapter 2: Flower bud formation, flower development and flowering	31
4	RM for L05 - A model to predict the beginning of the pollen season	33
5	RM for L06 - Physiology of Temperate Zone Fruit Trees	34
6	1 st RM for L07 - Physiology of Temperate Zone Fruit Trees	35

7	2 nd RM for L07 - Manipulation of growth and development by plant bioregulators	36
8	RM for L08+09 - Chapter 5 Growth, Growth correlations and Assimilate turnover	37
9	RM for L10 - Table Grapes in Cold Houses	39
10	1 st RM for L11-13 - CHAPTER 6 Fruit growth and fruit quality	40
11	2 nd RM for L11-13 - CHAPTER 7 Fruit Development and fruit ripening	41
12	RM for L13 - Growth and development in black currant (<i>Ribes nigrum</i>)	42
13	1 st RM for L15-16 - Advances in Fruit Aroma Volatile Research	43
14	2 nd RM for L15-16 - Advances in Fruit Aroma Volatile Research	44
15	3 rd RM for L15-16 - Advances in Fruit Aroma Volatile Research	45
16	1 st RM for L17+18 - Sustainable production systems for organic apple production	45
17	2 nd RM for L17+18 - Fruit quality in organic and conventional farming: advantages and limitations	46
18	RM for L19 - 9 Light Relations	47
3	Exam Questions and Answers	49
1	Yield and quality determinants and components	49
1.1	Characterise the development and importance of spurs and extension (long) shoots	50
1.2	Describe differences in bud development and structure between stone and pome fruits . .	50
1.3	Describe some important yield components in strawberry and in sour cherry	50
1.4	Describe some conditions which may affect the development of flower buds negatively . .	50
2	Yield and quality determinants and components	50
2.1	Describe important factors determining fruit set?	50
2.2	What is the importance of EPP?	50
2.3	What are important quality parameters for pollen and flowers?	50
2.4	Why and how do we use pollinators?	50
2.5	Are insects (fx bees) needed in pollination of self-pollinating crops?	50
3	Fruit development	50
3.1	Describe the general developmental phases in fruit development	51
3.2	Which sugars and acids are important in fruit development and how do they develop during fruit development? Example of species differences.	51
3.3	Which sugars are transported in the plant?	51
3.4	What is the role of starch in the carbon balance of an apple tree and an apple fruit?	51
4	Light use, vigor control and canopy management	51
4.1	Why do we manipulate the canopy structure in most fruit crops?	51
4.2	Describe the pruning response during the year. Why do we get differences in the growth response to pruning?	51
4.3	How does pruning affect fruit development and quality? (direct and indirect)	51
4.4	Characterise important factors (except from time in the year), which may influence the growth response to pruning?	51
5	Crop load and canopy management	51
5.1	How does a high fruit load influence photosynthesis and transpiration?	52
5.2	Explain the concept of source strength and sink strength	52
5.3	How do source-sink relationships develop during the season in an apple tree?	52
5.4	Why may some leaves be more important than others for fruit development?	52
5.5	Why do premature fruit drop occur?	52
6	Crop load management, fruit quality and vigor control	52

6.1	Give an example of a crop in which crop load has a strong impact on fruit development - and one where it does not.	52
6.2	Characterize the effects of fruit thinning on growth and development	52
6.3	When is it most optimal to perform fruit thinning? Why?	52
6.4	Explain why the optimal thinning strategy may dependent on the end use of the fruits. . .	52
6.5	Why do we not want fruits on a young tree the first year(s) after planting?	52
7	Preharvest factor management and quality	52
7.1	Characterise the differences in nutrient requirements of a vegetative growing and a fruiting plant?	53
7.2	Calcium is important for fruit quality. Why? - And why is the level of calcium low in many fruits, especially big fruits?	53
7.3	When and why are fertilizers often sprayed on the leaves and fruits in the production of apples?	53
7.4	Characterize the importance of potassium for fruit development	53
8	Preharvest factor management and quality	53
8.1	Describe the effects of nitrogen status on plant development	53
8.2	In which ways do nitrogen levels influence the yield components?	53
8.3	Impacts of nitrogen levels on fruit quality?	53
9	Preharvest factor management and quality	53
9.1	Describe the effects of stresses of nutrients and water on fruit development and quality. . .	53
9.2	Why are deficiency symptoms by some nutrients seen in the young leaves and by others in the old?	53
9.3	Describe how water stress can be used as a tool for growth control.	53
10	Fruit development	53
10.1	Describe some important factors for optimal fruit development in small and large fruited species. Are there differences?	54
10.2	What would you do to optimize fruit development and fruit quality in an apple crop? . . .	54
10.3	What is important for fruit development and quality in raspberry and strawberry?	54
11	Fruit maturity, harvest and quality assessment	54
11.1	How would you determine the optimal harvest time in apple?	54
11.2	Describe the problems and quality effects you might get, if you harvest either too early or too late.	54
11.3	Hand picking vs mechanical harvest - problems and benefits?	54
11.4	What are the main reasons for post harvest losses and what may be done to minimize it? .	54
12	Fruit maturity, cultivar variations and important quality parameters	54
12.1	When do aromas develop in fruits?	55
12.2	Characterise some important aroma substances and changes in aroma with maturity	55
12.3	Characterize the importance of harvest time on aroma development	55
12.4	What might affect aroma development pre and post harvest?	55
13	Fruit maturity, cultivar variations and important quality parameters	55
13.1	Characterise some important colour substances in fruits and berries	55
13.2	How does colour change with maturity?	55
13.3	What might affect colour development pre and post harvest?	55
13.4	What is the mechanism behind the occurrence of red clones in fruit cultivars (fx apples, pears and grapes)?	55

14	Cultivar variations and important quality parameters (fresh use and juice)	55
14.1	Characterise some of the most important (internal and external) quality characters, which may vary among cultivars in a fruit crop. (Fx strawberries or apple)	56
14.2	Which compounds are considered especially important in fruit and berries for human health and where are they located?	56
14.3	Which species are believed to be especially healthy to eat? Comment on the consumption of raw or processed fruits and berries.	56
15	Cultivar variations and important quality parameters (fresh use and juice)	56
15.1	How does the level of fruit ripening impact on juice processing and juice quality?	56
15.2	Which enzymes may be used in juice processing and why?	56
15.3	Comment on the effects of different juice processing steps on juice quality.	56
15.4	Why are juices pasteurised, and what are important factors for a successful pasteurisation?	56
16	Potentials for producing fruit and berry wines	56
16.1	Comment on the challenges and potentials in making fruit wine from different fruit and berries	57
16.2	High levels of acidity may be a problem. How may it be handled?	57
16.3	Characteristics of so called ‘cider apple cultivars’	57
16.4	Comment on the importance of ripening levels of fruit and berries for wine making	57
16.5	Characterize the process of cryo-concentration and the impacts on the juice quality and the potential for wine style development	57
17	Domestication of wild berries	57
17.1	Why may wild berries be attractive to domesticate?	57
17.2	Comment on some major challenges/barriers.	57
17.3	Describe important yield and quality components in wild/European blueberries.	57
17.4	Blueberries are one of few fruiting plants adapted to low pH soils. Comment on the challenges it causes in growing the plants.	57
17.5	Comment on the importance/impacts of propagation method in European blueberries. European blueberries.	57
4	Abbreviations and Explanations	58
	Appendices	60
1	Appendix 1 - Practical Exercise 01	60

Chapter 1

Lecture Notes

1 Introduction

The following lecture notes have been compiled by combining personal notes taken during class, personal highlights from the lecture slides, and targeted use of generative AI. The AI was provided with my notes and the highlighted slides, together with carefully designed prompts, to ensure that it focused on the most relevant and interesting aspects of each lecture. This approach aims to produce a coherent and focused summary that reflects both the course content and my individual learning perspective.

2 Lecture 01: Introduction - 01/09-2025

Fruit and Berry Crop Physiology and Quality or Fruit and Berry Crop Physiology, Quality and Use
Professor: Torben Toldam-Andersen, KU-PLEN

The course aims to provide broad knowledge on the physiology and growing of fruit and berry crops (including pit-, stone-, cane-, and bush fruits, other berries, nuts, and fruit vegetables), focusing at the whole plant and organ level. Key objectives are the development of specialized knowledge on a course topic, the ability to analyze a crop to identify important aspects in growing and quality, and understanding how fruit quality is influenced, specifically balancing productivity vs. quality. The curriculum seeks to build links between the production and quality of raw materials and the food science aspects of their use, including sensory aspects. The underlying idea is to address both the Plant science and the Food Science aspects of fruits and berries, emphasizing the importance of these crops for human health and addressing lacking knowledge regarding the potentials in different genotypes (raw materials).

The course focuses primarily on temperate fruits, nuts, and berries grown mainly in open field or tunnel systems. While fruit vegetables such as tomato may be included in the individual report, the emphasis includes common commercial systems, organic growing, and relevance for small-scale/home gardening. These crops are characterized as Diverse (many species and cultivars), high value crops, demanding intensive hand labor and resources, and often requiring manipulation at the single plant level. Although often fresh consumed/used, there is a large diversity in processed uses, such as juice, jam, fruit wine, vinegar, cakes, desserts, and ice cream. They are important to human health and provide pleasure through nice aroma and taste.

The course content is structured around eight main topic areas:

1. Genetic basis and Cultivar variation

2. Yield and quality components and determinants, including Organ development and interactions
3. Allocation of dry matter and nutrients, focusing on 'Sources' and 'sinks' in fruiting plants
4. Effects of pre-harvest factors on fruit quality, such as the importance of control of plant vigour and shape
5. Content and development of secondary compounds in fruits
6. Maturation, ripening, and assessment of optimal harvest time and quality aspects
7. Post-harvest usability (fresh consumption, cooking, industry processing, especially juice)
8. Sensory aspects of quality evaluation

Pedagogic methods utilize a mix of Lectures, Exercises ('Hands on'), Written assignments, Student lectures, and Excursions. Specialization is achieved through the individual report and the student lecture, which provide case specific supplements to the general lectures. The individual report is an opportunity to develop specific competence on a free topic relevant to the course (e.g., Bud dormancy, Aroma development, Canopy management). The report structure mandates an Intro (1-2 pages), Detailed presentation and discussion based on international literature (4-6 pages), and Summary/conclusion (½ - 1 page), not exceeding 8-9 pages in total (max 10 pages including figures). Students must use their own writing and avoid 'copy-paste' and plagiarism.

The assessment is an Oral exam lasting 25 minutes in total. The exam includes a short talk about the individual report, followed by two curriculum questions: one focusing on crop physiology and one on quality aspects. The final grade is based on 50% the individual report and 50% the curriculum questions.

3 Lecture 02: Yield and quality components - 01/09-2025

Professor: Torben Toldam-Andersen, KU-PLEN

The overarching aims in fruit growing are to maximise fruit yield and maximise fruit quality. The tools utilized to achieve these aims include Genetic choice (cultivars) and Growing technique. Fruit science focuses heavily on the physiological background informing the growing technique. Maximized Yield is typically expressed as tons/Ha (or hekto L/ha of juice or wine) and is fundamentally a product of the components Fruit number X Fruit size. Determinants influence the size or the level of development of single components within the genetic potential. These physiological elements are connected to growth and development and can be influenced by growing techniques, climate, and growing conditions.

Maximized fruit quality is less clearly defined than yield and may vary with fruit species and final use. Important quality parameters include Colour and other aspects of appearance (especially for Fresh consumption), Taste components (sugar, acid, aroma), and 'Health components' (vitamins, phenols) which are a focus of research. Fruit size serves as both a yield and a quality component. The presence of unwanted substances such as pesticides is also an important aspect of fruit quality. The optimal compromise is crucial because physiological and technical factors may have opposite effects on yield and quality components. The environmental impact of the growing technique used, such as carbon footprint and pollution, also requires knowledge.

The physiological elements that serve as yield and quality components and determinants include:

1. Planting system and Growing system (fx Rubus, strawberry), determining Number of plants/ha.
2. Plant size and structure, including Elongation growth, shoot type development, Bud development, and Flower bud initiation.
3. Bud number/plant, bud type, and the ratio of leaf bud/flower bud.
4. Flower development, specifically Number of flowers/cluster and Flower quality (e.g., Number of seed primordia, Position in cluster).

5. Pollination, fertilization, and initial set, defining Initial fruit number/flower.
6. Fruit drop (June drop), which determines the Final fruit number/initial number of fruits.
7. Fruit growth and fruit development, the Leaf/fruit ratio, and specific factors like Number of seeds/fruit (Pollination) and Amount of flesh/seed (in Strawberries), leading to Fruit size and quality.
8. Yield/ha (Fruit number x fruit size) and Fruit quality (content).

Specific crop examples illustrate the variability: Sour cherry yields may vary drastically, from 3 to 18 tons/ha, due to factors like Bud death (sometimes reaching 90%) and Fruit set percentage. Unstable yields in sour cherry 'Stevensbær' pose a significant problem for the Market (Industry). In strawberries, vegetative growth (runner production) is influenced by long days and high temperature, while flower cluster formation is promoted by short day and low temperature. Pollination is very important in strawberries as the final fruit size depends on the Achenes/berry. For grapes, yield components include Number berries/cluster x weight/berry and Number of clusters x weight/cluster, influenced by factors like pruning, thinning, and the growing system. Analysis of 51 grape cultivars harvested in 2018 showed a high correlation ($R^2 = 0.9516$) between Yield kg/plant and total sugar (Fruktose + Glukose g/plante).

4 Lecture 03: Bud and shoot development - 01/09-2025

Professor: Torben Toldam-Andersen, KU-PLEN

This lecture explores Bud and shoot development, covering Meristems, Bud types and shoots, and seasonal Growth patterns. Embryogenesis initiates plant development by establishing the primary meristems and determines the Apical ↔ basal axial development, including the Shoot apex, Hypocotyl (The stem), Root, Root apical meristem, and Root cap. It also establishes Radial patterning, such as Epidermal cells, Cortical tissue, and the Vascular cylinder. The initial stages of development include Cotyledons (The first leaves). Development transitions from meristem to shoot, detailing the structure of the shoot apex in 3D and the Transition from vegetative to floral meristem. Variation exists among species in how strong the xylem or wood develop, with Trees having strong development and Bushes and canes exhibiting weak development. The purposes of leaves include Production of assimilates (the source of carbon), Water transpiration resulting in uptake and transport of water and nutrients, Production of hormones, and Control of water status/cooling.

Buds are defined as locations of growth with a potential for development. The Types (fates) of buds include Vegetative, Generative, Sleeping (not dead), and Dead. Buds determine the plant's dimensions by forming shoots of different lengths, such as Short shoots (spurs), emphasizing the Importance of the shoot's position. In strawberries, the vegetative bud results in a side crown or a 'runner'. Pome fruits, exemplified by apple and pear, possess mixed buds, while Stonefruits utilize 'naked' buds. The topography of flowerbuds differs between pome- and stonefruit, which raises questions about which structure bears flowers in the terminal position on short shoots and the Importance of this difference.

Seasonal growth patterns involve a Flush of growth in spring and early summer, followed by Growth termination or indefinite growth, depending on the species. Terminal bud formation also occurs, alongside Adventive buds formed along with shoot development. The timing of terminal bud formation depends on Shoot type (short spur is early, long terminal shoot is late) and Vigour level (strong is late, weak is early). These factors interact: low vigour results in fewer and more short shoots, whereas high vigour leads to more and longer shoots.

Factors influencing seasonal growth include:

1. Shoot type and vigour.

2. Correlative inhibition within the plant, defined as 'communication among buds'.
3. Apical dominance, which influences adventive buds and inhibition. This involves Polar gravitropic transport and the activation of adventive buds upon Removal of the terminal bud. The underlying Auxin - Cytokinin balance varies among species (peach < apple < sweet cherry) and strongly influences the Branching pattern.
4. Competition for assimilates, specifically the relationship between fruits ↔ vegetative growth.
5. Number of growing meristems in top/root ratio.
6. Tree age, where Older trees produce many shoots but exhibit weak growth, and Young trees produce few and strong shoots.
7. Management and climate factors, such as Pruning, water and nutrient availability.

Development after terminal bud formation Happens suddenly, and differentiation continues inside the bud(s). A compact shoot is formed, and development continues until bud break next spring, though the intensity of this differentiation varies (dormancy).

5 Lecture 04: Flowers and flowerbud development - 05/09-2025

Torben Toldam-Andersen, KU-PLEN

The lecture addresses Flower and flowerbud development, encompassing Phases and theories, Structure - node development, Factors, and specific examples including Apples, Cherry, Currant, Raspberry, and Strawberry.

Flowerbud formation occurs across three distinct phases:

1. Induction (May - June)
2. Initiation (July - September)
3. Differentiation (July - May)

Historical theories regarding flowerbud formation include the 1822 concept that the flower bud is formed through April - June (induction) and then 'filled out' through July - October (initiation - differentiation). Later, the 1920's highlighted the C/N-ratio as decisive, while the 1970's focused on the Cytokinin/Gibberellin-ratio and Source-Sink relationships, stressing availability and activity. Node development speed influences the bud's fate: slow development results in remaining Vegetative, while relatively fast development leads to Floral buds. The time required for node formation is termed the plastochron.

The structure of the flower bud is akin to a compressed shoot. In apple, a flower bud contains 3 bracts, 6 real leaves, 3 transitional leaves, and 9 bud scales, totaling 21 nodes, with a minimum critical node number approximately 18-20. The flower organs are formed sequentially from the outside inward: sepals are formed first, followed by petals, which are formed second. The 'King flower' is formed first. Pomefruit (apple, pear) flower buds are "mixed" buds, often terminal, and are considered "strong" buds (high critical node number). Apple buds are less developed in the dormant stage, with a larger proportion of differentiation occurring in the spring. Conversely, Stonefruit (cherry, plum) flower buds are "naked" buds, always leaf buds in the terminal position, and are "weak" buds (low node number). Cherries reach an advanced stage of development before entering dormancy, which may render them more sensitive to winter damage. Sour cherry (*Prunus cerasus*) leaf buds contain 7 nodes, while flower buds contain 13 nodes.

Flower bud formation in tree fruit is affected by growing techniques related to Vigour or Source-sink dynamics:

1. Variety

2. Tree age
3. Rootstock
4. Dormant pruning
5. Root pruning
6. Light condition
7. Fruit load
8. Thinning
9. Nutrition
10. Irrigation

For shoot growth, a minimum length (e.g., >15 or preferably 20 nodes per shoot) is necessary before initiation. Initiation usually spreads upwards the shoot, except for the terminal bud, and lower buds do not form flowers. Very strong (long) shoot growth, potentially induced by Gibberellins, may counteract formation. Short shoots (spurs) easily form flowers and show greater bud density, a higher percentage of live buds, a higher percentage of flowerbuds, and more flowers/bud than long shoots. Climate factors play a role, as flower initiation normally happens if day length is <12-14 hours, and decreasing temperature has a positive effect.

In raspberries (*Rubus*), initiation requires reaching a minimum physiological stage, and is promoted by short day and low temperatures, typically starting from the top of the cane earliest in shoots where growth has terminated. Normally 2/3 of buds become flowers, but low positions on the cane are usually without flowers. For yield, the Number of laterals/cane and flowers/lateral are the most important components. Strawberries exhibit types ranging from obligate short day (June bearers), where day length is critical (<11-14 hours), to day neutral types that initiate at both short and long days. Vegetative growth (runners) is promoted by long day and high temperature. Warm periods in late autumn have positive effects on flower development, leading to more flowers/bud. This differentiation affects flower quality; for instance, strawberry 1. order flowers have 400 pistels, while 4. order flowers have only 80 pistels. In currants, the terminal flowers in the string are often poorly developed, resulting in few seeds, small berries, and a high risk of drop.

6 Lecture 05: Bud dormancy - 05/09-2025

Torben Toldam-Andersen, KU-PLEN

This lecture details Bud dormancy, covering its development, breaking, and specific studies on Sour Cherry 'Stevnsbær' buds in winter. Key concepts include Growth phases during the year, Dormancy terminology, and Metabolic activities during dormancy. The lecture introduces the concept of chilling units (CU) and Growing Degree Hours (GDH). Chilling requirement estimates for various fruit and nut species are noted. Dormancy models, including the calculation of Chill Units (Saure, 1985), and the Growing Degree Hour model are essential tools. A GDH curve was specifically developed for 'Montmorency' cherry (Anderson and Richardson 1986). Studies examining 'Redhaven' peach flowers and lateral/terminal buds explored the effect of temperature and length of chilling period on percent budbreak and GDH demand for budbreak (Scalabrelli and Couvillon 1986). Chilling periods of 600, 1340, and 2040 Chilling Hours were analyzed.

The yield components of sour cherry are multifaceted and include:

1. Tree size
2. Shoot number and type
3. Bud type
4. Number of flower buds

5. Flowers/bud
6. Bud death
7. Fruit set%
8. Number of fresh flowers in spring
9. Initial total number of fruits
10. Fruit drop
11. Final number of fruits
12. YIELD

Floral bud mortality is a critical factor. Dormancy levels in sour cherry buds can be determined by forcing shoots. Studies investigated the effects of irrigation and nutrients. Forcing of buds showed an increase in water content of forced buds over the period of October 4 to November 29 when under +drip conditions (both tip and base) compared to -drip. Shoot studies indicated a relationship between High water content and high bud death, and Low water content and low bud death. Observed damage included a missing pistil or a necrotic pistil.

Research examined the Effects of urea (0.75% and 4% urea) on sour cherry. In May 1996, the 4% urea treatment significantly reduced Living buds (79.3% compared to 91.3% in Control, LSD 4.0) and decreased Flowers/tree, Fruit set% (7.9% compared to 12.8%), and Berries/tree. Shoot length and Buds / meter shoot were not significantly affected. Furthermore, in November 1998, buds on annual shoots treated with 4% Urea showed reduced ABA content (1675 ng/drymatter versus 3948 in Control) and increased Water content (118.0% of drymatter versus 108.1% in Control), although Bud size was not significantly different. Freeze tests were conducted to determine the hardiness level in 'Stevnsbær' at -6°C (laboratory) and -7°C (field). Bud death was analyzed in relation to rootstock (Weirroot, Colt, Pavium Grundstamme) and frost exposure during winter. Counting flowers is noted as a considerable task.

7 Lecture 06: Flowering, pollination and fruit set - 08/09-2025

Torben Toldam-Andersen, KU-PLEN

The physiological processes of flowering, pollination, and fruit set are critical determinants of Yield, alongside factors such as Flower number and quality, Effective Pollination Period (EPP), Pollen quantity and quality, Pollen transfer, Initial fruit set, and June drop, leading to the Final fruit set. Flower quality is defined as "The capacity to develop into a fruit if pollinated with the right pollen at the right time". Indicators of flower quality include Ovule longevity, Flower weight, and Number of flowers per cluster (position in cluster). Stone fruit quality depends on carbohydrate reserve availability, while pome fruit quality depends on burse leaf area.

Fertility varies, categorizing species as Self fertile (Self compatible), such as apricots, peaches, strawberries, and grapes, or Self sterile (Self-incompatible), such as hazel, pear, and apple. Sterility is not absolute; in normal self-sterile conditions, own pollen mostly fails, but cross-pollination often results in faster pollen tube growth. Parthenocarpy is fruit development without fertilisation, without seeds. Some plum cultivars, such as Opal and Victoria, are self-fertile. Self-incompatibility is controlled by S alleles at the S-locus. Self-fertile sweet cherry cultivars (e.g., Stella, Lapins) are universal pollinators, often possessing a mutated S4 allele. Parthenocarpy or Apomixis (fruit or seed development without fertilisation) may be mistaken for self-compatibility.

Important flowering related terms include:

1. **Monoecious:** Having pistillate (female) and staminate (male) flowers on the same plant.

2. **Dichogamy:** Opening of male and female flowers at different times on a monoecious plant, which ensures cross-pollination.
3. **Protandrous:** Male flowers open before female flowers.
4. **Protogynous:** Female flowers open before male flowers.

Pollen tube growth requires its own reserves at the stigma surface, followed by interaction with style tissue (proteins, carbohydrates). The Effective Pollination Period (EPP) defines the time when fertilization and fruit development occur. EPP varies with flower quality and is reduced by factors such as low nutrient status and high cropping level.

The required quality of pollination depends on the species:

1. For **large fruited species** with intensive flowering, pollination does not need to be optimal, but should be optimal if flowering is weak ($< 40 - 50\%$ of max). Heavy crop load leads to small fruits.
2. For **small fruited species**, yield is mainly determined by fruit number, and pollination should always be optimal.
3. For **species with several pistels - embryos** (where fruit size is seed-number related), pollination should be optimal.

Pollinator demands include Quantity (e.g., Malus types like M. Floribunda), Quality (compatibility, ability to germinate at low temperatures), and Timing (overlapping flowering periods). Triploid cultivars (e.g., Gråsten apple) do not produce fertile pollen.

Pollen transfer occurs via Wind (for self-fertile species like sour cherry), Insects (for cross-pollinators), or artificially. Critical distance for transfer is 10-15 m. Honeybees (47%) and Soil living bees (25%) are common pollinators. Critical flying limits are 13-14°C for Bees and 8-12°C for Bumblebees. Bumblebees visit 2-3 x as many flowers as bees. Experimental trials on 'Stevnsbær' sour cherry showed that the addition of 'artificial wind' (6 km/t) increased fruit set (%) and yield (kg/tree) compared to controls, and that early pollination (Day 1) resulted in higher fruit set than later pollination (Day 2). Pollinator branches and hand pollination also improved fruit set, demonstrating that pollination is often not good enough.

8 Lecture 07: Fruit Thinning - 08/09-2025

Professor: Torben Toldam-Andersen, KU-PLEN

Fruit Thinning is a necessary practice carried out in Large fruited species, specifically Apple, Pear, Plum, Peach, and Grapes (where grape clusters may be seen as similar to a big fruit). The primary objectives of thinning are to maximise fruit size distribution, prevent the breaking of branches, improve flower development, and stabilize cropping (prevent alternance). Thinning also aims to improve quality, resulting in a higher sugar%, Dry matter+acid, and potentially improved firmness, although larger fruits are naturally softer. A negative aspect is that Increased size may lead to an increase in 'Bitterpit'.

The Timing of thinning is crucial. Early thinning is important as it affects both return bloom and fruit size. Late thinning has no effect on return bloom, and the effect on size decreases with delay in time. Very late thinning (starting at ripening) primarily affects quality by increasing sugar accumulation during ripening. Thinning timing effects on size were observed in apple, and in 'Clara Frijs' pear on three different rootstocks.

Thinning can be performed using various methods:

1. Hand thinning.

2. Hormones: BA (benzyladenin/Cytokinine), Ethephon, NAA (1-naphthylacetic acid, 'Pomoxon'). (Note: NAA is Not registered in DK).
3. Fertilizers: ATS (ammoniumthiosulfate), lime sulphur.
4. Organic compounds: Oils, soaps, salts.
5. Other: shade, mechanical.

Fertilizers such as ATS (ammoniumthiosulfate) function by burning the stigma and are applied before pollination, often requiring several applications to be effective. ATS may burn leaves, but is low cost, effective, and environmental friendly. To reduce leaf damage, fast drying must be ensured by using a low amount of water and spraying on dry leaves, and the weather must stay dry after spraying. Urea is similar to ATS but less effective, and Lime Sulphur is used in plums but is less effective in apple. The general mode of action for many organic compounds tested is the burning of the stigma, often associated with leaf damage when effective.

Mechanical thinning, developed in Germany, requires optimized tree training, but risks removing the highest quality buds. Constraints include thinning delicate and fragile fruitlets at an early stage without bruising the remaining fruit or removing too many leaves and spurs. Strong and sturdy training structures are necessary, especially when using M9-style rootstocks whose trees are firmly attached to the top wire. Shade experimentally induces thinning through increased internal competition and can be applied for a short period. Leaf damage caused by fertilizers is avoided in shade methods because the resulting permanent reduction of carbon supply is not wanted.

Factors affecting the efficiency of thinners include Temperature and humidity (increased effect with high temp and high humidity), Volume of spray, Additives (surfactants, which can cause leaf damage), and the benefit of Repeated treatments. Cultivar differences exist, as Strong flowers are difficult to thin. Thinning is enhanced if trees have a weak root system, drought stress, Nitrogen deficient status, few flowers in the cluster, large yield the year before (indicating weak flowers), young trees in strong growth (high vigor), or bad pollination (few bees). Thinning is decreased if trees are in good balance (growth and nutrition), fruits sit on horizontal or weeping branches, many flowers are in clusters with well developed leaves (strong flowers), trees are open, or pollination is optimal.

Thinning serves as a growth control tool. If growth is weak, thinning should be early, especially on weak shoots. If growth is strong, thinning should be late, keeping most fruit on strong shoots. The top of the canopy can be used to judge vigour, particularly in species like Solaris which has a strong canopy. Management techniques such as Topping send carbon down to clusters, enhancing growth in shoots which are not topped and stimulating lateral development. Removing inner leaves ensures good ventilation and facilitates later 'green harvest' or thinning. Furthermore, dense clusters may be 'tipped' before they close up, and big clusters (Fx Bolero) may be reduced in size. A high crop load can be sustained on a strong root stock.

9 Lecture 08+09: Source-sink and carbon allocation - 08/09-2025

Professor: Torben Toldam-Andersen, KU-PLEN

This lecture addresses Source-sink and carbon allocation, focusing on definitions, effects, internal competition, seasonal changes in allocation, and the relationship between source-sink dynamics and fruit size. The basis of growth is founded on the fact that fruits generally contain about 80-90% water, and of the dry matter, at least 90%

originates from assimilates formed in the leaves. Assimilates must be translocated from leaves to fruit. The end product exported is mostly sucrose, but in Pome and Stone fruit, sorbitol is the end product.

The source strength is defined as the ability of assimilating tissue to produce and export assimilates, quantified by Source size (leaf area/leaf number) multiplied by source activity. Sink strength is the ability of importing organs to attract and metabolize assimilates, quantified by Sink size (fruit number, number of growing shoot tips) multiplied by sink activity.

Cropping significantly impacts plant physiology, as demonstrated by apple studies. High cropping stimulates Photosynthetic intensity (140-210 relative to 100 without/few fruits), Translocation intensity (160-175), and Stomata opening degree (165). However, high cropping shifts Dry matter production allocation: while total dry matter production increases slightly (112 vs 100), the dry matter allocated to roots drops sharply (<1 vs 23), and 81 relative units are allocated to fruits. The sugar + sorbitol concentration decreases during high cropping (80 relative units). Increased CO_2 level leads to increased Sorbitol% (13.4 vs 12.2), Sucrose% (3.6 vs 2.7), and Transported $14^\circ C$ (% of uptake: 59.6 vs 53.6).

Internal competition involves 'Supply - Demand'. Apple has a strong sink strength but a relatively weak source, whereas Black currant has a weak sink strength but a strong source. Internal competition, characterized by the Fruit/leaf ratio, is strongly important in large fruited species such as apple, pear, plum, and large sweet cherry, necessitating thinning. However, the effect is weakly important in small fruited species like currants, blueberry, raspberry, and sour cherry. Sour cherry 'Stevnsbær' showed no difference in total dry matter (%), sugar (%), acid, or colour when comparing low (7 kg/tree) versus normal (25 kg/tree) cropping levels.

Source and sink strength of organs vary dynamically with development and time. Source strength increases with shoot growth, light, and temperature. Types of competition include internal competition between shoots and fruits, and between fruits and flowers.

There are two primary ways to increase fruit size:

1. By changing the fruit/Leaf ratio, thereby increasing Assimilate availability, which results in a Positive correlation between size and dry matter concentration (internal quality).
2. Through an increase in sink strength (or sink 'vigour'), such as by K fertilisation, which results in a Negative correlation between size and dry matter (internal Quality).

At the tree level, no correlation exists between size and quality.

10 Lecture 10: Table Grapes - growing techniques and cultivars - 12/09-2025

Professor: Torben Toldam-Andersen, KU-PLEN

This lecture focuses on aspects connected to the tunnel growing of table grapes and berries, specifically within an unheated tunnelsystem. The slides cover inherent benefits and challenges of this system. The experimental tunnel was established in 2007, and is relatively small (5m wide, 2,75m tall). It uses an internal trellis system allowing training in 3 levels (low, medium, and high stem plants) for approximately 120 plants. The training system is simple and similar for all: 2 Codon on low, medium or high trunk height, typically using short 2 bud stabs later developed into double stabs with (3)-4 shoots. Ventilation is implemented in both sides and utilizes a fan.

Management priorities are defined across seasonal stages:

1. **Budbreak and the early growth:** Focus on the importance of humidity, Shoot thinning (when and how), selecting the best shoot on a spur to carry the fruit, and selecting the best shoot for next year.
2. **Flowering and fruit set:** Identifying Important factors and management, and Cluster thinning?.
3. **The summer period:** Canopy management (How are the shoots treated?), and managing Diseases (which are most important? prevention or reduction methods?).
4. **Late summer and autumn:** Fruit development (including cluster thinning and shaping), determining what to do, when, and why, and establishing When to harvest?.

The descriptor system used for testing table grape cultivars in Denmark (VitiNord Nov 2015) included 38 cultivars. The descriptor employs a scale from 1-9 for scoring parameters, where 9 is generally the best, most wanted, or most attractive. However, the scale may be modified so 1 may not always be the 'less wanted', or 5 may be the best from a growing perspective. The system uses 25 Culture technical parameters (e.g., C1 Vigor, C3 Formation of laterals, C6 Fruit-fullness from the second eye, C17 Resistance to peronospera), 36 Technological parameters (e.g., T5 Cluster weight, T9 Berry shape, T23 Sugar content when ripe, T30-32 Aroma), and 25 Morphological parameters.

Specific parameters reflect desired growing characteristics; for example, T7 (Berry weight at BBCH 89) scores 9 for Very large (>7 g), while C3 (Tendency to formation of laterals) scores 9 for Very weak occurrence (1 lateral/main shoot) because low occurrence saves Work with canopy management. C10 (Density degree of the cluster) rates both very dense (1) and very loose (1,1 or 3,3) low, while Appropriate (7) and Very appropriate (9) densities are rated high. The Relative productivity index (C25) is calculated as (cluster number/no of main shoots) x weight of 1 cluster in gram. Cultivars are ranked based on an average equivalent score derived from summing and normalizing the Culture technical and Technological parameter scores.

Highly ranked cultivars include Augustostuzi Muskotaly, Conegliano 218, New York Muscat, and Palatina. These cultivars generally achieve a high sugar content (score 7) of about 19% brix, and acid levels (score 5-7) between 6 to 6,8 g/L. The cultivar Augusta exhibits medium vigor, Above medium fruit fullness from eye no 2, and Good resistance against Oidium, Peronospera, and Botrytis. Augusta clusters are large (780 g), and the berries are very large, exhibiting a very fine crunchy structure and very high eating quality. Augusta begins harvest in week 37 in the plastic tunnel and provides a qualitative and quantitative improvement in relation to its parents, 'Italia' and 'Königin der Weingarten'.

11 Lecture 11+12: Fruit development of large fruited species - 15/09-2025

Professor: Torben Toldam-Andersen, KU-PLEN

The development of large-fruited species involves examining fruit structure, growth curves, volume changes in apple, gravity, intercellular air space, and changes in fruit shape during development. Fruit shape, as noted by Westwood (1978), is influenced by climate, with cool climates often resulting in more pronounced shapes (greater length/diameter ratio) compared to warm climates. Pear shape (e.g., 'Conference') is known to be affected by seeds, as demonstrated by parthenocarp fruit. Fruit size is determined very early, influenced by cell size and thinning practices. The relationship between fruit size and yield dictates that a significant leaf area is required per fruit; for example, 30 leaves at 20 cm² per leaf equals 600 cm² or approximately an A4 sheet/fruit.

Quality effects observed in plums (Vangdal 1982) show strong interactions between fruit load, size, and sugar content. For instance, reducing the number of fruits per spur significantly increases sugar content; fruits/spur at

0.14 resulted in 11.1% total sugar and 7.6% sucrose, compared to 0.98 fruits/spur yielding 5.6% total sugar and 2.4% sucrose. Similarly, larger fruit diameter (>40 mm) correlates with higher sugars (13.1% total sugar) than smaller fruit (30-35 mm, 11.1% total sugar). Increased skin color (80%) also shows correlation with increased sugar levels (13.2% total sugar).

Yield relations in fruit growing can be modeled as a Michaelis-Menten kinetic system. If Y is yield t/ha and x is the number of fruits/ha, then $Y = (k \cdot x)/(K + x)$. Fruit size is defined as $Y/x = k/(K + x)$. The physiological parameters k and K are crucial: k represents source activity, determined mainly by total assimilate production (potential max. yield), and $1/K$ represents sink activity. The term k/K is the potential maximal fruit size. Comparison across different tree statuses shows that older, well-established trees typically exhibit lower sink activity ($1/K$ lower) but possess a larger source capacity (k) than young trees. Dense trees demonstrate lower source activity.

Internal quality, maturity, and harvest time are studied alongside external qualities like size, focusing on internal chemical changes during development. Apple C-14 labelling studies conducted end June and end July tracked chemical changes. In apple fruit development, both sorbitol (Sor) and sucrose (Suc) are unloaded into the cell wall space. Sor is taken up via sorbitol transporter (SOT). Suc is transported either directly by sucrose transporter (SUT) or converted to fructose (Fru) and glucose (Glc) by cell wall invertase (CWINV), followed by transport via hexose transporter (HT). In the cytosol, Sor converts to Fru via sorbitol dehydrogenase (SDH). Suc converts to Fru and Glc via neutral invertase (NINV) or to Fru and UDP-glucose via sucrose synthase (SUSY). Glc and Fru are phosphorylated by hexokinase (HK) and fructokinase (FK). F6P enters glycolysis/TCA cycle, and G1P is utilized for starch synthesis. Sucrose can be re-synthesized using UDPG and F6P via sucrose phosphate synthase (SPS) and sucrose-phosphatase (SPP). Finally, most non-metabolized Fru, Glc, and Suc are stored in the vacuole using special tonoplast transporters, where Suc can also be broken down by vacuolar acid invertase (vAINV).

12 Lecture 13+14: Fruit development of small fruited species / berries - 15/09-2025

Torben Toldam-Andersen, KU-PLEN

This lecture details the development of small fruited species, focusing on fruit types, growth phases and curves, size factors, chemical changes, and mechanisms for sugar and acid accumulation. Aggregate fruits are derived from many ovaries. Growth curves show that berries and stone fruits have a distinct Stage II characterized by slow size enlargement. A berry, such as a grape, gooseberry, or a currant, is a multiseeded fruit derived from a single ovary. The cell division period, specifically the first weeks, is critical in fruit development for both small and large fruited species. Fruits primarily consist of water (85-90%) and 10-15% dry matter, with about 90% of the dry matter originating from assimilates produced in the leaves.

Internal competition can be very strong during the cell division phase due to vigorous growth in all organs. Fruit drop in black currant plants was observed in shaded plants, but effects were minimal or absent in sun-exposed plants. For strawberries, size is a linear function of achene number/berry at a given achene density. Berry weight is calculated as $(\text{total achene number/berry} - C) \times F$, where C is a correction factor depending on the cultivar, and F is the flesh density in g/cm^2 . While maximal swelling yields 6 *achenes/cm*², it is often lower, resulting in 8-12 *achenes/cm*², with water being probably the most important factor. In black currant, berry size is positively correlated to shoot growth intensity. Conversely, berry size is negatively correlated with the concentration of total and soluble solids, although acid content increases. This suggests the main effect on black currant composition is via sink activity. Cultivar variation is often a larger factor influencing development than growing techniques or

climate. Fruit refixation of CO_2 occurs.

The relative sweetness of common compounds (scale where Sucrose =100) is Fructose: 173; Glucose: 74; and Sorbitol: 50. Relative sourness (scale 1-9; 9=most sour) is highest for Malic acid (8.2), followed by Tartaric acid (4.3) and Citric acid (3.6). Chemical analyses of black currant cultivars 'Tenah' and 'Ben Nevis' reveal changes in sugars (Sucrose, Glucose, Fructose) and acids (Citric and Malic acid) over time. A study noted that 1992 was a very warm summer. There is a relationship between water content and citrate content in black currant. Light affects fruit quality; the Sucrose/citric acid ratio in black currant was 0.28 in the Sun top compared to 0.15 in the Shade bottom.

Regarding vascular mechanisms, in the early phase (pre-veraison), the xylem is fully functional, transporting water and nutrients into the fruits. Post-veraison (developmental phase 3), the xylem loses its conductivity, leaving the fruit connected to the plant via phloem transport, which delivers sugar; the central connection to the seeds is maintained. Studies showed that when gas exchange via the berry surface is reduced by a coating with Vaseline, the sugar accumulation is impaired. Furthermore, Vaseline coated fruits combined with leaf removal resulted in fruit drop.

Fruit quality varies across berry types:

1. Black currant has the highest recorded ranges for Total dry matter (%) (17-28%) and C-vit mg/100 g (95-253).
2. Red currant exhibits the highest Acid g/100 g range (2.7-4.1).
3. Strawberry has the lowest Total dry matter (%) (7-11%) and Acid g/100 g range (0.8-1.2).

13 Lecture 15+16: Aromas in fruits - 22/09-2025

Mikael Agerlin Petersen, FOOD-DCB

This lecture explores Aroma in fruits, specifically addressing advances in volatile research, differences in aroma patterns, and the influence of the fruit/leaf ratio. Flavour is defined as the combination of basic tastes (Sour, Salt, Sweet, Bitter, Umami), odour (orthonasal and retronasal), and trigeminal stimuli. Aroma compounds are organic compounds with a certain volatility that interact with our olfactory receptors (not taste). They are effective in very small amounts; for example, 10 ppb of 2-Methoxy-3-isopropylpyrazin corresponds to mixing one teaspoon into a small swimming pool, resulting in a significant odour.

Aroma analyses are typically performed using GC-MS analyses. Gas Chromatographs (GC) cannot inject large amounts of water, lipids, sugars, or amino acids, but accept gas ($\approx 1\text{mL}$) or organic solvents ($\approx 2\mu\text{L}$). Headspace methods are frequently used, such as Dynamic Headspace Sampling (DHS), where headspace is continuously removed and collected/concentrated on a trap, ensuring equilibrium is never reached. The GC separates mixtures of volatiles, while the Mass Spectrometer (MS) utilizes an electron beam to create charged fragments of molecules, which are filtered by a Quadropole.

The aroma of a food is normally complex, resulting from a combination of many aroma compounds. Small peaks in a chromatogram (which may show >75 peaks) can also be important.

Fruit volatile compounds are mainly comprised of:

1. Esters
2. Alcohols
3. Aldehydes
4. Ketones

5. Lactones
6. Terpenoids
7. Some sulfur compounds

The specific content of these compounds determines, to a very high degree, the distinctive sensory quality of a fruit.

Aroma compounds are produced during ripening. Climacteric fruits show a distinct production controlled by ethylene. Studies on transgenic 'Royal Gala' apple (AO3) unable to synthesize ethylene confirmed that exposure to exogenous ethylene induces ripening. The genes involved in aroma biosynthesis are not coordinately regulated by ethylene; typically, only the first and final steps are ethylene regulated. Alcohol Acyl Transferase (AAT), which catalyzes the last step of volatile ester biosynthesis, is regulated by ethylene gene expression. Varieties like 'Golden Delicious' (high ester production) and 'Granny Smith' (low ester production) represent phenotypic extremes in volatile production.

The ethylene receptor blocker 1-MCP (1-Methylcyclopropene, "SmartFresh") binds to receptors without triggering a response. 1-MCP treatment immediately after harvest influenced the aroma profile in 'Ildrød Pigeon' apples after cold storage. 1-MCP is approved for minor use in closed storage rooms but is under re-evaluation. Aromas are generally under genetic control, but production can be influenced by Climate, Soil, and Fertilisation. The effect of the fruit/leaf ratio on aroma quality has been studied using different levels of fruit thinning (no thinning, moderate 10 cm, and heavy 20 cm) in potted apple trees.

Specific fruits contain characteristic compounds: Strawberries have one of the most complex aromas (≈ 350 volatile compounds). Furanol (2,5-dimethyl-3(2H)-furanones) is a key component, and Methyl anthranilate is responsible for the typical spicy-aromatic and flowery note of wood strawberry (*Fr. vesca* L.). Banana fruity top notes derive from volatile esters like isoamyl acetate and isobutyl acetate. Citrus characteristic volatiles include aliphatic aldehydes (e.g., Octanal) and monoterpenes (e.g., Limonene). Raspberry aroma is particularly influenced by α -ionone, β -ionone, and raspberry ketone.

14 Lecture 17+18: Organic growing of fruit and berries 22/09-2025

Maren Korsgaard, Inst. of Plant and Environment, mkor@PLEN.ku.dk

This lecture provides guidelines and legislative background for the organic production of fruit and berries. The main vision for organic production is built upon four foundational principles:

1. HEALTH: Sustain and enhance the health of soil, plant, animal, human, and planet as one and indivisible.
2. ECOLOGY: Based on living ecological systems and cycles, working with and emulating them.
3. FAIRNESS: Build on relationships that ensure fairness regarding the common environment and life opportunities.
4. CARE: Managed in a precautionary and responsible manner to protect the health and well-being of current and future generations and the environment.

Legislation governing organic production is based on EU 834/2007 (2021) and Guidelines of Organic agriculture in Denmark (Landbrugsstyrelsen). The EU Green Deal aims for 25% of agricultural land to be organic by 2030, rising from 10,5% in EU27 in 2022.

In Denmark, the total organic cultivated area was 295.233 ha in 2024, representing 11,1% of the Danish agri-

cultural land, showing a small decline (2,7%) related to 2023. The organic fruit and berry-acreage is increasing. The largest organic fruit areas in Denmark (2024, Ha) include Apple (616 Ha, 42,2% of total DK fruit-area) and Hazelnuts (504 Ha, 83,4%). Conventional Danish fruit production is often not profitable, and organic production is also risky business due to low yield per hectare. The organic apple-yield per hectare is approximately 5,5 ton/ha, but at least 15 ton/ha is necessary for a profitable organic apple-production. Organic production utilizes specific approved inputs. Organic fertilizers are always organic Nitrogen-fertilizers, including Farmyard-manure, Green manure, Rock phosphate, and Potassium sulphate*. Agents allowed for pest and disease regulation include I: Azadirachtin (Neem), I: Plant oils, F: Copper-products, F: Sulphur, and F: Potassium bicarbonate (Armicarb).

Studies suggest that organic agriculture is the most environmental friendly agricultural system. Although the climate impact per kg food is the same as conventional production, organic soil contains 10% more carbon and sequesters more C (+256 kg C/ha/year), and emission of nitrous gasses is 24% lower. Local production is emphasized, as transport and cooling represent the largest environmental impact of apples. Danish apples in season have a very low climate impact (0,1 kg CO_2 -eq./ kg food). Quality of organic products includes absence of pesticide-residues (apart from background pollution). Organic products also contain approximately 12% more health-promoting substances than conventional products, showing slightly higher content of total antioxidants and dry matter (taste). Organic apples have 4-7% more sugar.

To reduce risk in organic apple growing, planning should focus on dry climate locations, robust varieties, and well-drained soil. Preventing fungus diseases involves choosing weak-growing rootstocks, robust varieties, open pruning, low nitrogen-level, and potentially covering trees with a roof. Apple scab can be reduced by spraying sulphur or sodium bicarbonate. Insect damage can be prevented using beneficials by sowing flowering stripes and putting up nest boxes for insect-eating birds.

A trial involving five model organic apple orchards in 2004 found that the gross profit varied from - 83.000 kr/ha to +141.000 kr/ha, averaging 39.000 kr/ha, which was comparable to conventional production (\approx 35.000 kr/ha). Main problems identified were attacks from aphids and scab.

15 Lecture 19: Light utilisation and canopy management 26/09-2025

Torben Toldam-Andersen, KU-PLEN

The physiological context of fruit production emphasizes that Biomass Yield is determined by the formula: (available light)x(% light intercepted)x(Photosynthesis)-Respiration. Of these factors, the percentage of light intercepted is the only one that can be directly manipulated. Efficient utilization of available light is of key importance in light-limited conditions, as Productivity is defined as a linear function of intercepted PAR (Photosynthetically Active Radiation). Photosynthesis and carbon allocation must be understood in this production physiological context, particularly concerning yield and quality components. A strong sink, such as a high number of fruits per m^2 leaf area, stimulates a higher source activity, resulting in increasing Net photosynthesis. An optimal leaf area of 600 cm^2 /fruit is required to support approximately 1 g CO_2 /apple/day.

Canopy management techniques are essential for manipulating light interception and distribution. Light distribution is crucial, affecting development from the single leaf up to the canopy level. The light level must not be too low to form flowers, requiring a minimum of 30% of full light for flower bud development. Light effects influence dry matter allocation, plant shape, fruit size, and colour. Data indicates that at only 10% relative light level, the percentage of large fruits drops to 22% and the percentage of fruits more than a quarter red drops to 5% (Based

on Hansen 1995). The artificial application of post-harvest light underlines the very strong effect of light on red over colour development.

Light interception is fundamentally connected to the Leaf Area Index ($LAI = m^2 \text{ leaf area} / m^2 \text{ ground area}$). While light interception can be maximized through increased LAI, the optimal strategy requires a compromise between maximum interception and effective light distribution, ensuring the 30% minimum light requirement is met. An overall light interception of 50-65% and an LAI of 1.5(-2) is considered optimal. The theoretical maximum light interception (F_{max} ground area). While light interception can be maximized through increased LAI, the optimal strategy requires a compromise between maximum interception and effective light distribution, ensuring the 30% minimum light requirement is met. An overall light interception of 50-65% and an LAI of 1.5(-2) is considered optimal. The theoretical maximum light interception (F_{max}) is reduced by light transmitted between trees to the ground (T_f) and light transmitted through the canopy (T_c). If LAI reaches 3, the canopy is generally deemed too dense.

Methods applied to manage light interception include:

1. Planting system (distances, orientation).
2. Canopy shape and size.

Increased light interception can also be achieved by maximizing Foliage density, using Multi-row systems, and increasing tree height in Single row systems. The leaves must be "arranged" in 3D in an optimal way. Row orientation effects show variation across cultivars and rootstocks.

The importance of shoot type is evident as the leaf area on longer shoots is not correlated to yield, but the area on spurs is. Spur leaves are critical because they fast become exporters of assimilates, supplying the developing fruit during the first 5-6 weeks after flowering, which are critical for fruit set (fruit number) and cell division (fruit size). Spurs are highly fruitful due to a high density of flowers.

16 Lecture 20: Canopy management/General techniques - 26/09-2025

Torben Toldam-Andersen, KU-PLEN

Canopy management encompasses methods aimed at manipulating growth and development in fruit crops. The primary aims are to control tree shape and esthetics, influence cropping and fruit development, increase fruit size and quality, facilitate the renewing of shoots, and improve overall productivity. Management is also critical for addressing space limits, improving work ease for machines and picking, and controlling tree height and shoot orientation. Furthermore, canopy management helps prevent and control diseases by removing sick shoots/parts, creating open trees or canopies, and optimizing spraying efficiency.

Growth regulating methods function by affecting different parts of the tree.

1. **Top/Root Mechanisms:** Planting density, Root stock/spur types, Pruning (top), and Root pruning affect both the Top and Root. These methods change the Top/root ratio, leading to 'root hormones' becoming dominant.
2. **Top-Only Mechanisms:** Chemical growth regulators, Arching/ringing/saw cuts (affecting Stem/transport), Bending, and Thinning (fruits, flowers) affect the Top.
3. **Root-Only Mechanisms:** Ground covers and Fertilizers/water affect the Root.

The removal of tissue (shoot or root tips) is equivalent to the removal of hormone producing tissue. Affected

hormones include Cytokinins, Gibberellins, and Auxin, with Auxin sensitivity being crucial in controlling Apical dominance. Pruning methods include the Shortening of shoot, Thinning/removal, and Shoot Positioning. Pruning intensity and timing significantly affect growth: Winter 'dormant' pruning is growth promoting, while Summer 'green' pruning is growth weakening. Summer pruning reduces root growth, delays bud break, and reduces total shoot growth, resulting in shorter shoots and more spurs. Medium late summer pruning may stimulate flower bud development, but late pruning risks reduced hardiness. The regrowth response is stronger from many small cuts than from a few larger cuts, and shortening cuts are stronger than thinning cuts (for similar mass removed). Vertical branches respond with stronger growth than horizontal branches or those at 30°.

Pruning affects source and sink dynamics. It increases light penetration, which leads to changes in leaf structure and increased Photosynthesis. Strong pruning may result in compensating growth responses, such as increasing leaf size (e.g., from 9.0 cm² /leaf (No pruning) to 16.0 cm² /leaf (Strong pruning)).

1. **Effects on Fruit Set:** Dormant pruning can be positive (due to improved water/nutrient/C availability) or negative (in case of high vigor). Early green pruning is positive due to a decrease in competing sinks.
2. **Effects on Yield:** Pruning generally delays cropping in young trees. In older trees, the decrease in fruit number caused by pruning can be compensated by increased fruit size (e.g., 181 fruits/tree at 123 g/fruit (No pruning) yields 22.3 Kg/tree compared to 146 fruits/tree at 145 g/fruit (Strong pruning) yielding 21.1 Kg/tree).
3. **Effects on Quality:** Fruit size increases due to increased light (source), increased root/top ratio (sink activity), and younger tissue/increased leaf/fruit ratio. Color development is generally improved if moderate growth is induced.

The total effect of pruning depends on the level and time of pruning, as well as the vigor level and density of the tree canopy.

Pruning systems include the Spindel (central leader) and Espalier. The Slender spindel emphasizes ensuring a balance between the top (starting strong, becoming weak later) and the basis (3-5 permanent branches). Training involves maintaining a strong top shoot in young trees, preventing branching on side shoots, and removing or spur pruning strong shoots. Growth types, such as Spur types and Tip cropping types, exhibit different branching and renewal characteristics, necessitating adjustments in management practices, such as bending or tying. The principles of Zahn (1998) state that side shoots more than 50% of the trunk diameter form a split or bushy crown.

17 Lecture 21: Juice processing and Quality - 29/09-2025

Torben Toldam-Andersen, KU-PLEN

The lecture outlines the main concepts related to Fruit Juice and their importance in juice quality, recognizes the main processes involved in fruit juice processing, and identifies the theory linked to the sensory evaluation part of the course. Fruit juice is a product derived from fruits and berries, characterized by no addition of sugar, acid, aroma, or water; however, sweetened juice exists for products like Black currant and Sour cherry. Nectar is defined as juice combined with water and sugar.

Juice processing involves a sequence of steps: Harvest, Wash, Grinding, (Enzyme treatment), Pressing, Fining, Filtration, Pasteurizing, Juice storage tank, and Concentration (resulting in Aroma Concentrate). Juice can be either clear or cloudy. Key components determining juice quality include Sugars (Fructose, glucose, (sucrose)), Acids (Malic acid, citric acid), and Phenolic compounds (Quercetin, Caffeic acid, Cinamic acid, Catechin). Important ingredients vary by fruit: apple typically contains 9-14 g/100g sugar, 0.5-1.2 g/100g acid, and 5-25 mg/100g

Ascorbic acid. Sour cherry 'Stevnsbær' contains 18-22 g/100g sugar, 2,0-2,5 g/100g acid, and 10-15 mg/100g Ascorbic acid. Black currant contains 13-14 g/100g sugar, 3,0-3,5 g/100g acid, and 65-115 mg/100g Ascorbic acid. Anthocyanin content is highest in Black currant (350-450 mg/100 g).

Sensory evaluation, specifically the correlation between the sensory impression of sugar/acid and the measured sugar/acid ratio, is critical for quality assessment, using a sensory scale from -5 (very sour) to +5 (very sweet), where 0 is the optimal sugar/acid ratio. Sensorically important aroma compounds are generally volatile (e.g., Acetateesters, Butanoateesters, Butanol) or less volatile (e.g., Benzaldehyde, Bensylalkohol, Linalool, Geraniol, Eugenol, Terpenes). Cultivars exhibit diverse aroma profiles, as shown by heat map analyses of 116 Volatile Organic Compounds (VOCs) across 8 apple cultivars.

The Pressing Yield is defined as kg juice/kg fruit. The ripening stage is important: unripe fruits, with protopectin and insoluble midtlamella intact, offer good possibilities for run off, while very ripe fruits, possessing much soluble pectin and high viscosity, result in low press yields. The vacuole must be emptied for water and solutes during pressing. Pressing can be carried out using machines like a Belt Press or small hydropres for small-scale/hobby production. The resulting juice may be separated into Free-run and Press fractions.

Enzyme treatment of the pulp increases juice yield. Pectin degrading enzymes, such as Polygalacturonase and Pectin esterase, are used. Pectin consists mainly of galacturonic acid, with some acid groups esterified with methanol ($COOCH_3$). Enzyme treatment results in Lower viscosity, Better possibilities for run off, an Increase of juice yield (from 70-75% to 80-85%), Higher dry matter content, higher sugar, and Higher Methanol content.

Pasteurization, achieved by heating the juice to 90°C in 10-20 seconds, is necessary because, due to the low pH value of 3-4, only yeasts, fungus, and lactic acid bacteria can grow. Preservation methods for fruit juice include Concentration, aimed at reducing the volume from 100 l to 15-20 l. Concentration is achieved by Evaporation, Freeze-/cryo- concentration, or Reverse osmosis (membrane technology).

A major challenge during concentration by heat evaporation is the loss of volatile aroma components, as Esters, alcohols, and terpenes are more volatile than water. If only 10% of the water is evaporated, the concentration of aroma components has decreased to approximately zero. The concentration process, as illustrated, separates a 100 kg fruit juice (10% DM) into a 90 kg juice fraction (11,1% DM) and a 10 kg water + aroma fraction. Evaporation yields 20 kg concentrate (50% DM) and 10 kg water vapour + aroma components, which, via a Retification column and Condensator, yield 1 kg Aroma concentrate. Aroma components in apple juice are more volatile than those in strawberry juice, requiring a greater volume to be used as aroma fraction from strawberry juice.

Loss of anthocyanin and aroma during juice processing occurs through:

1. Loss to the press cake
2. Destruction by heating
3. Evaporation
4. Loss by clarification

The loss of the ester Ethyl butanoate and the concentration of anthocyanins decrease significantly across processing steps such as heating, enzyme treatment, pressing, clarification, and filtering. Cryo-concentration is also a method used, involving cooling curves of sucrose and glucose.

18 Lecture 22: Canopy management/ special techniques - 29/09-2025

Torben Toldam-Andersen, KU-PLEN

This lecture summarizes special techniques in Canopy management, specifically Summer pruning and Bending of shoots, which are utilized to manipulate growth and development. Summer pruning aims to open up too dense trees and is relevant for Espalier training systems. During summer pruning, unnecessary shoots are removed. Remaining shoots should be handled as follows: leaders, weak shoots and spurs are not to be touched, while others are pinched after 5-6 leaves. Several figures illustrate summer pinching techniques: Figure A shows the first summer pinching on shoot A at point b. Figure B illustrates the second summer pinching where shoot A developed shoot c, which is pinched at d. Figure C shows a second summer pinching where two shoots developed (c and d); in this case, the shoot c is removed by a cut at e, and shoot d is pinched at f, indicating that two developing shoots are considered too much. Figure D shows a similar scenario but with even stronger growth; the process involves removing the lower and weakest shoot. Figure E demonstrates a very strong growing shoot that was cut down to b in spring, resulting in all 3 buds starting to grow; the lower shoot e is pinched at g, and shoots c and d are removed with a cut at f. In problematic scenarios aimed at developing a fruiting spur (Figure F), where a spring cut at b led to three shoots (c, d, e), the lower shoot e is pinched at g, and c and d are removed at f, in the hope that the remaining shoot is weak enough to develop a spur. Another problematic scenario (Figure G) involves a shoot where lower side buds failed to develop into flowers, and only the terminal bud grew; here, the shoot is pinched short right above the base rosette of leaves as low as possible to force the lower buds to react. Figure H confirms the success of this stimulation two months later, showing a good spur development and the new shoot f pinched at g. See lecture slides for illustrations.

Bending of shoots has the primary aims of achieving Changed growth allocation and Flower bud development stimulation. Upright shoots and top exhibit the largest vigour. The degree of bending results in Reduction of vigour and Compensating growth. The degree of bending, which changes the allocation of growth and flower development, is demonstrated by quantitative data (Shoot angle, Shoot growth on bended part of branch, Shoot growth above the bended branches, Number of flowers year 2): an angle of 30° resulted in 522 cm shoot growth on the bended part and 4 flowers, while 120° resulted in 428 cm growth on the bended part and 95 flowers. Timing of bending is also crucial, as early bending results in many shoots. Studies comparing bending timing showed that March bending resulted in 28.3 shoots/tree, while bending on 1. August resulted in 8.9 shoots/tree, with shoot length varying between 32.9 cm and 37.3 cm. The mechanisms of bending involve Hormones and Apical dominance, specifically relating to Auxine and Ethylene (stress). It is important to be aware of the growth level (vigor); if the tree is weak (e.g., due to a weak root stock), bending may cause it to stop growing. Bending is an important tool in young trees.

19 Lecture 24: Nutrients, use and effects on development - 06/10-2025

Torben Toldam-Andersen, KU-PLEN

This lecture details the physiological aspects of Nutrients and fertilisation, covering Root development, Mobility, Storage, Nutrient demand and uptake, and Effects on yield components. Mobility in soil is pH dependent, where

$R_t = pH + 1/2$. Due to nutrient uptake (ion exchange) by plants, pH will fall. Approximately 4 tons/ha of Calcium Carbonate is needed every 4 years to increase soil pH with 0.5. Optimal R_t values for nutrients range from 4.0-10.0, for instance N is 5.8-8.0 and Mo is 7.0-10.0. Mobility in plants classifies nutrients as Mobile (N, P, K and (Mg)), Less/partially mobile (S, Mo), or Immobile (Ca, B, Fe, Mn, Cu and Zn). Symptoms of deficiency depend on mobility, appearing on young leaves/top of plant, older leaves/Base of plant, or the whole plant, in addition to differences in coloring at the leaf level.

Regarding Storage, nutrients move from leaves to tree storage in Autumn. N and P storage approximates 50%, K storage is about 10%, and Ca, Mg storage is only about 1%. Early leaf drop may reduce storage N. Urea spray can increase leaf N and storage N in apple. Seasonal changes show that Arginine is the storage form, and Asparagine is the transport form of soluble N. In Spring, storage moves to new growth, causing concentration to fall. Low storage results in less shoot growth and lower fruit set. Remobilisation occurs in spring, confirming that new uptake is important, requiring early nutrient supply.

Nutrient demand and uptake correlate with seasonal growth. The leaves form the major pool for nutrients, leading to the largest demand early season. Seasonal accumulation in apple fruits shows that K, P (and N) accumulation runs parallel with fruit growth, while Ca accumulation occurs only in early season. Total uptake estimates suggest N about 60 kg/ha, K about 60 kg/ha, and P and Mg about 10 kg. The type of growth influences demand. Vegetative growing plants demand double the amount of most nutrients, except K. Fruits contain very Low Ca, but are high in K.

The effects of nutrients on yield components show that nutrient use (demand) is similar across fruit species. Small fruited species with a short period of fruit development show less impact of fruits on nutrient demand. Strawberry growers may vary nutrient solutions based on three growth phases:

1. until flowering,
2. fruit development,
3. after harvest.

Flowering intensity in apple depends on cropping level and N status the year before. Moderate N level (1.9-2.6% N in leaves) results in a higher percentage of maximal flower number than low N level (1.6% N in leaves). Fruit set (relative value of flower number) increases with N% of leaves just after flowering. For Ribes, shoot growth is linked directly to bud number, flowers, fruit number, and yield. 'Stevnsbær' and Elderberry also exhibit positive responses at high N levels, while Raspberry shows less positive responses. Strawberry response depends on the growing system: young plants in a short lifetime system need higher N levels, whereas older plants need less N to prevent them from becoming too dense.

20 Lecture 25: Nutrients use and effects on quality - 06/10-2025

Torben Toldam-Andersen, KU-PLEN

This lecture focuses on the effects of nutrients on quality components in fruit and berry crops, covering Nitrogen (N), Phosphor (P), Potassium (K), Calcium (Ca), and Magnesium (Mg), along with micro nutrients, optimal status determination, and application timing.

Nitrogen (N) negatively affects apple quality components, resulting in reduced red color (due to shade and a direct effect), more green fruits, and reduced firmness. N also reduces taste, decreases the concentration of acids and

sugars, and potentially reduces aroma. Furthermore, N reduces storability. In berries, N results in less sugar, but has no effect on acid. N affects growth and development, including vegetative growth and generative growth (flowers, fruit set). Its effects on fruit development include colour (red and green) and aroma. N effects are often indirect, dominating via the fruit/leaf ratio or carbohydrate availability, although direct effects may also exist.

Phosphor (P) is not a critical nutrient in DK but may improve storability. Potassium (K) is characterized by a high demand in fruits. However, high K and high yields can be negatively correlated to bud development. K can also increase biannual bearing and induce deficiency of Mg and Ca due to the K/Ca+Mg ratio. Thus, it is crucial to be careful not to use too much K.

Calcium (Ca) is important for storability. Studies show the percentage of bitter pit attack is a function of the calcium content of apple fruits. The incidence of 'Møsk' (tissue breakdown) is also correlated with Ca content in fruits. Ca is only mobile in the xylem and not in the phloem. Seasonal uptake in fruits of Ca occurs during development. Sprays with CaCl_2 during late season may increase the Ca level. Xylem function, as observed in Braeburn' and Granny Smith' fruits assessed at 64 and 67 DAFB, shows differences in vascular function.

Magnesium (Mg) is part of Chlorophyll. Deficiency of Mg leads to necroses and leaf drop, which can be treated with leaf sprays of MgSO_4 early season. Boron (B), Zinc (Zn), and Iron (Fe) are the most important micro nutrients. There is a danger of excess levels of micro nutrients, and their content may be found in fungicides (e.g., Dithane). Leaf sprays with these minerals are frequently used.

Optimal nutrient status determination using symptoms of deficiency is often seen too late. Soil samples are difficult in fruit cultures as they vary with depth, in and between rows, drip irrigation, soil types, etc.. Leaf samples are considered the basis of next years application. They should be taken in mid - late August, specifically from the middle of annual shoots.

Regarding application: N should be applied early spring, with high levels requiring 2/3 early spring + 1/3 June (mid summer). K should be applied in winter (KCL). Ca is applied via leaf sprays of CaCl_2 (0.75%) late season. Micro nutrients are applied via leaf sprays early season. Nutrients should be applied in the tree row, and Fertigation is used a lot.

21 Lecture 23: Stresses and effects on quality - 10/10-2025

Torben Toldam-Andersen, KU-PLEN

This lecture addresses stresses and quality in fruit and berry crops, covering the use of water stress, damage from low and high temperature stress, prevention of damage, and damage caused by wind and biological stresses.

Low temperature stresses include freezing damage to berries (braun and dry out) and freezing in open flowers. Damage from frost (freeze) protection with water is based on the principles of high thermal mass of water and latent heat. Latent heat is the energy transferred during a phase change. The phase change from liquid water to ice transfers 80 cal of latent heat, which is crucial for frost protection. In comparison, 1 cal (4.17 Joule) is required to increase the temperature of 1 g of water by 1°C, and 600 cal of latent heat is transferred during a phase change from vapour to liquid water. Frost damage, post bloom and Winter damage are also covered.

The water vapour content of the air is important: a high content results in a slow temperature drop during the night, while a low content results in a quick temperature drop. The location near the coast is important for protection against frost damage. In spring, the minimum temperature near the coast is higher and the maximum is lower, which results in delayed Budbreak and lowered frost risk. Data for 'Red Boskoop' apples (1975-2022) indicates

that the beginning of flowering is now >3 weeks earlier than in 1975, corresponding to a temperature rise of >1.5 K since 1975. This makes frost protection by irrigation increasingly important. Frost protection methods include Overhead sprinklers and Water tubes with micro sprinklers at each tree, as well as using Heaters/burners. Radiative frost occurs on clear and calm nights and is very local. Techniques to mitigate frost include using Wind mixers (e.g., in Washington State, US) and sand cover (e.g., Soil protection in China).

High temperature and water stresses include fruit cracking due to water problems. This has been observed in Washington, US and Ullensvang, Norway. Sunburn damage results from too much sun. Hail, especially when combined with strong wind, can be very harmful. Hail can also severely damage the shoots. Rain, hail and sun protection methods are available.

Wind stress/damage requires the establishment of shelters. Pest and disease stresses lead to various negative outcomes:

1. Reduce leaf carbon assimilation and export
2. Deformities of fruits (spots, cracks, splitting)
3. Less attractive
4. Lower quality
5. Lower yield
6. Fruit rot
7. Bad taste
8. Lower storability

The shoots can also be severely damaged.

22 Lecture 28: Potentials in fruit and berries for fruit wine production in Denmark - 10/10-2025

Torben Toldam-Andersen, KU-PLEN

This lecture explores the Potentials for (Danish) fruit wines through the 'NATVIN' Project, focusing on Apple cultivars, the content of apple juice, the importance of concentration, wine potentials, the process (How to do it), Quality impact, and the Benefits compared to alternatives to cryo concentration. For ordinary juice containing 12% Brix, the composition is typically 120 g/L sugar, 8 g acid, and 2 g sorbitol, resulting in 110 g Fermentable Sugars (FS). This concentration allows for a yield of 52 g alcohol or 6,5% vol Alc.. Wine acidity levels dictate the requirement for malolactic fermentation (MLF), defined as the conversion of malic acid to lactic acid by bacteria's (*Oenococcus oeni*). Moderate acid levels facilitate dry wines without MLF, while high levels necessitate MLF or wines with residual sugars (RS).

The implementation of cryo concentration expands wine potentials, increasing 12% Brix to 20% brix. This results in sugar content rising from 110 to 180-185 g/L, achieving 10,8% vol alc.. Acid levels concurrently rise from 5-8 g/L to 10-13 g/L, often requiring some MLF. Sparkling wines require Bottle Fermentation of added sugar to 4-5 bar, adding +1-1,2% vol alc., thereby reaching 12% total alcohol. Still wines made from 22% brix can reach 12% vol alc., generally requiring MLF. High Concentrate wines achieve 30-35% Brix, suitable for Sweet dessert wines, 'ice wines', although high acidity is a challenge, potentially exceeding >20 g/L.

The cryo concentration process involves freezing the juice, then thawing it slowly to yield concentrated juice. The resulting ice holds 2-5% Brix. Thawing often utilizes 1000 L tanks, where juice from multiple tanks is collected

into one. This method can result in an aroma explosion, with +40 aromas increased with more than 3x from control juice wines to high concentrated wines. Specific volatile increases include Dodecanoic acid, ethyl ester (314x) and Decanoic acid, methyl ester (178x). Only Ethyl lactate was observed to disappear, reduced 24x. Fufural content is also monitored, raising questions regarding Maillard reactions and/or caramellisation. Alternatives, such as Membrane filtration techniques (Reverse osmosis, Cross flow filtration), are problematic for cloudy juice due to high pectin content, necessitating viscosity reduction, and strong filtered juice may be difficult to ferment.

Berries pose unique challenges due to relative low in sugar and high or very high acid content (20-35 g/L), primarily citric and malic acid. Since these acids only form soluble salts, chemical reduction is impossible. Biological reduction via MLF is risky due to potential off flavour (Diacetyl)?, compounded by low or very low pH. Berry juice also exhibits very characteristic aromas, high phenolic content and often high in colour, and lots of pectin. Solutions involve Blends with water or a base of apple or pear. High acid species include:

1. Sour cherry 'Stevnsbær'
2. Black currant
3. Red and White Currant
4. Gooseberries
5. Rhubarb
6. Black berries

Potential techniques and styles include Sparkling, Liquor (juice+alcohol), Fortified wines, Ice wines, and Still wines. The average soluble solids in 114 Gooseberry cv's is 11,8%.

23 Lecture 30: Growing of wild berries - 20/10-2025

Martin Jensen, senior scientist, martin.jensen@food.au.dk

This lecture addresses the domestication of European Blueberry (*Vaccinium myrtillus* L.) for a new berry crop, focusing on wild fruits and the plant's morphology. The European Blueberry is an erect, woody, rhizomatous shrub that grows to a height of 5-90 cm. It exhibits tiller growth from the rhizome in a 3-angled axis, developing abundant annual shoots in a complex structure. Leaves are glabrous acute, 1-3 cm long, and flowers occur singly or in pairs in axil leaves, reaching 4-7 mm in diameter with a pink corolla. The plant forms patches through rhizome growth, with aerial shoots reaching 20-30 cm. Native wild populations of *Vaccinium myrtillus* are present in all regions of Denmark.

Wild harvested bilberries in Sweden are part of a larger yield in Nordic countries, estimated at 500 million kg per year, though only 5-8% is exploited. Traditional berry picking is often for personal consumption and local markets, involving a small and fragmented industry that exports to East Asia, with a price typically 20-30 DKK/kg. Yield in nature ranges from 0-450 kg/ha. Hand picking often employs manual rakes, with 3-4000 Asian people flying in annually to pick. For comparison, wild harvesting of lowbush blueberry (*Vaccinium Angustifolium*) in Canada, conducted as semi-cultivation on 2x50.000 ha of wild plants with a two-year production cycle, yields 99 million kg of berries per year.

European blueberry is of interest due to its status as a 'Superfruit', attractive taste, and high price. Potential health effects include lowering cholesterol and blood pressure, improving cognitive performance, and exhibiting antibacterial and vision benefits. It possesses a higher concentration of antioxidants than highbush blueberries. Currently, there is no commercial orchard production, relying only on 'natural collection'. Manual harvesting leads to high costs, limiting production and supply, and is deemed non-sustainable. Market demand is increasing

globally, and there are no reported cultivars or breeding/selection efforts done. Orchard production using selected cultivars and mechanical harvesting could create new product options due to lower costs, leveraging the fact that it is a native Danish species with available climate-adapted genetic resources.

The domestication of European Blueberry requires overcoming critical barriers:

1. Significant improved berry yield/plant is needed (selection and breeding), as maximum harvest in nature is too little (around 450 kg to 1 ton/ha).
2. Significantly more efficient and non-expensive methods of vegetative propagation are required, as past success has been minimal.
3. Development of a sustainable orchard concept/design adapted to mechanical harvesting is necessary.
4. Development of efficient mechanical harvesting is critical to reduce the cost of harvest, as Canadian solutions may not be adaptable.

The domestication project involved collecting clones, establishing a common garden trial of >100 genotypes, characterizing these genotypes for plant growth performance, berry yield, and berry quality. The established collection includes 113 clones from 40 populations across two sites, used for gene conservation, clonal comparison, and as a source of superior genetics.

Studies on fertilizer composition established that liquid fertilizers with N values of 200 and 400 ppm N damage and kill plants after 1-3 months. Optimal fertilizer concentration is 50-100 ppm N (EC = 0,3-0,6). 100 ppm N is optimal but only slightly better than 50 ppm in young seedlings, while 50 ppm was found better in cuttings. 25 ppm N suggests P deficiency (red leaf colour). A liquid fertilizer composition was developed for continuous use, using collected rainwater for irrigation, characterized by low levels of Ca, Mg, and Fe.

Berry quality analysis of 53 genotypes in 2016 showed mean total soluble solids of 9.1°Brix, total titratable acid of 0.895 g citric acid eqv./100 g, and a pH of 3.14. Total anthocyanins averaged 330.4 mg CGE*/100 g. Superior clones identified in 2016 achieved high berry production, with the top clone yielding 115 g/plant/year, leading to a potential theoretical yield of 16.56 tons/ha/year (assuming 16 plants/m²). The average yield of all 106 clones was 3.49 tons/ha/year. Observations 11 years after planting showed top genotypes yielding 90-150 g/plant, equaling a potential yield of 8-13 tons of berries/ha (assuming 9 plants/m²). The relationship between mean berry production and the mean weight of the 10 largest berries showed a positive correlation ($R^2=0,3553$). The research has demonstrated efficient and low-tech methods for cutting propagation, a liquid fertilizer composition, and characterized over 100 genotypes, pointing to possible superior genotypes for cultivation and further breeding.

24 Lecture 31: Bioactive compounds in fruit and berries - 20/10-2025

Martin Jensen, senior scientist, martin.jensen@food.au.dk

This lecture examines Bioactivity, defined by the effects or possible effects in humans from oral intake of compounds, relating to maintaining health, preventing disease, or resulting in a direct effect in the primary body or indirectly through a prebiotic or antibacterial effect. Regulatory categories include Medicine, Plant medicinal compounds, Functional foods, Supplemental foods, Food, and Novel Foods. Health claims, which must be based on scientific evidence evaluated by EFSA, include 'Function Health Claims' (related to growth, psychological functions, slimming), 'Risk Reduction Claims' (e.g., plant stanol esters reducing blood cholesterol), and 'Health Claims referring to children's development'. Documentation supporting health claims ranges from epidemiological evidence and in vitro tests to human clinical intervention tests, specifically recommending double blinded -

placebo controlled experiments for confirmation.

Examples of authorized EFSA claims include Pectins (reduction of blood glucose rise and maintenance of normal blood cholesterol) and Melatonin (alleviation of subjective feelings of jet lag and reduction of time taken to fall asleep, e.g., in sour cherry). Non-authorized claims include antioxidants from pomegranate juice and anthocyanins from elderberries.

Fruit and berries contain compounds with potential health effects, categorized as:

1. Anti-bacterial, Anti-viral, and Prebiotic effects.
2. Anti-oxidant, Anti inflammation, Anti- diabetic effects.
3. Protective against CVD - cardio vascular diseases (e.g., LDL cholesterol lowering, blood pressure lowering).
4. Anti-carcinogenic, Anti-mutagenic, Anti-toxic, detoxifying effects.

Estimated plant phenolics exceed 8000 polyphenolic compounds, including over 4000 flavonoids. Antioxidant capacity (FRAP, mmol/100 g) varies greatly, with Dog rose scoring 39.46 and Apple (Gold del) scoring 0.29. Genetic factors (cultivars), maturity, climate variation, cultivation methods, processing, and storage all cause variation in the concentration of single compounds, such as anthocyanins.

Bioavailability studies indicate that anthocyanins and large polymeric tannins generally show low uptake and low concentrations in blood. Anthocyanin metabolites reach maximum concentration after 1-2 hours and are depleted after approximately 10-12 hours. Clinical trials on single fruit species are few, often showing limited effects due to too low concentrations of active compounds in raw fruit. Research on Aronia suggests extracts or concentrates may function as functional food for disorders related to oxidative stress, but more confirmatory clinical trials are needed. A mechanism for cholesterol lowering is the ability of fruit and berries to bind and excrete bile acids in the intestines. Therapeutic effects may be achieved through additive effects in portfolio diets. Specific compounds like Galactolipid (GOPO) and two triterpene acids from Rose hip fruit are associated with pain reduction in osteoarthritis, as GOPO partly inhibits chemotaxis of human peripheral blood PMN's. Future research requires rigorous experiments, focusing on understanding compound metabolizing in the gastro-intestinal tract and modulation by bacterial flora.

25 Lecture 32: Water management and water stress - 24/10-2025

Torben Toldam-Andersen, KU-PLEN

Water is the most abundant constituent of plant tissues and functions as an excellent solvent for salts, other solutes, and gases. The majority of absorbed water is lost to the surrounding air via transpiration. Insufficient water limits horticultural and agricultural production in many areas. Future trends suggest an increase in extreme weather situations, such as intense rain events or droughts. Producers must therefore regulate their irrigation strategy based on crop needs and weather situations. Climate change is projected to increase irrigation needs; for instance, by the 2050s, eastern, southern, and central England are predicted to have greater irrigation needs than those currently experienced anywhere else in England.

Water flow through the plant (soil → plant → atmosphere) is driven by a gradient in water potential (ψ). Water potential is defined as the energy state of water in the environment, which governs water availability. The total water potential is calculated as $\psi = \psi_p + \psi_s + \psi_g + \psi_m$, where ψ_p is pressure or hydrostatic potential (turgor), ψ_s is osmotic potential (binding to solutes), ψ_g is gravitational potential, and ψ_m is matric potential (binding to

surfaces). In soil, the matric and osmotic potentials are the main components, whereas in plants, hydrostatic/turgor and osmotic potentials dominate. Net photosynthesis is correlated with stomatal conductance in apple leaves under varying water stress levels.

When the soil dries, the plant initiates physiological reactions. Moderate soil drying results in a slight decrease in root water potential, an increase in ABA (abscisic acid) in roots, and increased root growth. Subsequently, ABA and pH increase in the xylem, while nitrate decreases. The leaf water potential is kept stable, although stomata gas-exchange and leaf expansion decrease, ABA in the leaf increases slightly, and nitrogen in the leaf decreases slightly. Severe soil drying causes a strong decrease in root water potential, strong increases in root ABA, and strong root growth. The leaf water potential decreases, stomata gas-exchange (including A_{max}) decreases strongly, leaf expansion decreases, and nitrogen in the leaf decreases. Water stress affects vegetative development (root and shoot growth), flower bud formation (impacting the next year's crop), yield, and quality.

Plant water use and consumption depend on various factors:

1. Water reserves, which are determined by soil type.
2. Transpiration and evaporation, which are determined by temperature, wind, and humidity.
3. Leaf area, which depends on plant species, growing system, plant age, and plant shape.
4. Root size, which depends on plant species, growing system, plant age, and rootstock.

Cropping influences physiology in apple, leading to higher values compared to uncropped controls for photosynthetic intensity (100 vs. 140 - 210) and transpiration intensity (100 vs. 150 - 300). The water use coefficient measures L water used/m² leaf area/mm water evapotranspiration, and water use efficiency measures kg fruit/L water used.

Regulated water stress can be utilized to benefit yield and quality in fruit trees and vine production. Sensitivity varies greatly across species and cultivars. Shoot growth stops when new leaf development is reduced by water stress (increasing negative potential), which effectively controls shoot growth and induces terminal bud formation. Growers must control the water status to remain within the 'window' between the effect on shoot growth and the effect on photosynthesis to maintain dry matter production. Mild stress stimulates root growth, while strong stress reduces roots, particularly near the soil surface. Competition for water can be utilized, as green cover evaporates 50-90 mm/month (about 2-3 mm/day) compared to bare soil which evaporates 25-35 mm/month (about 1 mm/day), and covering with straw can save approximately 60 mm in a season.

Measurement techniques include Tensiometers, which use a porous ceramic cup to reach equilibrium with the soil solution's water potential, and Time Domain Reflectometry (TDR), which measures volumetric soil water content using electromagnetic waves. Water management strategies are applied to crops such as Apple and pears, Strawberry, Stone fruit, and Bush and cane fruits (e.g., Black currant and raspberry), sometimes in combination with root pruning.

Water stress reduces fruit set, berry size, and inflorescence initiation and development. The period most sensitive to stress is the first few weeks after flowering, where berry enlargement can be permanently limited. Just before ripening ('veraison'), there is little effect on fruit expansion. During ripening, mild stress enhances sugar and color content and reduces acid, promoting ripening. Strong stress during ripening can delay and reduce sugar and color. Reduced shoot growth limits competition between vegetative sinks and fruit development.

26 Lecture 33: Vitamins and colours in fruit and berries - 24/10-2025

Torben Toldam-Andersen, KU-PLN

The internal quality of fruits, specifically concerning vitamins and colour, is influenced by climate effects and growing techniques. Fruit colours are derived from three main compound classes: Green colour is attributed to Chlorophyll a and b; Yellow colours originate from Carotenoids, including Carotins ($C_{40}H_{56}$) like Beta-carotin and Lutein, and Xanthophylls ($C_{40}H_{56}O_2$) such as Violaxanthin and Neoxanthin; and Red colour is due to Anthocyanins. Seasonal changes affect the concentration of the four main carotenoids in 'Golden del.' apple.

Anthocyanins are the dominant red colours found in berries. Based on data from Kaack 1975, the dominant anthocyanins include:

1. Black currant: Cyanidin-3-glucosid, Delphinidin-3-glucosid
2. Elderberry: Cyanidin-3-glucosid, Cyanidin-3-sambubiosid
3. Strawberry: Cyanidin-3-glucosid, Pelargonidin-3-glucosid
4. Cherry: Cyanidin-3-glucosid, Cyanidin-3-sophorosid

The synthesis of Anthocyanins begins with green leaves and light producing carbohydrates. Cyanin (anthocyanidin) is produced in the presence of light, and adding galactose (sugar) completes the synthesis of anthocyanin. This process involves the Pentose phosphate pathway, Shikimic acid pathway, Glycolysis, and Malonacid pathway, with glycosidation as the final step.

Colour development is regulated by factors including Fruit/leaf ratio (light indirect), Nitrogen, Light (specific via PAL), and Temperature. The ratio of Kg fruit/Kg leaves or Number of fruits/kg leaves influences the yellow and green colour of 'Golden delicious' apple. Temperature effects are observed during early and late stages of ripening. In unripe fruit, Light combined with warm temperature increases fruit growth and sugar accumulation. In mature fruit, red colour is enhanced by light and warm temperature, as well as by light and cold (night) temperature. Climate affects red colour development, demonstrated by treatments of Control, 50% shade, and Warm night conditions applied to Jonagold apples from mid-August to October. Post-harvest colour development is also influenced by artificial light exposure on bagged fruit. In Black currant, high Nitrogen (N) levels increased the concentration and content per fruit of chlorophyll and carotens. Leaf removal increased the concentration of chlorophyll and carotens, but decreased the content per fruit due to resulting smaller fruits. Anthocyanin concentration was affected only to a limited degree by N level or leaf removal, although fruits in the top of the canopy exhibited higher anthocyanin content compared to berries at the basis, which suggests the importance of light conditions and assimilate availability.

Vitamin C comprises Ascorbic acid plus De-hydro-ascorbic acid. Vitamin C content varies significantly across species in mg/100 g Fresh Weight (FW), ranging from Apple (5-10) and Pear/Grape (5) to Orange (50-60), Black currant (180), and Rose hip (400-2000). Vitamin C is formed before the ripening phase, with concentrations affected by ripening and storage effects, Light (sun/shade, south/north, peel), and other factors like fertilizers/nutrients, often through light or developmental degree. Vitamin C degrades very fast. The distribution of Vitamin C within an apple shows highest concentration in the Epidermis (70 mg/100 g fw), declining significantly in the inner flesh (6 mg/100 g fw) and core (5 mg/100 g fw), leading to the conclusion: do not peel an apple. Light exposure also affects content, with the south side of an apple fruit having 34 mg/100g fw compared to 20 mg/100g fw on the north side. The effect of development on Vitamin C is varied; it remains stable in Strawberry (e.g., 50→54 mg/100 g fw from unripe to ripe) and Apple (e.g., 10→13), but decreases in Black currant (219→140) and Sour Cherry

(20→12). Gooseberry cultivars show an average of 30.9 mg/100 g fresh weight. Storage conditions (Controlled Atmosphere, Ventilated, Cold Storage, or room temperature) also impact Vitamin C levels.

27 Lecture 34: Fruit harvest and postharvest - 27/10-2025

Torben Toldam-Andersen, KU-PLEN

Fruit development involves specific phases of growth and the role of ethylene, leading up to harvest. Harvest time represents a crucial compromise between optimizing internal quality and maximizing storability. Internal quality is assessed via chemical changes, while external qualities include size and colour. Seasonality significantly impacts fruit characteristics. In the early season (e.g., strawberries), a cool climate results in low sugar and high acid content, and phenology favors the first and largest fruits, with cultivar differences being notable. The main season is often warm and dry, accelerating development, which increases the danger of harvesting too late and resulting in overripe fruit. The late season (e.g., apples) is cool and humid, characterized by lack of light, and involves the last and smallest fruits, alongside increasing risks of pests, diseases, and fruit rot, impacting storage products.

Key harvest criteria are essential for balancing the goals of high eating quality and maximum storability. Important parameters include fruit size, colour change, firmness, removal force, sugar/acid ratio, starch content, aroma, and the levels of ethylene, O₂, and CO₂. The Streif index is calculated as $\text{firmness (kg/cm}^2\text{)} / \text{sugar (\%)} \times \text{starch index (0-10)}$ and is dependent on both cultivar and climate. Cultivar differences are highly significant, especially regarding firmness in pears, which may change very fast. Prediction models, such as those relying on the temperature sum in 60 days after flowering for apples or the first 40-50 days determining phenology for cherries (Vitrup Christensen, 1973), aid in determining optimal harvest dates. For black currant, as fruits do not ripen uniformly, the harvest time is a compromise reached when some fruits begin to dry out and drop; machine harvest requires care to avoid damaging shoots and leaves. The optimal harvest date for storage can be related to the development of butylacetat (butyl etanoate), which is the most abundant aroma in apple juice.

Harvest and post-harvest losses can be very high, reaching 20% in Gross sale + Detail, and 20% for the consumer. Fruits are heavy, water-containing, alive, and undergo respiration and senescence, giving them short keep ability. They are fragile and sensitive to bruising and wounding, which causes ethylene release, increased water loss, and infection risks. High losses stem from damaged/reduced appearance, increased respiration and water loss, and infections. Prevention methods include careful harvest, appropriate boxing, and managing whether the stalk is included. Mechanical harvest methods include platforms for apples/pears, machinery for industry crops (e.g., 'Stevnsbær'), cane fruit harvesters, and portal harvesters for grapes in adapted growing systems.

Post-harvest physiology is strongly governed by temperature. Higher respiration equates to shorter shelf life. The Q10 ratio, which measures the respiration ratio at two temperatures 10°C apart, is typically 2-3, but can be 3-4 for fruits, emphasizing the dramatic effect of temperature. Fast cooling is necessary; for instance, strawberry Q10 \approx 4, meaning shelf life drops from 7-8 days at 0-2°C to about 2 days at 10°C. A delay in cooling of 6 hours can result in 50% higher water loss and loss of C-Vitamin and sugar. Fruits are categorized by ripening patterns:

1. Climacteric: Respiration and ethylene production increase (e.g., Apple, Pear, Banana).
2. Non-climacteric: Respiration gradually falls during development, indicating non-ethylene sensitive ripening (e.g., Cherry, Strawberry, Citrus).

To minimize drying out, fruits should not lose more than max 2-3% of water. Water deficit is the primary factor, heavily dependent on temperature. Water loss occurs 30% via lenticels and 70% via the cuticle. Minimization strategies include cooling, wrapping/foils, waxing, and the use of humidifiers in stores.

SmartFresh, a commercial brand of 1-MCP (1-Methylcyclopropene, C_4H_6), is structurally related to ethylene and functions as a plant regulator to inhibit ethylene action by binding to its receptors. 1-MCP effects conclusions show that early treatment (at harvest) strongly reduces aroma development, while later treatment (after light exposure) only results in partial effects. 1-MCP maintains higher firmness. Furthermore, treatment applied before light exposure reduces the fruit's ability to develop red over colour. The application time significantly affects the concentration of specific aroma compounds like propanol and ethyl acetate in pigeon apples.

Chapter 2

Literature résumés

This section of the course notes is designed to streamline access to the key findings from each reading material (RM), providing a concise and accessible overview of essential information. Created through experimentation with various AI platforms, this chapter also serves to enhance prompt engineering skills, exploring diverse methods of note-taking for maximum efficiency and clarity. The procedures for creating these summaries have varied, but all methods share a common approach: each RM has been fully read, with summaries and notes prepared after completing each respective subsection. By using these AI-co-op'ed approaches, these notes aim to be both a reliable reference and a resource for continuous improvement in capturing complex microbiology concepts.

1 RM for L02 - Fruit Yield and Quality. Components and Determinants

Fruit cultivation will normally be intended to maximize fruit yield and fruit quality [1]. The available tools for achieving this are variety selection (genetic selection) and choice of cultivation techniques, the physiological basis of which is treated in subsequent chapters. Maximizing fruit yield, often expressed as tonnes (or hekto litre in case of wine or juice) per hectares (ha), requires no further definition. Fruit yield is ultimately determined by fruit number per hectares and fruit size, understood as a product of several components. While component weights vary across fruit species, the main components can be summarized as shown in Table 1 and illustrated for cherry in Figure 1 [1]. Determinants define the amount or degree of development of each component within the genetic potential. Plant number per ha is primarily determined by the growing technique, mainly the selected planting system. The cultivation system may also determine factors such as the number of shots in raspberries and the number of rooted runners in strawberries. Other determinants are primarily physiological factors associated with growth and development, although they may also be influenced to some extent by growing technical interventions. The first main section of the course includes a review of these physiological determinants, serving as a basis for explaining and understanding the effects of different cultivation technical interventions.

Maximizing fruit quality is also a primary target in growing, although fruit quality cannot be defined nearly as unambiguous as fruit yield. Fruit quality includes a variety of components that vary with fruit type and fruit use, involving different weightings in different contexts. Fruit size is an important quality component, especially for table fruit. Colour and other factors related to fruit appearance are also important, particularly those factors that are part of EU standards for the grading of the various fruit species. These components are of specific interest to fruit growers because they contribute to price differences between different grades. Components attached to fruit taste and enjoyment value—e.g., sugar, acid, and aromatics content—are also important and are expected to become

even more important in the future. For industry fruit, such as sour cherries, emphasis is placed on juice colour and acidity, and for blackcurrant, on vitamin C content, despite there being no real price differences (quality premium). The concept of fruit quality can also include aspects such as the content of foreign substances like residuals from spraying of chemicals, which are of increasing importance in the complex of fruit quality. Societal priorities for health and environmental concerns, etc., also gain increasing importance in a growing technical context, and these factors are discussed in relevant, growing technical lectures.

Determinants such as flower development, and particularly fruit growth and fruit development, affect the quality components. Since physiological and growing technical factors may have opposite effects on yield and fruit quality components in some cases, the fruit grower's task is to seek the best overall compromise.

The components determining Yield per hectare and fruit quality, along with their primary determinants, follow a hierarchical path, as outlined in Table 1 [1]:

1. Number of plants / ha is determined by \leftarrow Planting System, and \leftarrow Growing System (Rubus, strawberry).
2. Bud Number / plant, bud type is influenced by \leftarrow Plant size, structure, \leftarrow Elongation growth, shot type development, and \leftarrow Bud Development.
3. Number of inflorescences / bud is determined by \leftarrow Flower initiation.
4. Number of flowers/cluster is influenced by \leftarrow Flower initiation and \leftarrow Flower Development.
5. Flower quality is determined by Number of seed primordia / Flower and Position in cluster.
6. Initial fruit set / flower is influenced by \leftarrow Pollination, fertilization, initial setting.
7. Final / initial fruit set is affected by \leftarrow Fruit drop (June drop).
8. Fruit Size is determined by Number of seeds / fruits (\leftarrow Pollination) and Quantity fruit flesh/seed, and is influenced by \leftarrow Fruit Growth, Fruit Development.

2 RM for L03 - Buds and Bud Development

Buds are growth points which can develop into shoots, determining the plant's future extension growth (dimension), or develop flowers, thereby determining the cropping [2]. Bud number and bud type are important components influenced by plant size and structure, which determine the number and distribution of various shoot types, consequently influencing bud number and type at the start of the growing season. Terminal buds, and sometimes the upper side buds on last year's long shoots, may develop new annual shoots. Lateral buds are formed in the axils of the leaves as the stem grows, culminating in terminal bud formation, with this elongation growth increasing the number of lateral buds. Other lateral buds on long shoots and end buds on short shoots may remain as short shoots or spurs, completing their short terminal growth with a terminal bud. Short shoots may mature well-developed lateral buds (often developing flowers), typical of fruiting spurs on cherry. Sleeping eyes (buds) only break after a strong stimulus, such as pruning. Flower buds of apple and pear are mixed buds, containing leaves and flowers, plus a new small bud (bourse-shoot bud), meaning a flower bud is usually followed by a new growth point. Stone fruit, conversely, has 'simple' or 'naked' flower buds, containing only leaves or flowers, resulting in the loss of a growth point at that position upon flowering, which can lead to bare areas on the shoot, typical of some sour cherry cultivars.

Elongation growth, and its distribution, determine bud development during the growing season. Terminal buds on short shoots are formed early (June). Lateral buds on longer annual shoots are formed in line with leaf development, with terminal bud formation occurring from June until September, depending on growth intensity. Elongation growth and bud development are subject to correlative inhibition mechanisms, including apical dominance, which involves a polar transport of growth substances (auxins, gibberellins) from terminal buds that inhibits

lateral bud growth. Heavy crop load can strongly reduce shoot elongation growth and the number of lateral buds due to competition for assimilates. Terminal bud formation occurs earlier if the ratio is high, see equation 2.1:

$$\text{Root activity} = \frac{\text{Root mass} \times \text{Root activity}}{\text{Number of Growth Points in the Top}} \quad (2.1)$$

Mature trees tend to have earlier terminal bud formation (possibly June-July) than young trees (often not before September). Elongation growth is enhanced by increasing temperature and by increased nitrogen and water supply (see sections 19.2.1 and 20.2.3).

The formation of a terminal bud is an abrupt process where bud scales develop instead of leaves, forming a compressed shoot inside with primordials for transitional leaves, true leaves, and possibly flowers. Bud break in the same growing season (re-growth) may occur exceptionally where correlative inhibitions are broken sharply, such as by removing shoot tips or performing early summer pruning. This is only possible until a certain date; with apple, for example, up to approximately 1st August, after which buds enter endo-dormancy.

Buds enter the endo-dormant phase when they can no longer break due to strong influences on correlative inhibitions. Bud dormancy is lifted by a species- and variety-specific period of low temperature, termed 'chilling'. Efficient chilling temperature is usually calculated as temperatures below +7°. In the temperate climate zone, the buds go through a physiological period of dormancy where there is no visible growth, which, if not met, can cause poor development or bud fall. The duration of dormancy varies by species (Table 1-2 shows Black currant ending dormancy in December, Apple in March). The depth and length of dormancy can be influenced by growth conditions in summer and geographical location, with cool summers accelerating termination. Dormancy is controlled by growth regulating substances, with Absciscic acid (ABA) having the strongest relationship with dormancy, while Gibberellins (GA) and/or cytokinins are growth-promoting substances. In areas with too little chilling, cultivation can be enabled by initiating the process with drought, followed by artificial defoliation (e.g., spraying with 10% urea) and then applying a spray to interrupt dormancy, such as 0.1% active DNOC +4% emulsified crude oil at the first watering.

Woody plants propagated by seeds must reach a certain age before flowering, termed the juvenile period or phase. This period may be short (e.g., Rubus and Ribe species), or long (e.g., 4-5 years for apple, 6-8 years for pear). The period between the end of the juvenile phase and first natural flowering is the transitional period. The juvenile period is characterized by different leaf forms and, sometimes, the transformation of short shoots into woody thorns (e.g., pear). To shorten the juvenile period in apples, experiments have reduced the period to approximately a third of normal using continuous growth in greenhouse conditions with increased day length and CO₂ addition.

3 RM for L04 - Chapter 2: Flower bud formation, flower development and flowering

A proportion of the buds formed during the growing season develop into flower buds, a process that is a very important yield component [3]. Flower bud formation begins with flower initiation, which occurs when there are no inhibitory effects. The time of initiation is typically summer - late summer, occurring in the approximate order of: Ribes, stone fruit, pome fruit, strawberry, and Rubus. This initiation is followed by flower differentiation.

Flower bud formation is most detailed studied in apple (pome fruit), primarily because it is an important crop and because it is frequently a limiting factor, especially concerning alternation (every two-year cropping). Studies show that a bud must undergo development before initiation can occur. For the apple variety 'Cox's Orange', a

critical node number of about 20 must be formed inside the bud. If the total number of nodes reaches 20 or more, a flower bud is formed, with a flower primordial in the terminal position and others in the corners of bracts (usually six to seven flower buds in total). The earlier a bud is formed (like spur buds), the longer time it has to develop, making flower formation easier. If node development is too rapid, the bud may break again in the same year (re-growth), causing delays that inhibit flower formation. Microscopically, the first sign of initiation is an increase in width of the growing point from 12-15 to 18-25 cells. Different theories exist regarding the mechanism, often involving positive and negative-acting factors, such as the Carbon/Nitrogen (C/N) ratio, where starch accumulation promotes flower bud formation. Flower bud formation may be associated with a high auxin / gibberellin content. The Source / sink ratio and flower bud formation are positively correlated. A strong negative correlation between fruit number per tree (fruit / leaf ratio) and flower bud formation explains the phenomenon of alternation. Reduced fruit growth rate, resulting in smaller fruits, leads to very large flower bud formation. Increased photosynthetic activity due to CO₂ addition is accompanied by an increase in flower bud formation. Increased shoot growth generally correlates negatively with flower bud formation. However, on young trees, recent studies suggest that a positive correlation can be found between shoot growth and flower bud formation on annual shoots, particularly when nutrients and water are supplied by drip irrigation. This allows the buds on annual shoots to reach their lower critical node number (14-16).

In stone fruit, alternation is typically less pronounced than in apple, possibly due to fruit size-related sink strength. In 'Stevnsbær', most buds on short shoots bloom, and the negative correlation between shoot growth and flower bud formation is less strong, primarily affecting longer annual shoots. Flower initiation may depend on the rate of development rather than a critical node number.

In *Ribes*, flower bud formation requires relatively short days (<12-14 hours) and low temperature. There is no direct effect of fruit number on flower bud formation; however, good cropping reduces shoot growth, leading to fewer buds and thus fewer flowers per bush the following year.

In strawberry (short-day types), day length is the crucial factor, with initiation occurring under Danish conditions when day length falls below 11 to 14 hours. Low temperature also promotes initiation. Vegetative growth (e.g., runner formation) is promoted by long days and high temperatures.

Flower development determines flower quality, which relates to the ability of the flower to set fruit. The differentiation of flower parts follows a sequence:

1. Sepals and petals differentiate rather quickly after initiation.
2. Stamens and carpels differentiate next.
3. Anthers (pollen sacks) and ovules often only differentiate next spring.

In cherry ('Stevnsbær'), bud death and abortion of flower primordia can cause mortality rates between 30-80% of initial primordia. In strawberries, the inflorescence is a cyme. With increasing flower order (primary, secondary, etc.), development becomes less complete, leading to inferior quality, characterized mainly by a decreasing number of pistils (Achenes) per flower.

Once bud dormancy is over, flowering time is mainly determined by temperature. The duration of the flowering process is also controlled by temperature; hot conditions may reduce it to one week, while cooler years may extend it beyond two weeks.

4 RM for L05 - A model to predict the beginning of the pollen season

The aim of the present study was to test phenoclimatographic models, comprising the Utah phenoclimatology Chill Unit (CU) and ASYMCUR-Growing Degree Hour (GDH) sub-models, on the allergenic trees *Alnus*, *Ulmus*, and *Betula* in order to provide a method to predict the beginning of the pollen season [4]. This type of model relates environmental temperatures to rest completion and bud development. Flowering is a phenological event resulting from a long period of development, beginning with the initiation and differentiation of buds into flower and vegetative buds during the summer. Falling temperatures cause a gradual change into a phase of winter rest with little or no growth activity. After a period, which length apparently depends on the climate and plant species, the plant gradually reverts to a phase of active growth in the spring.

As the phenologic parameter, 14 years of pollen counts (1977 to 1990) from a Burkard Volumetric Spore Trap in Copenhagen were used. The observed dates for the beginning of the pollen seasons were defined from the pollen counts as a fixed percentage (2.5%) of total counts. The observed dates of first bloom and pollen counts deviate profoundly from one year to another, with *Alnus* deviating from the 30th December to the 1st April, *Ulmus* from the 21th February to the 2nd May, and *Betula* from 2nd April to the 9th May. The chilling requirement was expressed using Chill Units (CU), where one CU is defined as one hour at 6°C, the optimum Chill Unit temperature for fruit trees. CU calculations began at the first day yielding positive chilling values. The experimentally determined temperatures for the GDH function included an optimum temperature of 25°C and a critical temperature of 36°C. The base temperature for GDH was changed from 4°C to 2°C, giving a slightly better correlation.

The models used were:

1. A fixed day model, using only the GDH model with 1st January as fixed initiation point.
2. A CU/GDH model, with a fixed sum of Chill Unit requirement as the initiation point for subsequent GDH accumulation.
3. A dynamic CU/GDH model, based on a dynamic relationship between CU and GDH.

In the ordinary CU/GDH model, the statistically estimated requirements for the trees were: *Alnus*: CU = 1550, GDH = 200; *Ulmus*: CU = 1850, GDH = 700; and *Betula*: CU = 1900, GDH = 2446. The minimum standard deviation for *Betula* was obtained at a CU estimate of 1900, giving an average of 2446 GDH. The dynamic CU/GDH model, equation 2.2, adjusted the necessary GDH to budbreak according to the CU obtained, proposing a non-linear S-curve function:

$$GDH = C + \frac{(D - C)}{1 + \exp\left(\frac{CU - CU_0}{A}\right)} \quad (2.2)$$

When compared with observed dates, the dynamic model provided the best predictions: within 2-4 days for *Alnus*, 8-10 days for *Ulmus*, and 3-5 days for *Betula*. This deviation size is acceptable given the large temporal variation from year to year. It is concluded that the CU and GDH relationships defined for fruit trees are generally applicable and give a reasonable description of the growth processes of other trees. *Alnus* tends to be completely regulated by temperature, but other parameters like the photoperiod apparently must be involved for *Ulmus* and *Betula* if predictions are to be further improved. The involvement of a dynamic relationship, though complicating the model, tends to improve the simulation of gradual changes during dormancy and budbreak. Furthermore, the results indicate that frost damage, detected using an LT₅₀ submodel (e.g., in 1981 and 1988), might be an important factor that can strongly affect total pollen counts.

5 RM for L06 - Physiology of Temperate Zone Fruit Trees

Fruit set plays an important role in modern fruit production, as a large yield of fruit is only expected if conditions for pollination and fruit set are favorable. Fruit set is a complex series of physiological events where no single element can be limiting to the overall process [5]. This overall process begins with the transfer of viable pollen to a receptive stigma, followed by germination, pollen tube growth, nutrient supply, and the successful fertilization of the mature embryo sac, which must occur before subsequent growth of the embryo. The events involved in fruit set are grouped into three main categories: the flowering process itself; circumstances that influence pollination and fertilization; and conditions conducive to fruit set without fertilization. All processes connected with fertilization are influenced by regulators, nutrition, and rootstock type, all of which are controllable by the grower [5].

The biology of bloom describes the process in which sepals and petals slowly enlarge, and the stigma(s) and stamens are exposed [5]. Bloom is considered to begin when 12-15% of the flowers are open and ends when 95-100% of the flowers are open. Cultivars generally do not bloom at the same time, and ensuring effective pollinizers requires that the bloom period of the main variety and the pollinizer overlap sufficiently. The timing of flowering varies from year to year, regardless of the absolute date, due to the differential sensitivity of various stages of bloom development to temperature stimuli [5].

The majority of fruit trees are self-incompatible, defined by the inability of fertile hermaphrodite seed plants to produce zygotes after self-pollination. Self-pollen is usually genetically inhibited from germination or pollen tube growth, preventing fertilization. However, even in self-fruitful cultivars, cross-pollination usually sets heavier crops. Triploid cultivars contain more pollen, which is generally less viable and has less value as a pollinizer [5]. Temperature has a great influence on the amount of pollen produced, with long cold winters or cold temperatures in early spring leading to a reduction in the number and vitality of pollen grains [5]. Bees are considered the most important pollinator insect [5].

The concept of the Effective Pollination Period (EPP), introduced by Williams in 1966, describes the limited period immediately following flower opening during which fertilization is possible [5]. The EPP duration equals the longevity of the ovule minus the time required for pollen tubes to reach the embryo sac. Ovule longevity varies significantly by cultivar; for example, the ovule longevity of 'Delicious' apple was estimated at 5 days, compared with 7-8 days for 'Jonathan'. The growth rate of compatible or partially compatible pollen is dictated by temperature. Pollen tube growth responds to temperature in a linear fashion, and a temperature response index can estimate the time required for pollen tube growth, compounded based on daily mean temperatures [5].

For a good fruit set, three requirements are sequential: first, the development of a strong flower bud during the previous fall, requiring sufficient photosynthate and nitrogen supply; second, a certain temperature range during and soon after bloom to ensure good pollination, pollen tube growth, and fertilization; and third, a relatively high photosynthate supply for the young developing fruit after fertilization [5]. Failure to satisfy these factors results in a poor fruit set and early drop of the young fruit. Early shedding of fruit is a regular feature of fruit set, often appearing in four waves at 12-14 day intervals. The third and fourth drops, called the 'June drop', are more conspicuous due to the larger size of the dropping fruit and involve a complete abscission process that includes the formation of ethylene.

Parthenocarpic fruit set, where fruit sets and grows to full size without fertilization, is most widespread in pears among temperate zone fruits. Pear varieties can be classified into four groups based on their ability to set parthenocarpic fruit [5]:

1. Not parthenocarpic: 'Pap Pear'.
2. Parthenocarpic fruit is shed at the time of June drop: 'Hardy'.

3. Variably parthenocarpic: 'Bosc', 'Clapp Favorit', 'Diel', 'Madame du Pois', 'Oliver Serres', 'Bartlett'.
4. Consistently parthenocarpic: 'Arabiska', 'Hardenpont', 'Passe Crassane', 'Pringall'.

In most fruit, the seed is the source of Gibberellic acid (GA), which is sensitive to temperature, explaining the dependence of parthenocarpy on high temperature around bloom [5].

6 1st RM for L07 - Physiology of Temperate Zone Fruit Trees

The utilization of plant reactions to changes in the fruit/leaf ratio requires consideration of the growth phase of the fruits, which varies between cultivars. Vigor, growth form, and cluster structure are also factors to be taken into account.

In fruit science, an organ with the status of an importer, such as strong growing shoot tips or developing fruit, is termed a "**sink**," importing dry matter and energy. Conversely, exporting organs, such as fully developed leaves that produce more sugar through photosynthesis than they use, are termed a "**source**". The surplus sugar is exported to competing sinks throughout the plant. The main source capacity covering the grapes' import/sink needs is located in the bottom 2/3 of the foliage wall.

The development of berries is roughly divided into 3 phases [6]:

The effects of affecting the fruit/leaf ratio must take into account the growth phase of the fruits, which are roughly divided into 3 phases characterized by different developmental processes. Phase 1 is dominated by cellular divisions and strong berry growth, quickly increasing berry size. This phase determines both the number of berries per cluster (based on the percentage of fruit set) and the size potential of the berries; increased cell divisions result in larger potential berry size. Reducing the number of bunches during Phase 1 decreases the fruit/leaf ratio, thereby increasing available resources, including carbohydrates (from leaves) and minerals (root intake). The result is an increase in both the percentage of fruit set and the number of cells per fruit, with the effect being stronger the earlier the action is taken. Early fruit (cluster) thinning can increase fruit set if the prospect for fruit set is less than desired. Simultaneously, there is a "risk" that the size of the berries increases, potentially resulting in too dense clusters.

The availability of resources for fruit development can be supported or enhanced by tipping shoots that have reached the desired height on the foliage wall. Strong growing shoot tips are defined as "sinks" because they import dry matter and energy, as their new young leaves are unable to cover their own needs. Conversely, exporting organs, such as fully developed leaves that produce surplus sugar in photosynthesis, are "sources". When the shoot tip is removed, a competing sink is eliminated, and the plant sends more sugar resources downward to cluster development, which also has sink status. Tipping also eliminates the apical dominance of the shoot, encouraging side shoot growth, although a period exists when the resource surplus primarily benefits the development of young/new fruit clusters. In wine grape production, the goal is often concentrated grapes and a relatively large skin/fruit ratio (i.e., small berry size), so early cluster thinning is usually avoided.

For cultivars such as 'Solaris,' utilizing the ability of fruit growth activity to limit shoot growth is often preferred. Early fruit thinning in Phase 1 is generally done only if the shoot growth is less than desired (e.g., on young plants). If growth is strong, maximum fruit load is kept, and the amount of leaves may be reduced. Uneven growth rates can be evened by tipping vigorous shoots when they reach the desired height on the foliage wall, moving resources to strengthen weak shoots. Phase 1 typically occurs for approximately 5 weeks, from 'Sct. Hans' until the end of July.

Phase 2 involves decreasing cell division, dominant seed development, and acid levels reaching their maximum. Xylem transport ability decreases, and the plant/fruit development is most robust to stress influences, such as drought stress. If the fruit load is clearly higher than optimal for Phase 3 (maturation), a first coarse thinning (e.g., down to 2 clusters per shot) may be appropriate. Bunch size reductions can be considered in high-setting, high-risk varieties like 'Johanniter' or Pinot's, or in large-clustered cv's such as 'Bolero' and possibly 'Cabernet Cortis,' if the clusters have not yet closed up. This period is typically around the start to mid-August. Continuous trimming of significant summer side shoot growth is important. Cutting off a dense cluster tip may allow the grapes ample space, and it may be better to keep 2 such adjusted bunches per shot instead of 1 large and dense cluster.

Phase 3 is marked by color shift, progressing from green to more yellow shades. Berries grow in size through cellular expansion, resulting in decreased firmness and a doubling of grape size. If a first rough adjustment was not made in Phase 2, it can be done early in Phase 3. Managing this phase is an act of balance; weaker growth requires greater thinning, while stronger growth requires delaying thinning to use the fruit to slow growth. As foliage becomes denser, leaf thinning begins, focused on the grape zone, and side shoots are heavily controlled. Thinning later than 3 weeks before harvest will reduce the effect on sugar accumulation (higher brix value). The final fruit/leaf ratio should normally be established by the first week of September.

For varieties susceptible to bunch stem-necrosis, such as 'Rondo' and 'Cabernet Cortis,' gradual adjustments are particularly important to avoid/reduce shock treatments. Leaf removal usually begins early, focusing on inner leaves in the grape zone to create an "inner open box" while preserving the outer leaves as important sugar producers and for protection. Leaves are first removed on the east side in August to stimulate early morning drying. Gradually, the west side is opened to achieve the desired exposure of around 40-50% of the clusters. The main source capacity covering the grapes' import/sink needs is located in the main leaves in the bottom 2/3 of the foliage wall.

7 2nd RM for L07 - Manipulation of growth and development by plant bioregulators

The growth and development of deciduous fruit trees are traditionally regulated by practices such as choice of rootstock, pruning, training, fertilization, water supply, and adjustment of crop load by hand thinning. When these cultural practices are insufficient, Plant Bioregulators (PBRs) are applied. PBRs are natural or synthetic compounds that mimic or counteract naturally occurring hormones. Examples include the cytokinin BA, which promotes cell division and fruit growth, and AVG, which suppresses ethylene formation. PBR results are often variable due to a poor understanding of their mode of action, "carry over" effects in fruit trees, variation in environmental conditions affecting uptake and translocation, and the specific anatomy and metabolism of different crops and cultivars. Societal concern and high re-registration costs pose significant obstacles to the continued use and development of synthetic PBRs [7].

PBRs are generally applied as aqueous foliar sprays. A major constraint on effective delivery is penetration through the leaf surface, which is covered by the cuticular membrane (CM), a non-cellular lipoidal barrier. The effectiveness depends on the PBR's nature, concentration, formulation, droplet size, and carrier volume. Decreasing droplet size or increasing carrier volume per leaf generally increases uptake, translocation, and biological response. High-volume spray applications often provide greater consistency. Environmental factors greatly affect PBR uptake and response. For instance, uptake of NAA and NAAm in apple and pear leaves increased with temperature in the range 5°C to 35°C.

Regulation of Fruiting (Thinning)

Excessive fruit set must be reduced because fruits may not reach adequate size and quality due to insufficient leaf area per fruit. Overcropping also reduces or inhibits flower-bud formation, which leads to a biennial pattern of cropping. Hand thinning is the most accurate method, but chemical thinning greatly reduces cost and time.

To reduce excessive fruit set, several PBR strategies can be followed:

1. The initiation of floral primordia may be partially inhibited.
2. Flowers can be killed or prevented from setting fruits at the time of blossoming (chemical flower thinning).
3. Thinning may be postponed until fruit set can be judged and, if abundant, chemicals aimed at promoting the drop of young fruitlets can be used (chemical fruit thinning).

The third method (chemical fruit thinning) is usually preferred as the least risky strategy. However, a too long delay in thinning diminishes the desired effects on fruit growth and flower-bud formation. For 'Empire' apples, fruit weight increased significantly when hand thinning occurred at bloom, 10, or 20 days after bloom compared to 40 days after bloom.

Flower thinning is crucial for cultivars that set abundantly, as later fruitlet thinning may be ineffective. Two flower-thinning strategies exist:

1. Using caustic compounds, such as ATS, that damage or severely desiccate stigmas and styles to prevent effective pollination. These compounds may also act by stimulating wound-ethylene evolution.
2. Using chemicals, such as ethephon, that enhance ethylene formation in the flowers. Ethephon application at anthesis increases ovule senescence and reduces auxin basipetal transport, leading to enhanced flower abscission.

Flower thinning

Flower thinning success may advance fruit maturity. Thinning activity is promoted by cool, cloudy, wet periods preceding application, and is enhanced by high temperatures after application, which increases competition for assimilates between sinks (shoot tips and fruitlets). For long-term sustainability, the most promising thinning compounds are those that occur naturally (e.g., BA), degrade rapidly (e.g., ethephon), or disappear quickly (e.g., NAA, ATS).

8 RM for L08+09 - Chapter 5 Growth, Growth correlations and Assimilate turnover

Fruit growth is an important yield determinant and affects fruit quality. Fruits contain 80-90% water. Of the dry matter, at least 90% originates from assimilates formed in leaves by photosynthesis. In pome and stone fruit, the sugar-alcohol sorbitol is mainly formed and serves as the transport agent. Assimilate production and allocation form the basis for growth and fruit development. By incorporating radioactive CO₂ (¹⁴CO₂), one can track assimilate use [8].

Approximately 21% of the dry matter produced in leaves in autumn builds up storage nutrients, which are stored in the tree and utilized the following spring. A significant share, 18%, is stored in the roots. Storage nutrients include starch, sorbitol, and sugar. Fall and spring are often dominated by starch, while sorbitol and sugar proportions increase in winter. Most stored reserves (17 of 21%) disappear completely, assumed used for respiration

to generate energy for spring growth. Warm spring weather may lead to enhanced respiration and consumption of reserves, resulting in low carbohydrate content, reduced ovule development, lower flower quality, and reduced fruiting ability. Early autumn defoliation reduces carbohydrate storage, which subsequently reduces shoot growth the following year. High fruit load reduces root growth and stored carbohydrate reserves. This may reduce shoot growth and thickness growth early next summer and decrease the cell number per fruit. Storage materials play a role in other species; in strawberries, starch accumulates in the roots and is utilized during leaf and flower development in spring.

In the very early growth stages, storage material is widely used; flower buds and young shoot tips receive 50-65% of their building material from reserves, with the remainder from young leaves. New foliage is initially a net importer, but leaves become net exporters when they reach around half their final size. Vigorous-growing shoot tips are strong users of assimilates, dominating competition over young fruit. High young fruit drop is evident by particularly strong growth shoots. If shoot tips are removed, the assimilate demand of the shoot is temporarily reduced, and young fruitlet drop decreases. As shoot growth decreases and terminal growth is completed, only approx. 15% of what the leaves produce are retained in the shoot. Large quantities then support fruit growth, such as in Figure 5-2e where a fruit received approx. 35% of substance produced in leaves of annual shoots further out. Strong summer pruning (in August) that removes many annual shoot leaves can reduce fruit growth and dry matter content.

Although fruits are not strong competitors in the earliest developmental stages, their ability to attract assimilates (as strong sinks) becomes very large later. A fruit in July-August consumes approx. 80% of the assimilates formed in leaves on the same spur (Figure 5-2f). If there are many fruits, they can pull assimilates from other branches over distances of 1-1.5 m; in Figure 5-2g, 68% of assimilates formed in leaves on a branch without fruit were allocated to fruits on a branch with many fruits. In strongly growing apples, at least 95% of fruit carbohydrate consumption originates from leaf assimilates. Flowers and young fruits may contribute 15-33%, mostly at the 'tight cluster' stage.

The temporal increment of dry matter often results in an S-shaped curve (Figure 5-8). Spur leaves complete growth in June, followed by annual shoot leaves, shoots, small branches, main branches, stem, and roots, which often have their main growth in September-October. The strong import ability of fruits allows them to dominate competition for assimilates. At very high cropping levels, apple fruits can utilize nearly 70% of total dry matter production in a growing season. Vegetative growth is strongly inhibited by high fruit load. In young trees, compared to similar non-bearing trees, high fruit load may reduce:

1. Leaf growth to half.
2. Shoot growth to one-third to half.
3. Branch growth to a third.
4. Root growth increment even more.

The inhibitory effect is more powerful further from the fruit, correlating with the temporal coincidence of stem and root growth with the fruits' main growth period. To achieve strong vegetative growth in young trees, fruit should be removed as early as possible. The total photosynthetic production in young apple trees may be more than two times greater per leaf area unit in bearing than in non-bearing trees in July and August.

In blackcurrant, even with a much lower share of total dry matter accumulation in fruits compared to apples (Table 5-9), reduced leaf and shoot growth are found at a good cropping level. However, differences are much smaller on branches and especially roots compared to apple. This is likely due to the earlier fruit harvest (approx. August 1), which limits the fruit's effect on late-season growth organs like roots. Unlike apple, where the root/top ratio decreases with cropping, the opposite tends to be true for blackcurrant. The strong pull in assimilates exerted by

fruits causes a greater photosynthetic intensity and more rapid transport out of leaves. Assimilate retrieval from nearby leaves to growing fruit occurs with almost no loss, possibly due to CO₂ re-fixation within the fruit.

9 RM for L10 - Table Grapes in Cold Houses

This compendium note provides detailed information on growing vines in unheated glasshouses on a month-by-month basis, assuming the vines are well established. Maximum ventilation should be provided in January to keep buds dormant. In February, start maintaining a slightly higher temperature, opening ventilators near 7°C. Syringing with tepid water should only occur on sunny days, before midday, ensuring the house and vines are dry by evening. By March, the temperature can rise to about 10°C before admitting air. A moist atmosphere must be maintained by damping, sometimes twice a day on sunny days. Borders should be examined to a depth of 45 cm; if dry, water (30-40 mm), and repeat (10-20 mm) in a week, combined with manure water immediately after (never water manure on dry soil) [9].

Mildew is a common disorder. To prevent infection, spray with a fungicide or use sulphite smoke when shoots are 5 to 7 cm long. In early April, reduce young growths to two shoots per spur, retaining the terminal one and a strong shoot near the base for short spur pruning. Before the end of April, pinch young shoots at two to four leaves beyond the bunch. In May, gradually tie laterals down to the wires using a loop-tie (Fig. 3), preventing them from touching the glass. During flowering, maintain a fairly high temperature, keeping the top ventilator open day and night unless there is an exceptionally cold spell. Drier atmospheric conditions are desirable before midday for drying pollen. To pollinate, draw the hand gently down the bunches about midday, or tap cordons sharply around 9 a.m. for shy setters.

Immediately after berry set in June, the number of bunches must be reduced, retaining a small surplus. As berries swell, thinning should start in two stages using vine scissors, first removing all seedless berries and those pointing towards the centre of the bunch, retaining all tip berries. Overcropping must be avoided; a rough estimate for vines on the single cordon system is one bunch for every foot (30 cm) of cordon from the basal spur. A mulch of farmyard manure may be applied in June, ensuring ventilation is maintained to disperse fumes. In July, the berries undergo the first swelling (BBCH 77) followed by stoning (BBCH 77-79), a critical period where temperature should not fluctuate. After stoning, a second thinning is advised, suspending shoulders with raffia to maximize swelling (BBCH 80-89).

As berries approach maturity in August, gradually reduce atmospheric moisture. For the second swelling (BBCH 83), temperature can be increased by early closure of ventilators, accompanied by copious damping's. Scalding (discolored sunken patches) is caused by hot sun on moist tissues; ventilators must be opened slightly before the sun strikes the house. Ripening is categorized using BBCH stages, with total days for BBCH 70 to 89 ranging from 75 (very early) to 100 (very late).

Pest and Disease control includes managing Mealy bug with a 5% tar-oil wash when buds are dormant, or approved sprays during the season. Shankling is a condition caused by unhealthy roots or overcropping, requiring root defect correction, crop reduction, and maintaining pH and K/Mg balance. Deficiencies, such as Magnesium (yellowish orange discoloration), can be corrected by spraying 250g magnesium sulphate in 10L of water plus spreader.

By November, bunches should be removed, and maximum ventilation given to rest the vines; sub-lateral growth should be removed to ripen laterals. Pruning should occur immediately in December after leaf fall, cutting the lateral back to two buds to reduce bleeding. Cultivars tested at Pometet, such as those in the Interspecific resistant types (Group C), are ranked based on 25 culture technical parameters and 36 technological parameters. Very high ranked cultivars generally show high sugar content (score 7) \approx 19% brix and acid levels (score 5-7) between 6

to 6,8 g/L. Ventilation is crucial and should be managed using top ventilators first, avoiding drafts, and seldom requiring bottom ventilators except at flowering.

10 1st RM for L11-13 - CHAPTER 6 Fruit growth and fruit quality

Fruit growth determines fruit size and shape, which is an important yield and quality component. External and internal quality factors vary greatly between species. Botanically, fruit types include a pome (pear, fusion of ovary and receptacle), a drupe (peach, one-seeded from ovary), a berry (grape, multiseeded from single ovary), an aggregate fruit (blackberry, many ovaries of a single flower), and an epigynous or false berry (blueberry). Most fruit is high in water, often over 80%. Carbohydrates dominate the dry matter in most species, though nuts contain protein and fat. Fructose has the highest relative sweetness (173, with sucrose as 100). Malic acid or citric acid are most often the dominant organic acids [10].

At least 90% of the fruit's dry matter originates from assimilates formed in leaves via photosynthesis. In pome and stone fruit, the sugar-alcohol sorbitol is mainly formed and serves as the transport agent, while sucrose is the transport agent in most other species, including fruit bushes and strawberries. Approximately 21% of dry matter produced in leaves in autumn is stored as reserves (starch, sorbitol, sugar), with 18% stored in the roots. Storage reserves are crucial, as flower buds and young shoot tips receive 50-65% of their building material from them in early growth stages. Fruit growth is conceived to be affected by the assimilate level, which depends on the ratio of assimilate producing organs (sources) and utilizing organs (sinks). Source strength is the capacity to synthesize compounds for export, and sink-strength is the potential capacity for accumulation of metabolites.

Leaf/Fruit Ratio and Assimilate Turnover With increasing leaf/fruit-ratio, increased fruit growth, and increased concentration of total solids, soluble solids (sugar), and acid are observed in apples. Conversely, a larger fruit/leaf ratio causes a stronger "pull" in assimilates, which animates leaves to greater production (greater photosynthetic intensity) and faster transport, but results in a lower assimilate level and lower concentration of certain substances, such as fruit dry matter. The relationship between yield (y) and number of fruits (x) per hectare can be expressed as $y=kx/K+x$, potentially using k to express source activity and $1/K$ to express sink activity. High fruit load strongly inhibits vegetative growth in young trees, reducing leaf growth to half and shoot growth to one-third to half.

Source and Sink Activity

Increasing source activity, such as through more light or higher CO₂ concentration, leads to a greater "pressure" from the leaves, resulting in larger fruits and increased accumulation of soluble solids (sugar). Light is an important factor affecting source activity in the field. Increasing sink activity in the fruits, often affected by substances from the root (e.g., potassium), causes fruits to "pull" assimilates stronger, resulting in larger fruit size but a lower dry matter content. For example, a good potassium supply resulted in larger fruit (117 g/fruit) but lower soluble solids (15,1%) compared to fruits lacking potassium (98 g/fruit, 15,9% soluble solids). Cultivar differences in fruit growth rate potential, such as 'Graasten' being approximately 25% higher than 'Golden Delicious', are apparently due to genetic differences in fruit sink activity. Variation in Species

The importance of the leaf/fruit-ratio on fruit growth is less pronounced in small-fruited species. In plums, reduced fruit number increases fruit size and solids/sugar content. In small-fruited sour cherry ('Stevnsbær'), reducing fruit number gives no or only a small increase in size. In black currant, sink activity is the most important factor for fruit growth, and large fruits are negatively correlated with solids concentration. For strawberries, flower development

and quality are essential for berry size. Berry weight is a linear function of achene number per berry at a given achene density (Annex 6-23). Development of achenes (seeds) is important as they release growth substances (auxin) needed for berry growth. In strawberries, thinning effects on berry growth are small, and the impact of physiological and cultivation factors on sugar and acid content is very limited. The relative importance of key factors varies across species (Annex 6-6):

1. Leaf/fruit ratio shows a positive correlation (+) with Fruit size, Dry matter, Acid, and Yellow background color.
2. Source-activity (light) shows a positive correlation (+) with Fruit size, Dry matter, Acid, and Yellow background color.
3. Sink-activity shows a positive correlation (+) with Fruit size, but a negative correlation (-) with Dry matter (sugar) and Red over-color (apples).

11 2nd RM for L11-13 - CHAPTER 7 Fruit Development and fruit ripening

Fruit development is characterized by phases which gradually transition, resulting in an overall effect on the developed fruit that is quantitatively influenced by the source-sink ratio. Fruit development transforms into ripening, which is characterized by a decrease in size increment that eventually stops, as well as the synthesis of specific substances, such as colors and flavors. The fruit may be tree- or picking ripe (mature) for a longer period before becoming eating ripe (ripe), particularly with many apple and pear varieties, though berry species often unite these stages.

Size increment typically follows an S-shaped (sigmoidal) curve in apples, pears, and strawberries. Stone fruit growth follows a double S-shaped curve that can be divided into three stages:

1. Seed coat and endosperm develop.
2. Seed coat hardens (lignifies, or pit-hardening), and the embryo develops. Growth seems less powerful.
3. Growth is again rapid, and the flesh develops via cell expansion, increasing the proportion of intercellular space.

Black currants (inner fruits in the raceme) can also show a double S-shaped growth curve. Cell divisions are dominant during the first 10-20% of the fruit developmental period, equalling the first approximately 4 weeks after flowering in apple. Thereafter, the time is dominated by cell expansion, although in strawberries, cell divisions may occur over a major part of the growing period.

With fruit development, the attraction between cells decreases as changes shorten the molecules within the pectin fraction, decreasing fruit firmness. The content of solids generally increases, with soluble solids (sugar) rising particularly sharply at the end. Assimilates (at least 90% of dry matter) are transported mainly as the sugar-alcohol sorbitol in pome and stone fruit, and as sucrose in most other species, including fruit bushes and strawberries. In apples, total acidity decreases with time, partly because de novo synthesis mainly occurs early in the growing season (when sorbitol is largely transformed into acid, e.g., malic acid, by June 30), and partly because already formed acid is eventually metabolized during ripening. Sorbitol transported into the fruit later (after August 20) yields especially sugars, mostly fructose. Starch accumulates and subsequently decomposes in apples, contributing to the increase in total sugar content.

The development and maturation rate is influenced by the rate of turnover in each phase, largely affected by source-sink relationships, and the duration of each phase. Increased temperature promotes the metabolic turnover

rate and shortens the duration of individual phases, thus increasing development and maturation rate. In late apple and pear varieties, increased temperature leads to larger fruit (Annex 7-6) and a higher sugar content (Annex 7-7). For the sour cherry cultivar 'Stevnsbær', the number of days between full bloom and harvest can be calculated as $160 - (0.13 \times \sum T_{0-40})$, where $\sum T_{0-40}$ is the temperature sum in the first 40 days after full bloom.

Harvest criteria in apples and pears include fruit size, change in base color from green (chlorophyll decomposition) to yellow (carotenoids), formation of red over color (anthocyanins, which require light), pip color change (white to brown), formation of the abscission layer, and firmness. High sugar content, approximated by soluble solids (refractometer), and decreasing acidity are important quality factors. The degradation of starch, followed by iodine-potassium-iodine solution staining, is also used as a harvest criterion. Because a single criterion is too risky, an index is used, such as the equation 2.3 that shows the Streif index:

$$\text{Streif index} = \frac{\text{Fruit firmness (kg/cm}^2\text{)}}{\text{Refractometer (\%)} \times \text{starch value (0-10)}} \quad (2.3)$$

The index decreases during maturation. Optimal index values vary by variety, e.g., 'Elstar' 0.30 to 0.38. Fruits for fresh consumption are still picked by hand, while fruits for industry (e.g., sour cherries, black currants) are largely harvested mechanically.

12 RM for L13 - Growth and development in black currant (*Ribes nigrum*)

III. Seasonal changes in sugars, organic acids, chlorophyll and anthocyanins and their possible metabolic background

Potted black currant plants of 'Tenah' (one year old, 1992) and 'Ben Nevis' (three year old, 1993) were sampled weekly from shortly after anthesis until harvest to measure fruit growth as dry-matter increments and seasonal changes in soluble sugars, organic acids, chlorophyll, and anthocyanins. Fruit growth in *Ribes* may be divided into two major rapid growth phases separated by a relatively short transition period. The typical double sigmoid growth pattern includes a plateau between 25-38 d from fertilization. The first rapid growth period coincides with cell division and seed growth.

In the very early stage of fruit development, high concentrations of sugar, malic acid, and chlorophyll were observed. Sugar concentration was initially high, then dropped to a minimum about 3-4 weeks after full bloom. Sugars started to accumulate rapidly, followed by a declining rate during the last weeks of ripening. During ripening, sugars, especially glucose and fructose, accumulated. Fructose reached a higher level while sucrose attained much lower values.

For organic acids, the initial concentration of malic acid in young fruits was high, declining to a steady state level of about two thirds of the initial level. During ripening, malic acid concentration fell rapidly until harvest. Citric acid concentration increased rapidly along with fruit growth until harvest, resulting in citric acid dominating the acid content per fruit. The accumulation of citric acid was similar in both years/cultivars on a per fruit basis. The total acid content decreases because initial malic acid concentration is very low and malic acid dominates, initially accounting for 70-75% of total organic acids. The final malic/citric acid ratio was 1:6-10 at maturity.

The sugar/acid ratio was initially high (1.4-1.5), dropping dramatically to a minimum between 35 d after full bloom (0.42-0.45), before returning gradually to the initial level during the last growth phase. The total chlorophyll concentration increased during early development. On a per-fruit basis, chlorophyll content rose strongly until 35

d after full bloom and then remained constant. Anthocyanins, low in immature fruits, showed a very rapid increase during ripening.

Metabolically, immature green fruits have a high capacity to support their own growth and carbon requirement through photosynthesis and dark respiration. Photosynthesis may be supported by CO₂ fixation via PEPcarboxylase, leading to the formation of oxaloacetate (OAA) and subsequently malate. Citric acid accumulates strongly in both immature and ripening fruits. A close correlation was found between citrate content and water content per fruit ($R_2=0.97$). The large amount of citric acid suggests a high metabolic priority. The period of maximum sugar minimum concentration, spanning approximately 30-45 d from full bloom, may reflect a period of critical internal competition for resources.

13 1st RM for L15-16 - Advances in Fruit Aroma Volatile Research

Fruits produce a range of volatile compounds that form their characteristic aromas and contribute to flavor, which comprises sweetness, acidity, or bitterness (perception in mouth) and odor. Fruit volatile compounds are mainly comprised of esters, alcohols, aldehydes, ketones, lactones, terpenoids, and apocarotenoids. Aroma is a complex mixture whose composition is specific to species and often variety. Volatile esters often represent the major contribution in apple (*Malus domestica* Borkh.) and peach (*Prunus persica* L.). Many C₁₀ monoterpenes and C₁₅ sesquiterpenes compose the most abundant group, sometimes determining the characteristic aroma, such as the terpenoids S-linalool, limonene, valencene, and β -pinene in strawberry, koubo, and citrus. The volatile profiles of fruit are complex and vary depending on the cultivar, ripeness, pre- and postharvest environmental conditions. Volatiles are classified as primary (present in intact fruit tissue) or secondary (produced upon tissue disruption). The proportion of glycosidically bound volatiles is usually greater than that of free volatiles, making them an important potential source of flavor compounds [11].

Volatile profiles are highly species-specific. For example, more than 300 volatile molecules are reported in apples, with esters being the most abundant. Hexyl acetate, hexyl 2-methyl butanoate, hexyl hexanoate, hexyl butanoate, 2-methylbutyl acetate, and butyl acetate are prominent in 'Pink Lady' apples throughout maturation. In strawberry, the furanones 2,5-dimethyl-4-hydroxy-3(2H)-furanone (furanol) and its methyl derivative (mesifurane) are dominating aroma compounds. Esters, which are the most important group, cover 90% of the total volatiles in ripe strawberry fruit. Banana fruity top notes are primarily volatile esters, such as isoamyl acetate and isobutyl acetate. Terpenes, such as D,L-limonene and valencene, are the major class of compounds in citrus. Peach aroma is defined by C₆ aldehydes and alcohols (green-note) and lactones and esters (fruity aromas).

Factors affecting volatile composition include genetic makeup, maturity, environmental conditions, postharvest handling, and storage. Maturity is critical; immature fruits produce low quantities of volatiles and lose the capability of volatile production more readily during storage than mature fruit. C₆ aldehydes and alcohols are major contributors to the flavor of immature fruits, but their levels decrease drastically during ripening as furanone and ester production increases. Postharvest handling significantly influences volatiles. Storage temperature is fundamental. Storage under Controlled Atmosphere (CA) conditions (low O₂ and high CO₂) alters production, reducing the capacity to synthesize ethylene and aroma volatiles.

Volatile aroma compounds are biosynthesized via several pathways starting from lipids (fatty acids), amino acids, terpenoids, and carotenoids.

1. Fatty Acids Pathway: Fatty acids are major precursors for C₁ to C₂₀ straight-chain compounds. The β -

oxidation pathway forms volatiles in intact fruit, providing alcohols and acyl CoAs for ester formation by removing C₂ units. The Lipoxygenase (LOX) pathway produces saturated and unsaturated volatile C₆ and C₉ aldehydes and alcohols, typically when tissue is disrupted.

2. Amino Acid Pathway: Amino acids, such as alanine, valine, leucine, and isoleucine, are direct precursors. They undergo deamination or transamination to form α -keto acid, followed by reductions, oxidations, and/or esterifications to form aldehydes, acids, alcohols, and esters.
3. Terpenoids Pathway: Terpenoids are derived from the C₅ precursor isopentenyl diphosphate (IPP) and its isomer DMAPP. C₁₀ monoterpenes originate from geranyl diphosphate (GPP), formed in plastids via the MEP pathway. Terpene Synthases/Cyclases (TPSs) convert these precursors into diverse cyclic and acyclic compounds.
4. Carotenoid Pathway: Apocarotenoid derivatives are produced by the oxidative cleavage of carotenoids, catalyzed by Carotenoid Cleavage Dioxygenases (CCDs). Apocarotenoid volatiles are synthesized only at the latest stage of ripening.

14 2nd RM for L15-16 - Advances in Fruit Aroma Volatile Research

The formation of volatile aroma compounds is influenced by the ripening process of the fruit, including the effects of harvest date and storage conditions. A negative relationship exists between the fruit/leaf ratio and the availability of assimilates and sizes, as well as the concentration of total and soluble dry matter and acids. Fruit growth is conceived to be affected by the assimilate level, which depends on the ratio of assimilate producing organs (sources) and utilizing organs (sinks). The great import ability of the fruits means that they will dominate over other organs in the competition for assimilates when they are out of the early growth stages [12].

One-year-old apple trees of *Malus domestica* cv. Jonagored were divided into three fruit/leaf ratio groups: Low (130 fruit:leaf DM 10⁻³ kg/tree), Cropping median (172), and High (381). The study established that at a low fruit/leaf ratio, larger apples were produced, accompanied by higher concentrations of total dry matter, soluble solids, titrateable acids, and a lower firmness value, compared to apples from trees with a high fruit/leaf ratio. Fruit firmness decreased during the ripening period, and apples from trees with a high fruit/leaf ratio were softer than apples from trees with the low ratio at harvest.

Headspace Gas Chromatography (GC) analysis showed that aroma compounds consisted of approx. 20% esters, 71% alcohols, and 6% C₆ aldehydes. The total concentration of aroma compounds increased during the ripening period and was most pronounced at the lowest fruit/leaf ratio. Esters and alcohols were dominant, while C₆ aldehydes showed no significant difference between the fruit/leaf ratios. The higher aroma concentration associated with low fruit/leaf ratios is supported by the theory that greater availability of assimilates favors the accumulation of precursors for the synthesis of aroma compounds.

Ethylene concentration was measured during a 3-week ripening period after cold storage. The ethylene concentration was between 45 and 83 ppm on the first day and peaked five days later (15 November). The ethylene concentration was most pronounced at the lowest fruit/leaf ratio during the ripening period and was lowest in fruits from trees with a low fruit/leaf ratio during the last part of the ripening period. The greater availability of assimilates at a low fruit/leaf ratio favors accumulation of substrates for synthesis of aroma compounds, larger pools of sugars and organic acids, and greater levels of aroma compound precursors like acetyl CoA and fatty acids, which are transformed into the esters and alcohols found in the study. High fruit load strongly inhibits vegetative growth in young trees, reducing leaf growth to half and shoot growth to one-third to half.

15 3rd RM for L15-16 - Advances in Fruit Aroma Volatile Research

Fruit growth is an important yield determinant, with at least 90% of the fruit's dry matter originating from assimilates formed in leaves via photosynthesis. In pome and stone fruit, the sugar-alcohol sorbitol is mainly formed and serves as the transport agent, while sucrose is the transport agent in most other species, including fruit bushes and strawberries. Approximately 21% of the dry matter produced in leaves in autumn is stored as reserves (starch, sorbitol, and sugar), with a significant share (18%) stored in the roots. Storage material supplies 50-65% of the building material for flower buds and young shoot tips in the very early growth stages. The great import ability of fruits means they dominate competition for assimilates when they are out of the early growth stages. For example, a fruit in July-August consumes approx. 80% of the assimilates formed in the leaves on the same spur. High fruit load strongly inhibits vegetative growth, potentially reducing leaf growth to half and shoot growth to one-third to half in young apple trees. The effect is more powerful further from the fruit, coinciding temporally with the main growth period of organs like roots (September-October). This strong sink activity causes a greater photosynthetic intensity and more rapid transport out of leaves in fruiting plants.

Biological variation is substantial, and even within the same fruit cultivar, flavor and aroma can vary significantly. This variation affects uniformity and categorization strategies. Apples from three cultivars ('Elshof', 'Holsteiner Cox', and 'Ingrid Marie') were analyzed individually for sugar, acid, and aroma compounds, having been picked at maturity from four zones on the trees (top, bottom, east, west). Using Principal Component Analysis on the relative peak areas of 59 tentatively identified aroma compounds, the three cultivars were completely separated. For 'Ingrid Marie' apples, effects of position could not be elucidated due to inadequate juice yield.

A discriminant Partial Least Squares regression analysis revealed distinct differences based on position for 'Elshof' apples, separating apples from the top and bottom parts of the trees. The level of sugar (soluble solids) was highest in apples from the top (13.4% sugar) and lowest in apples from the bottom (12.3% sugar). This positional effect relates to differences in quality and ripening levels, likely associated with light exposure affecting photosynthetic activity. High soluble solids are generally associated with high production of esters. For 'Elshof', compounds high in the top included 2-Methylbutyl acetate (2400a) and 2-Phenylethyl acetate (2.7a), while Hexanal (1300a) and 2-Butanol (5.6a) were high in the bottom. For 'Holsteiner Cox', apples from the west side separated well, characterized by lower levels of specific compounds, including 4-Methyl-5-vinylthiazole (2.2b), Anethole (1.0b), and 3-Hexenal (16b). Differences between top and bottom were most evident in 'Elshof'.

The magnitude of variations in concentrations of characteristic odorants (e.g., 2-Methylbutyl acetate, pentyl acetate) is substantial, indicating that position on the tree is likely to affect sensory quality. It is concluded that when apples are sampled for categorization and characterization, it is extremely important to sample from all parts of the trees or clearly define the position from which the fruits are sampled. Data also indicates the importance of good light exposure on fruits to ensure high quality.

16 1st RM for L17+18 - Sustainable production systems for organic apple production

Organic apples often sold for fresh consumption must meet high quality requirements, obtaining the right size and being undamaged without important infections of pest and diseases. The organism most often causing damage

on the skin of apples is apple scab, which creates brown or black spots. This disease causes a big reduction in yield and quality in organic production. The yield reduction for growers not using copper for control was 86% compared to conventional production, depending on the cultivar and the year. Copper is an effective fungicide used in organic apple production in some European countries, but it has not been permitted in Denmark for the last 10 years, and the European Union wants to reject it from the list of permitted pesticides [13].

To improve quality and yield, research has been carried out on cultivars with low susceptibility, using combinations of fertilisation, rootstocks, and planting distance to prevent or reduce apple scab infections. Control has involved using the warning program RIMpro to predict severe ascospore discharge periods. The most important prevention method is to use genetic resistant apple cultivars or cultivars with less susceptibility. Small, open, rather slow-growing apple trees reduce infection possibilities. The resistance to apple scab was broken down in most cultivars, but 'Florina' was still fully resistant during the experiment, and 'Vanda', 'Retina', and 'Redfree' were less infected.

A high level of available nitrogen in the soil causes increasing growth and an extended growth season, which gives better infection possibilities for apple scab. Increased nitrogen also causes a decrease in the content of phenols in plant tissue. The effect of cover crops was investigated, comparing a permanent grass mixture (1), a clover grass mixture (2), and an annual cover crop (3). Fruits produced on trees managed with a permanent grass alleyway (1) had a lower nitrogen supply and obtained the best skin coloration. A lower nitrogen supply, especially during fruit development, resulted in more red fruits. Apple scab infections were more numerous on apples grown in the annual cover crop (3), which gave the largest nitrogen supply. Fruits from the grass alleyways (1) had the highest percentage of marketable fruits. Although the gross yield was bigger from trees grown in the annual cover crop (3), the marketable fruit crop was at the same level due to the higher percentage of disease infections.

Early end of vegetative growth and development of end buds in autumn are very important to reduce the possibility of late infection in autumn and thereby reduce the risk of spring infections from conidia wintering in woody parts. End bud development of 51 rootstocks showed that rootstocks with high winter hardiness also had the earliest end bud development. MM106, a rootstock with poor winter hardiness and the latest end bud development of the 51 rootstocks, should not be recommended for organic apple production. The weak rootstocks M9 and the Russian rootstock B9 are potentially suitable due to:

1. Early end bud development.
2. Weak growth (M9).
3. High winter hardiness (B9).

The fruit quality recorded consists of both outer quality (fruit size, fruit color, and damages caused by pest and diseases on the fruit skin) and inner quality (firmness and content of sugar and starch).

17 2nd RM for L17+18 - Fruit quality in organic and conventional farming: advantages and limitations

Fruit quality is essential for nutrition and human health, requiring urgent attention in current agricultural practices. Modern agriculture must address the challenges of sustainability, crop quality, and yield, corresponding to United Nations Sustainable Development Goals (SDGs). Conventional agriculture, which accounts for a huge part of global plant production, utilizes an average of 1.81 kg of pesticides and 28.48 kg of nitrogen per hectare. Fruit quality is a complex, multifactorial concept assessed by consumers based on organoleptic traits (appearance, aroma, flavor, and texture), maturity stage, nutritional value, and presence of harmful chemical compounds. Fruits

contain essential bioactive compounds, including vitamins (E, C, B group), essential mineral nutrients (K, Ca, Mg, P, S, Fe, Zn, Mn), flavonoids, isoprenoids, and anthocyanins. Immature fruit is characterized by excessive firmness, poor aroma, low sucrose, high starch, and green color [14].

Organic farming is a more sustainable production model than conventional agriculture. It can provide higher quality in some fruit crops due to the absence of synthetic fertilizers and pesticides, enhanced pollination, and reduced protection treatments, which boosts antioxidant compound production. A meta-analysis of 71 studies published over the past 5 years (2019-2023) showed that organic farming improved overall fruit quality in 38 (53.5%) cases, whereas only 6 (8.4%) showed negative effects. Effects appear largely species-specific, with grape berries, peppers, and apples benefiting most.

Organic farming inherently contributes to high-quality fruit through several properties: biologically active soil, management practices such as compost and green manure, enhanced pollination, low/moderate biotic stress, and the absence of harmful compounds derived from fertilizers and pesticides. Total pesticide content in organic farming soils is 70-90% lower than in conventional soils, and nitrate and nitrite residues are usually lower in organically grown crops. Long-term improved soil health is a key principle of organic farming. The low/moderate biotic stress inherent to organic systems, coupled with the absence of synthetic pesticides, leads to a controlled exposure to stress factors. This pathogen exposure can trigger the synthesis of plant defense metabolites, such as phenolics, terpenoids, and flavonoids. Salicylic acid and jasmonates are phytohormones involved in activating phenylpropanoid metabolism in response to biotic stress, enhancing compounds valuable for human nutrition. Furthermore, organic farms enhance pollination success.

However, organic farming is not without potential risks. The use of compost and organic fertilizers, such as manure, means a significant input of microorganisms, potentially increasing the occurrence of pathogenic bacteria like *Escherichia coli*, yeasts, molds, and *Bacillus cereus* in organically grown vegetables. Nonmicrobial contaminants, such as heavy metals (Cu and Zn), can accumulate from sources like municipal sewage sludge or animal manures. Copper-containing plant protection products used in organic farming can also contribute to heavy metal excess. Despite these risks, organic farms generally perform much better in terms of pesticide residues. Fruit quality is a multifactorial concept affected by factors such as stress intensity, pruning and crop load, genetic traits, and rootstocks in both conventional and organic models. The integration of all parameters is essential for achieving high quality and productivity in sustainable farming models.

18 RM for L19 - 9 Light Relations

Visible light in the 400-700 nm waveband is the driving factor of biomass production via photosynthesis. Although the production of dry matter in apple correlates to the amount of visible light intercepted by trees, dry-matter production does not automatically translate into increased yield of marketable fruit. Light plays a twofold role: influencing processes leading to the production of large quantities of high-quality fruit, and supplying energy stored in chemical form (carbohydrates). The quantity of light plants use is defined as Photosynthetic Photon Flux (PPF) or Photosynthetically Active Radiation (PAR). Overhead sunny conditions yield a PPF of around $2000 \mu\text{mol m}^{-2} \cdot \text{s}^{-1}$. Diffuse radiation can penetrate the canopy virtually in every direction, resulting in higher light levels in inner tree-canopy positions. Apple leaves absorb approximately 80% of incoming visible radiation [15].

Orchard management aims to maximize light interception, with efficient orchards intercepting 60-90% of available radiation, yielding high marketable production (up to $120\text{-}140 \text{ t ha}^{-1} \text{ year}^{-1}$). Orchard productivity is often higher when light interception is greater than 50% of available light. Total tree net CO_2 exchange (NCE) is found to be greater for fruiting trees than for non-fruiting trees when expressed per tree leaf area. Fruiting reduces light

interception due to lower leaf area, but stimulates the photosynthetic process, making the foliage more efficient. Yield ($\text{t} \cdot \text{ha}^{-1}$) is strongly correlated with whole-canopy Leaf Area Index (LAI) ($R^2=0.774$).

Apple leaves develop distinct morphological and physiological traits reflecting their light environment. 'Sun' leaves (exterior, well-illuminated) exhibit greater maximum photosynthetic capacity than 'shade' leaves (interior, low PPF). Shade leaves are typically denser, possessing more layers of cells, and higher nitrogen content. Shading spur leaves in the previous season was found to reduce the leaf photosynthetic rates.

The partitioning of carbon between different sinks (vegetative vs. reproductive) is complex. Primary spur leaves are vital early in the season, supplying carbohydrates for fruit set and cell division. They are the only carbon source until developing shoot leaves become net exporters (around 10-15 unfolded leaves). Early shading delays the transition of shoot carbohydrates to export. Heavy shading (70%) applied 5 weeks after full bloom (AFB) drastically reduced fruit growth and increased fruit drop. Low PPFs reduce flower-bud differentiation, fruit set, fruit size, soluble-solids concentration, and firmness.

Red fruit color formation (anthocyanin synthesis) is light-dependent, requiring light intensity and quality. This process is stimulated by UV-B radiation ($<320 \text{ nm}$) and enhanced synergistically by red light. Fruits grown under canopy shade show increased transpiration and shrivelling during storage.

Controlling light levels in the orchard improves quality. Pruning is used to maintain adequate light penetration. The optimum training system for maximizing light distribution is the slender spindle. Reflective materials (Reflectants), such as aluminum or tar-paper, increase light availability in the lower and inner parts of the canopy. Reflective ground cover applied 2-3 months before harvest improves skin color (via UV region effects), increases fruit size, and boosts soluble-solids concentration. Bagging fruits 4-6 weeks AFB, followed by bag removal, also enhances red color formation.

Chapter 3

Exam Questions and Answers

This chapter of the course notes compiles the exam questions for the course held in November 2025, along with their respective answers prepared by me. The purpose of this section is twofold: firstly, to provide a reflective exercise that consolidates understanding of the course material; and secondly, to document my comprehension of the course topics as assessed through the exam questions.

To ensure citation accuracy and academic transparency, NotebookLM has been employed as the primary generative AI platform. Its use has focused on verifying that all citations accurately reference the uploaded course materials and lecture slides provided by the professors. Beyond citation control, this section also represents an ongoing exploration of prompt engineering — refining interaction design to optimise AI output quality, precision, and academic reliability. Through this approach, the work aims to maintain a high academic standard while enhancing clarity, structure, and depth in written responses.

There are a total of 17 questions in the exam, each comprising between three and five sub-questions. The numbering of the sections in this chapter corresponds directly to the numbering of the exam questions, ensuring a clear and consistent structure throughout. Questions 1-9 address aspects related to crop physiology, while questions 10-17 focus on fruit quality, maturity, and usability. Each question is presented below, followed by its respective sub-questions and answers.

Questions within: Crop Physiology aspects

1 Yield and quality determinants and components

Shoot and bud development, growth and flower bud development

- 1.1 Characterise the development and importance of spurs and extension (long) shoots**
- 1.2 Describe differences in bud development and structure between stone and pome fruits**
- 1.3 Describe some important yield components in strawberry and in sour cherry**
- 1.4 Describe some conditions which may affect the development of flower buds negatively**

2 Yield and quality determinants and components

Flowers, pollination and fruit set (sterility and fertility)

- 2.1 Describe important factors determining fruit set?**
- 2.2 What is the importance of EPP?**
- 2.3 What are important quality parameters for pollen and flowers?**
- 2.4 Why and how do we use pollinators?**
- 2.5 Are insects (fx bees) needed in pollination of self-pollinating crops?**

3 Fruit development

Fruit development of small and large fruited species

- 3.1 Describe the general developmental phases in fruit development**
- 3.2 Which sugars and acids are important in fruit development and how do they develop during fruit development? Example of species differences.**
- 3.3 Which sugars are transported in the plant?**
- 3.4 What is the role of starch in the carbon balance of an apple tree and an apple fruit?**

4 Light use, vigor control and canopy management

Canopy management (pruning, growing systems, light use)

- 4.1 Why do we manipulate the canopy structure in most fruit crops?**
- 4.2 Describe the pruning response during the year. Why do we get differences in the growth response to pruning?**
- 4.3 How does pruning affect fruit development and quality? (direct and indirect)**
- 4.4 Characterise important factors (except from time in the year), which may influence the growth response to pruning?**

5 Crop load and canopy management

Carbon allocation (source-sink, fruit/leaf)

- 5.1 How does a high fruit load influence photosynthesis and transpiration?**
- 5.2 Explain the concept of source strength and sink strength**
- 5.3 How do source-sink relationships develop during the season in an apple tree?**
- 5.4 Why may some leaves be more important than others for fruit development?**
- 5.5 Why do premature fruit drop occur?**

6 Crop load management, fruit quality and vigor control

Thinning of fruits, how, why, when and effects

- 6.1 Give an example of a crop in which crop load has a strong impact on fruit development - and one where it does not.**
- 6.2 Characterize the effects of fruit thinning on growth and development**
- 6.3 When is it most optimal to perform fruit thinning? Why?**
- 6.4 Explain why the optimal thinning strategy may depend on the end use of the fruits.**
- 6.5 Why do we not want fruits on a young tree the first year(s) after planting?**

7 Preharvest factor management and quality

Use and management of nutrients

- 7.1 Characterise the differences in nutrient requirements of a vegetative growing and a fruiting plant?**
- 7.2 Calcium is important for fruit quality. Why? - And why is the level of calcium low in many fruits, especially big fruits?**
- 7.3 When and why are fertilizers often sprayed on the leaves and fruits in the production of apples?**
- 7.4 Characterize the importance of potassium for fruit development**

8 Preharvest factor management and quality

Effects of nutrients on yield and quality

- 8.1 Describe the effects of nitrogen status on plant development**
- 8.2 In which ways do nitrogen levels influence the yield components?**
- 8.3 Impacts of nitrogen levels on fruit quality?**

9 Preharvest factor management and quality

Effects of stresses on yield and quality

- 9.1 Describe the effects of stresses of nutrients and water on fruit development and quality.**
- 9.2 Why are deficiency symptoms by some nutrients seen in the young leaves and by others in the old?**
- 9.3 Describe how water stress can be used as a tool for growth control.**

Questions within: Fruit quality, maturity and usability aspects

10 Fruit development

Influencing factors

- 10.1 Describe some important factors for optimal fruit development in small and large fruited species. Are there differences?**
- 10.2 What would you do to optimize fruit development and fruit quality in an apple crop?**
- 10.3 What is important for fruit development and quality in raspberry and strawberry?**

11 Fruit maturity, harvest and quality assessment

Maturity measures, Harvest time and methods

- 11.1 How would you determine the optimal harvest time in apple?**
- 11.2 Describe the problems and quality effects you might get, if you harvest either too early or too late.**
- 11.3 Hand picking vs mechanical harvest - problems and benefits?**
- 11.4 What are the main reasons for post harvest losses and what may be done to minimize it?**

12 Fruit maturity, cultivar variations and important quality parameters

Aromas in fruits and effects on aroma development

12.1 When do aromas develop in fruits?

12.2 Characterise some important aroma substances and changes in aroma with maturity

12.3 Characterize the importance of harvest time on aroma development

12.4 What might affect aroma development pre and post harvest?

13 Fruit maturity, cultivar variations and important quality parameters

Colors in fruit and berries and effects on colour development

13.1 Characterise some important colour substances in fruits and berries

13.2 How does colour change with maturity?

13.3 What might affect colour development pre and post harvest?

13.4 What is the mechanism behind the occurrence of red clones in fruit cultivars (fx apples, pears and grapes)?

14 Cultivar variations and important quality parameters (fresh use and juice)

Cultivar characterization and uses. Fruit composition and human health

- 14.1** Characterise some of the most important (internal and external) quality characters, which may vary among cultivars in a fruit crop. (Ex strawberries or apple)
- 14.2** Which compounds are considered especially important in fruit and berries for human health and where are they located?
- 14.3** Which species are believed to be especially healthy to eat? Comment on the consumption of raw or processed fruits and berries.

15 Cultivar variations and important quality parameters (fresh use and juice)

Juice processing and juice quality

- 15.1** How does the level of fruit ripening impact on juice processing and juice quality?
- 15.2** Which enzymes may be used in juice processing and why?
- 15.3** Comment on the effects of different juice processing steps on juice quality.
- 15.4** Why are juices pasteurised, and what are important factors for a successful pasteurisation?

16 Potentials for producing fruit and berry wines

Challenges and opportunities

- 16.1** Comment on the challenges and potentials in making fruit wine from different fruit and berries
- 16.2** High levels of acidity may be a problem. How may it be handled?
- 16.3** Characteristics of so called ‘cider apple cultivars’
- 16.4** Comment on the importance of ripening levels of fruit and berries for wine making
- 16.5** Characterize the process of cryo-concentration and the impacts on the juice quality and the potential for wine style development

17 Domestication of wild berries

Challenges and opportunities

- 17.1** Why may wild berries be attractive to domesticate?
- 17.2** Comment on some major challenges/barriers.
- 17.3** Describe important yield and quality components in wild/European blueberries.
- 17.4** Blueberries are one of few fruiting plants adapted to low pH soils. Comment on the challenges it causes in growing the plants.
- 17.5** Comment on the importance/impacts of propagation method in European blueberries. European blueberries.

Chapter 4

Abbreviations and Explanations

Topic	Abb.	Description
Leaching	n.a.	<i>leaching refers to the process by which substances, such as ions, minerals, or nutrients, are removed or lost from the soil. This often occurs due to water penetrating the soil and displacing these substances</i>

Bibliography

- [1] Torben Toldam-Andersen. “FRUIT YIELD AND QUALITY. COMPONENTS AND DETERMINANTS”. Based on Hansen 1995. 2010.
- [2] Torben Toldam-Andersen. “BUDS AND BUD DEVELOPMENT.” Based on Hansen 1995. 2010.
- [3] Torben Toldam-Andersen. “CHAPTER 2 Flower bud formation, flower development and flowering”. Based on Hansen 1995. 2011.
- [4] Torben B. Andersen. “A model to predict the beginning of the pollen season”. In: *Grana* 30 (1 1991), pp. 269–275. DOI: [10.1080/00173139109427810](https://doi.org/10.1080/00173139109427810).
- [5] Miklos Faust. *Physiology of temperate zone fruit trees*. 1989.
- [6] Torben Bo Toldam-Andersen. “Green harvest/fruit thinning: How, when, how much and why? – Or how we best utilize the reactions of the plant to changes in the fruit/leaf ratio”. In: *Vinpressen* 4 (2017). Associate Professor University of Copenhagen.
- [7] SJ Wertheim and AD Webster. “Manipulation of growth and development by plant bioregulators”. In: *Fundamentals of temperate zone tree fruit production* (2005), pp. 267–294.
- [8] Torben Toldam-Andersen. “CHAPTER 5 Growth, Growth correlations and Assimilate turnover”. Based on Hansen 1995. 2011.
- [9] Torben Bo Toldam-Andersen. *Table Grapes in Cold Houses*. Compendium note for the course: Fruit and Berry Crop Physiology and Quality. Copenhagen, Denmark, 2016.
- [10] Torben Toldam-Andersen. “CHAPTER 6 Fruit growth and fruit quality”. Based on Hansen 1995. 2011.
- [11] Muna Ahmed Mohamed El Hadi et al. “Advances in fruit aroma volatile research”. In: *Molecules* 18.7 (2013), pp. 8200–8229.
- [12] Leif Poll et al. “Availability of assimilates and formation of aroma compounds in apples as affected by the fruit/leaf ratio”. In: *Physiologia plantarum* 97.2 (1996), pp. 223–227.
- [13] Hanne Lindhard Pedersen. “Sustainable production systems for organic apple production”. In: *DARCOF II-Danish Research in Organic food and Farming systems 2000-2005* (2006), pp. 12–15.
- [14] Sergi Munné-Bosch and Núria F Bermejo. “Fruit quality in organic and conventional farming: advantages and limitations”. In: *Trends in Plant Science* 29.8 (2024), pp. 878–894.
- [15] Luca Corelli Grappadelli. “Light relations.” In: *Apples: botany, production and uses*. CABI Publishing Wallingford UK, 2003, pp. 195–216.

Appendices

1 Appendix 1 - Practical Exercise 01

PE1: Tropical Crop products

Group n. 01

Group members:

- Lucas Daniel Paz Zuleta, TZS159

Photo of your culinary preparation



List the tropical products used:

White rice, Pequi, Okra, Black beans, lentils, Cassava, salad (mix; rocula, spinach), Assorted Cherry tomatoes, Pineapple, Lentils, Olive oil, Palm hearts, and Jílo.

Discuss the potential macro nutrients composition of your dish (Use chatgpt):

Potential Macronutrient Composition of the Dish

- **Carbohydrates:**

White rice, cassava, lentils, and black beans are major carbohydrate sources, providing both starch and dietary fibre. Pequi and pineapple add natural sugars. Okra, cherry tomatoes, salad greens, palm hearts, and jiló contribute smaller amounts of carbohydrates, mainly fibre.

- **Proteins:**

Black beans and lentils are the primary plant-based protein sources. Spinach, arugula, and other salad vegetables contribute minor amounts of protein.

- **Fats:**

Olive oil and pequi are the main fat sources. Pequi contains monounsaturated fats, while olive oil contributes healthy unsaturated fats. Small contributions may also come from palm hearts.

- **Fibre:**

High levels of dietary fibre come from legumes (black beans, lentils), okra, cassava, salad greens, cherry tomatoes, jiló, and pineapple. Okra in particular also adds soluble fibre (mucilage).

This dish is **balanced**:

- **Carbohydrates** from rice, cassava, and legumes.
- **Proteins** mainly from legumes.
- **Fats** from olive oil and pequi.
- **Fibre and micronutrients** from vegetables, fruits, and jiló

[H]