



Aroma Development and Volatile Composition in Mango

A Theoretical Report on the Biochemical Pathways, Ripening Physiology, and Quality Determinants in mango.

Lucas Daniel Paz Zuleta, TZS159

MSc students at the University of Copenhagen

Last compiled: 28-10-2025

[Link to Git repository](#)



Preface

This written report has been prepared as part of the course NPLK14014U - Fruit and Berry Crop Physiology and Quality at the University of Copenhagen. The course explores the physiological and biochemical factors determining fruit development, yield, and quality in relation to fresh consumption and processing. Emphasis is placed on how genetic, environmental, and management factors influence internal and external fruit quality, including the formation of secondary metabolites and sensory attributes such as aroma.

The present project is a theoretical study on the development and composition of aroma in mango, *Mangifera indica* L., The focus is on the biochemical formation of volatile organic compounds (VOCs) and their relationship to fruit ripening and sensory quality. The report integrates physiological and biochemical knowledge to describe how preharvest conditions, cultivar differences, and post-harvest handling affect aroma expression and the perception of fruit quality.

The assignment was carried out by Lucas Daniel Paz Zuleta (TZS159), MSc student at the University of Copenhagen, as an individual written report fulfilling the course requirements for NPLK14014U - Fruit and Berry Crop Physiology and Quality.

Summary

Signatures

Copenhagen, 28-10-2025

A handwritten signature in black ink, appearing to read 'L. D. Paz Zuleta', is written over a horizontal line.

Lucas Daniel Paz Zuleta (TZS159)
28-10-2025

Contents

1	Introduction	1
1.1	Background	1
1.2	Focus on Mango	1
1.3	Aim and Scope	1
2	Aroma Composition in Mango	1
2.1	Chemical Classes of Mango Aroma Compounds	1
2.2	Key Aroma Compounds in Mango	2
2.3	Variation in Aroma Profiles Among Mango Varieties	4
3	Biochemical Pathways of Aroma Compound Formation	4
3.1	Overview of Biosynthetic Pathways	4
3.2	Enzymes Involved in Aroma Biosynthesis	4
3.3	Terpenoid Pathway (MEP and MVA Pathways)	4
3.4	Fatty Acid Derivative Pathway	4
3.5	Amino Acid Derivative Pathway	4
4	Environmental and Genetic Factors Influencing Aroma Production	4
4.1	Pre-harvest Factors	4
4.2	Post-harvest Factors	4
4.3	Processing Implications on Aroma Retention	4
5	Analytical Techniques for Aroma Compound Identification	4
5.1	Extraction and Analysis Methods	4
5.2	Quantification and Sensory Evaluation	4
6	Comparative Perspective	4
7	Conclusion	4

1 Introduction

Fruits are complex and heterogeneous structures that comprise a wide variety of metabolites that serve as precursors to volatile organic compounds (VOCs), such as carbohydrates, fatty acids, and pigments [1]. The mango fruit is no exception of this complexity and have in recent years been the focus of many studies, studying the activity of volatiles in the fruit [2].

1.1 Background

Mango is a tropical fruit which belongs to the Anacardiaceae family and is scientifically known as *Mangifera* [2]. It is popularly characterised as a sweet, juicy, aromatic fruit with a low fibre flesh [3]. It is primarily cultivated in tropical and subtropical regions, where it is of significant economic importance [3]. The annual tropical production of mango is over 46 million tons and is thereby the most produced tropical fruit after banana [4]. The perishable nature and susceptibility to post-harvest losses and diseases pose challenges for the mango industry, restricting the production and potential [3].

1.2 Focus on Mango

The mango is an important fruit crop worldwide, valued for its high nutritional content, with significant levels of fibre, vitamin C and β -carotene [5, 4]. However, beyond its nutritional benefits, the mango is particularly renowned for its distinctive aroma, which plays a crucial role in consumer preference and marketability [6]. s of quality and freshness [3]. A study by Badar et al. (2016) found that aroma is one of the most important quality attributes that influence consumer preference and acceptance of mango fruit [6]. The main aroma contributors in mango are the VOCs; aldehydes, alcohols, esters and ketones [7]. Among these, 3-carene, limonene, β -pinene, acetaldehyde, ethanol and hexanal [7]. Understanding the origin and behaviour of these volatiles is therefore essential for improving fruit quality, post-harvest handling, and processing applications.

1.3 Aim and Scope

The aim of this report is to explore the formation, composition, and significance of VOCs in mango fruits. The main focus is on how these compounds contribute to the fruit's aroma and overall quality, both in terms of maintaining freshness and enhancing consumer appeal. The scope includes an overview of the chemical classes and key VOCs identified in mango, the metabolic- and enzymatic pathways involved in their biosynthesis, and the influence of pre- and post-harvest conditions on their abundance. Finally, the report outlines the methods commonly applied for the extraction and identification of VOCs and their relevance for assessing fruit quality and consumer perception.

2 Aroma Composition in Mango

The complexity of aromas in mango is due to the variety of VOCs that are present in the fruit's matrix. These compounds arise from different biochemical pathways and contribute to the overall sensory experience of the fruit [3]. These pathways are most prominently manifested during the ripening processes of the fruit. Also visual and textural changes occur during ripening, which further influence the perception of aroma [5, 3].

2.1 Chemical Classes of Mango Aroma Compounds

The volatile composition in fresh mango fruits has been extensively studied, revealing a diverse array with several hundred identified volatile compounds, occurring in free form in the fruit [4]. By calculating the flavour dilution factor (FD) of the volatile compounds using gas chromatography-olfactometry (GC-O) analysis, it has been possible to identify the key aroma compounds for the odour-active fraction [4].

Mango aroma is primarily composed of several chemical classes. The most dominant classes is monoterpene hydrocarbons, which account for 90.2% of the total volatile compounds in fresh mango, as illustrated in Figure 1.1. Other significant classes include lactones (4.1%) and sesquiterpene hydrocarbons (3.0%) [4].




Figure 1.1: Distribution [%] of chemical classes of volatile compounds in fresh mango. Adapted from Bonneau et al. (2016) [4].

2.2 Key Aroma Compounds in Mango

In this subsection, the key aroma compounds identified from fresh mango fruits in the study by Bonneau et al. (2016) are discussed [4].

Terpenes and Terpenoids

Terpenes represent the largest class of mango aroma compounds, derived from isoprene units via the terpenoid biosynthetic pathway [8]. They include both monoterpenes and sesquiterpenes, which together account for the majority of mango volatiles [4]. Terpenes and terpenoids are found in many different natural sources, including fruits, plants, animals, microbes, and fungi. The terpenes belong to the largest class of secondary metabolites in nature and consist of five connected carbon atoms, known as isoprene units. These carbon units can be assembled in thousands of ways [9]. The terpenoids are further subcategorized into five sub-groups based on the number of isoprene units they contain: monoterpenes (C₁₀), sesquiterpenes (C₁₅), diterpenes (C₂₀), sesterterpenes (C₂₅), and triterpenes (C₃₀) [9].

Monoterpene hydrocarbons A total of 11 monoterpene hydrocarbons were identified as key aroma compounds in fresh mango [4]. The most significant ones include: α -phellandrene, γ -terpinene, δ -3-carene, β -myrcene, α -terpinene, limonene, β -phellandrene, and α -terpineol.

Monoterpenes are the smallest molecules in the isoprenoid family with conserved hydrocarbons [8]. They share the formula C₁₀H₁₆ and over 400 different chemical structures have been classified as such [8]. The key monoterpene hydrocarbons identified in mangos are reported to have significant impact on the overall odorants [4].

Sesquiterpene hydrocarbons Compared to monoterpenes, sesquiterpenes are larger molecules with the formula C₁₅H₂₄. In mango fruit, the study by Bonneau et al. (2016) identified four key sesquiterpene hydrocarbons, including: α -gurjunene, α -copaene, β -caryophyllene, and α -caryophyllene [4].

Alcohols and Aromatic Alcohols

Alcohols and aromatic alcohols are, like terpenes and terpenoids, major contributors to the key aroma of mango, though they differ in chemical structure and biosynthetic origin. A study by Singh et al. (2010) highlighted the role of alcohol dehydrogenase (ADH) in the enzymatic reduction of aldehydes leading to the formation of these compounds [10].

A total of nine alcohols were identified as key aroma compounds in fresh mango [4]. The most significant ones include: 2-methyl-1-propanol, 1-pentanol, (*E*)-2-penten-1-ol, (*Z*)-2-penten-1-ol, 1-octanol, 1-butanol, 3-methyl-1-butanol, 2-decanol, 1-hexanol, and (*Z*)-3-hexen-1-ol [4].

Aldehydes and Aromatic Aldehydes

Aldehydes are volatile compounds primarily formed through the oxidation of unsaturated fatty acids, such as α -linolenic acid, via the lipoxygenase (LOX)–hydroperoxide lyase (HPL) pathway. These reactions generate C₆ and C₉ aldehydes, often referred to as green leaf volatiles, which are associated with fresh and grassy notes in the mango fruit aroma [11]. In mango, this pathway becomes particularly active under chilling stress, leading to increased levels of compounds such as *1-hexanal*, *(E)-2-hexenal*, and *(Z)-3-hexenal* before visible quality loss occurs.

In fresh mango, a total of six aldehydes were identified as key aroma compounds [4]. The most significant ones include hexanal, *(E,E)*-2,4-heptadienal, nonanal, and *(E,Z)*-2,4-heptadienal [4].

Ketones

handling

Esters

handling

Furans

handling

Lactones

handling

2.3 Variation in Aroma Profiles Among Mango Varieties

3 Biochemical Pathways of Aroma Compound Formation

3.1 Overview of Biosynthetic Pathways

3.2 Enzymes Involved in Aroma Biosynthesis

3.3 Terpenoid Pathway (MEP and MVA Pathways)

3.4 Fatty Acid Derivative Pathway

3.5 Amino Acid Derivative Pathway

4 Environmental and Genetic Factors Influencing Aroma Production

4.1 Pre-harvest Factors

4.2 Post-harvest Factors

4.3 Processing Implications on Aroma Retention

5 Analytical Techniques for Aroma Compound Identification

5.1 Extraction and Analysis Methods

5.2 Quantification and Sensory Evaluation

6 Comparative Perspective

7 Conclusion

Bibliography

- [1] Norma S. Pano-Farías et al. “Direct immersion single drop micro-extraction method for multi-class pesticides analysis in mango using GC–MS”. In: *Food Chemistry* 237 (2017), pp. 30–38. ISSN: 0308-8146. DOI: [10.1016/j.foodchem.2017.05.030](https://doi.org/10.1016/j.foodchem.2017.05.030). URL: <https://doi.org/10.1016/j.foodchem.2017.05.030>.
- [2] Ming Guo et al. “Nutritive value of active volatile components of Anacardiaceae mango and their effects on carrier proteins function”. In: *Food Research International* 168 (2023), p. 112779. ISSN: 0963-9969. DOI: <https://doi.org/10.1016/j.foodres.2023.112779>. URL: <https://www.sciencedirect.com/science/article/pii/S0963996923003241>.
- [3] C. F. Chin, E. Y. Teoh, M. J. Y. Chee, et al. “Comparative Proteomic Analysis on Fruit Ripening Processes in Two Varieties of Tropical Mango (*Mangifera indica*)”. In: *Protein Journal* 38 (2019), pp. 704–715. ISSN: 1573-4943. DOI: [10.1007/s10930-019-09868-x](https://doi.org/10.1007/s10930-019-09868-x). URL: <https://doi.org/10.1007/s10930-019-09868-x>.
- [4] Adeline Bonneau et al. “Aroma compounds in fresh and dried mango fruit (*Mangifera indica* L. cv. Kent): impact of drying on volatile composition”. In: *International Journal of Food Science and Technology* 51.3 (Jan. 2016), pp. 789–800. ISSN: 0950-5423. DOI: [10.1111/ijfs.13038](https://doi.org/10.1111/ijfs.13038). eprint: <https://academic.oup.com/ijfst/article-pdf/51/3/789/59497344/ijfs13038.pdf>. URL: <https://doi.org/10.1111/ijfs.13038>.
- [5] D. A. Aguirre-López et al. “Compuestos orgánicos volátiles presentes en el aroma de 17 frutas exóticas en Colombia: revisión”. In: *Revista Colombiana de Investigaciones Agroindustriales* 10.2 (2023), pp. 117–142. DOI: [10.23850/24220582.5208](https://doi.org/10.23850/24220582.5208).
- [6] Hammad Badar, Anoma Ariyawardana, and R. Collins. “Mango value preferences of consumers in Pakistan”. In: *Acta Horticulturae* (July 2016), pp. 439–446. DOI: [10.17660/ActaHortic.2016.1120.68](https://doi.org/10.17660/ActaHortic.2016.1120.68).
- [7] Angélica Moreno et al. “Análisis del perfil de compuestos volátiles del mango (*Mangifera indica* L. var. Tommy Atkins) tratado por métodos combinados”. In: *Revista Colombiana de Química* 39.1 (2010), pp. 61–72. ISSN: 0120-2804. URL: <http://www.redalyc.org/articulo.oa?id=309026683005>.
- [8] Benedict J. Barras, Taotao Ling, and Fatima Rivas. “Recent Advances in Chemistry and Antioxidant/Anticancer Biology of Monoterpene and Meroterpenoid Natural Product”. In: *Molecules* 29.1 (2024). ISSN: 1420-3049. URL: <https://www.mdpi.com/1420-3049/29/1/279>.
- [9] Shagufta Perveen and Areej Al-Taweel, eds. *Terpenes and Terpenoids*. Open access publication. DOI link: <https://doi.org/10.5772/intechopen.71175>. London, United Kingdom: IntechOpen, 2018. ISBN: 9781789847772, 9781789847765, 9781838815295. DOI: [10.5772/intechopen.71175](https://doi.org/10.5772/intechopen.71175). URL: https://mts.intechopen.com/storage/books/6530/authors_book/authors_book.pdf.
- [10] Rajesh K. Singh et al. “Differential expression of the mango alcohol dehydrogenase gene family during ripening”. In: *Phytochemistry* 71.13 (2010), pp. 1485–1494. ISSN: 0031-9422. DOI: <https://doi.org/10.1016/j.phytochem.2010.05.024>. URL: <https://www.sciencedirect.com/science/article/pii/S0031942210002189>.
- [11] Velu Sivankalyani et al. “Chilling Stress Upregulates α -Linolenic Acid–Oxidation Pathway and Induces Volatiles of C6 and C9 Aldehydes in Mango Fruit”. In: *Journal of Agricultural and Food Chemistry* 65.3 (2017), pp. 632–638. ISSN: 0021-8561. DOI: [10.1021/acs.jafc.6b04355](https://doi.org/10.1021/acs.jafc.6b04355). URL: <https://doi.org/10.1021/acs.jafc.6b04355>.