

Local variability of vegetation structure increases forest resilience to wildfire

Michael J Koontz^{1,2}, Malcolm P North^{2,3}, Stephen E Fick^{4,5}, Chhaya M Werner⁶, Andrew M Latimer²

¹Graduate Group in Ecology, University of California, Davis; Davis, CA

²Department of Plant Sciences, University of California, Davis; Davis, CA

³Pacific Southwest Research Station, U.S.D.A. Forest Service; Mammoth Lakes, CA

⁴U.S. Geological Survey, Southwest Biological Science Center ⁵Department of Ecology and Evolutionary Biology, University of Colorado, Boulder; Boulder, CO ⁶Center for Population Biology, University of California, Davis; Davis, CA

Abstract. The long-term persistence of forest ecosystems hinges on their resilience to changing conditions. Measurements of resilience are critical for these valuable ecosystems, but are challenging to capture at relevant scales given a forest's temporal longevity and vast spatial extent. Wildfire plays a key role in structuring the vegetation of many forests, and vegetation structure can feedback to influence wildfire behavior. Local variability of vegetation structure can enable a forest stand to withstand wildfire disturbance and retain its essential identity and function— a hallmark of a resilient system. We investigate the system-wide generality of this phenomenon over the entire spatial extent of the yellow pine/mixed-conifer forest system in the Sierra Nevada mountain range of California. We develop an approach to measure wildfire severity and local vegetation structure using Landsat satellite data and texture analysis across a 33-year time series of all known wildfires covering greater than 4 hectares in this region. We find that, on a system-wide scale, greater variability in local forest structure reduces the probability of a high severity wildfire. We find the most support for this relationship at the smallest spatial extent of vegetation structure tested, indicating this phenomenon manifests at a local scale. Variable forest structure thus makes yellow pine/mixed-conifer forest in the Sierra Nevada more resistant to inevitable wildfire disturbance on average, and may increase the probability of long-term forest persistence. Efforts to maintain or increase vegetation structure variability in forests, such as allowing fires to burn under some conditions, should be continued.

Introduction

Biological systems comprising heterogeneous elements can retain their fundamental properties in the face of regular disturbance. This ability of a heterogeneous system to absorb disturbances, reorganize, and to persist within a domain of stability with respect to its identity, structure, function, and feedbacks is termed resilience (Holling 1973; Gunderson 2000; Folke *et al.* 2004; Walker *et al.* 2004). Resilience has been demonstrated in complex biological systems characterized by a variety of different types of “heterogeneity” including genetic diversity (Reusch *et al.* 2005; Agashe 2009; Baskett *et al.* 2009), species diversity (Tilman 1994; Chesson 2000; Cadotte *et al.* 2013), functional diversity (Gazol & Camarero 2016), topoclimatic complexity (Ackerly *et al.* 2010, @Lenoir2013), and temporal environmental variation (Questad & Foster 2008). An emerging paradigm in forest ecology is that spatial heterogeneity in the structure of vegetation on the landscape

can confer resilience to disturbances such as wildfire, drought, and insect outbreaks (Stephens *et al.* 2008; North *et al.* 2009; Virah-Sawmy *et al.* 2009). In California, increasing temperature coupled with increasing drought frequency exacerbate water stress on trees during “hotter droughts” (Park Williams *et al.* 2012; Millar & Stephenson 2015). Further, a century of fire suppression policy has led to drastic densification and homogenization of forest structure in the Sierra Nevada (North *et al.* 2015). Wildfire regimes have changed in these forests such that fires are bigger and burn more at higher severity (Miller & Thode 2007; Cansler & McKenzie 2014; Harvey *et al.* 2016). Changes in wildfire disturbance regimes are particularly suited to catalyze catastrophic shifts in ecosystems because of their feedback with spatial forest heterogeneity at multiple scales. Thus, western North American forests are experiencing novel, “unhealthy” conditions (*sensu* Raffa *et al.* (2009)) that are liable to upset the feedbacks between forest structure and pattern-forming ecological disturbances that historically stabilized the system and made it resilient (Raffa *et al.* 2008; Millar & Stephenson 2015). Forests are of high management priority (Hansen *et al.* 2013; Crowther *et al.* 2015; Millar & Stephenson 2015; Trumbore *et al.* 2015), thus it is critical to understand the mechanisms underlying the effect of spatial heterogeneity in forest structure on forest resilience.

Resilience of forest ecosystems is fundamentally challenging to quantify because forests comprise long-lived species, span large geographic extents, and are affected by disturbances at a very broad range of spatial scales. The ease or difficulty with which a disturbance changes a system’s state is resistance, and it is a key component of resilience (Walker *et al.* 2004). In yellow pine/mixed-conifer forests of California’s Sierra Nevada mountain range, wildfire disturbance alters the system state at landscape scales with the deaths of overstory trees. Thus, a more resistant forest system should generally experience lower tree mortality when a fire inevitably occurs.

Wildfire severity describes the effect of a wildfire on an ecosystem— often the amount of vegetation mortality (Sugihara & Barbour 2006). Wildfire severity can be measured by comparing pre- and post-fire satellite imagery for a specific area, but this usually requires considerable manual effort for image collation and processing, followed by calibration with field data (Miller & Thode 2007; Miller *et al.* 2009; De Santis *et al.* 2010; Cansler & McKenzie 2012; Veraverbeke & Hook 2013; Parks *et al.* 2014; Prichard & Kennedy 2014; Edwards *et al.* 2018; Fernández-García *et al.* 2018). Efforts to measure severity across broad spatial extents, such as the Monitoring Trends in Burn Severity project (Eidenshink *et al.* 2007), are unsuitably subjective for rigorous scientific analysis though they serve their intended management purpose admirably (Kolden *et al.* 2015). Automated efforts to remotely assess wildfire have arisen, but they tend to focus on more aggregate measures of wildfire such as whether an area burned or the probability that it burned rather than the severity of the burn (Bastarrika *et al.* (2011); Goodwin & Collett (2014); Boschetti *et*

al. (2015); Hawbaker *et al.* (2017) but see Reilly *et al.* (2017) and (???). Here, we present a method to automate the measurement of wildfire severity using minimal user inputs: a geometry of interest (a wildfire perimeter or a field plot location) and an alarm date (the date the fire began). This information is readily available in many fire-prone areas (such as California, via the Fire and Resource Assessment Program; http://frap.fire.ca.gov/projects/fire_data/fire_perimeters_index) or could be derived using existing products (such as the Landsat Burned Area Essential Climate Variable product described in Hawbaker *et al.* (2017)). Further, the flexibility of this approach facilitates collaborative calibration with field-collected wildfire severity data.

Vegetation characteristics such as canopy density (Rouse *et al.* 1973; Young *et al.* 2017), moisture content (Asner *et al.* 2015), insect attack (Näsi *et al.* 2015), and even functional diversity (Asner *et al.* 2017) can be measured using remotely-sensed imagery. Texture analysis of imagery can quantify ecologically relevant environmental heterogeneity across broad spatial scales (Wood *et al.* 2012). Texture analysis was originally developed for image classification and computer vision, and it characterizes each pixel in an image by a summary of its neighboring pixels (Haralick *et al.* 1973; Connors *et al.* 1984). Ecologists have successfully used texture measurements to augment predictions of species richness (Huang *et al.* (2014); Stein *et al.* (2014); Tuanmu & Jetz (2015) but see Culbert *et al.* (2012)).

Creation and maintenance of spatial heterogeneity

Forest structure is defined by the size and distribution of trees on the landscape. Differences in tree crown heights characterize vertical structure, while differences in the rooting locations of trees characterizes horizontal structure (North *et al.* 2009). Competition for light, water, and other resources can yield aggregations of trees within favorable microsites, as well as areas containing trees that are more widely spaced trees where resources are more limiting (Clyatt *et al.* 2016). Demographic processes of dispersal, recruitment, and mortality affect forest structure by adding or subtracting whole trees. Reciprocally, forest structure can also influence these pattern-forming processes; for example, vegetation overstory density alters microclimate and changes understory tree demographic rates (Larson & Churchill 2012; De Frenne *et al.* 2013; Ford *et al.* 2013). The stabilizing effects of these reciprocal processes in forests are hallmarks of a resilient system (Folke *et al.* 2004). In the Sierra Nevada range of California, one of the strongest feedbacks between forest structure and pattern-generating ecological process is wildfire, which affects hundreds of thousands to millions of hectares of forested area per year in the Sierra Nevada (Larson & Churchill 2012; Park Williams *et al.* 2012; Millar & Stephenson 2015).

Wildfire interacts dynamically with the forest structure (Westerling *et al.* 2006; Larson & Churchill 2012; Park Williams *et al.* 2012). Wildfire can affect future forest structure by changing demographic rates of individual trees (e.g. increasing growth or germination via increasing light or nitrogen availability), but its most lasting impact to forest structure is in the pattern of killed trees left in its wake (Larson & Churchill 2012). Wildfire behavior is inherently complex and is influenced by local weather, topography, and heterogeneous fuel conditions created by departures from the average fire return interval at any particular place (Sugihara & Barbour 2006; Collins & Stephens 2010). For instance, high tree density and presence of “ladder fuels” in the understory increase the probability of crown fire that kills a high proportion of trees (Stephens *et al.* 2008). A heterogeneous forest can largely avoid overstory tree mortality because a reduced amount of accumulated ladder fuel decreases its ability to get into the crown (where mortality is more likely to result), because wide spacing between tree clumps interrupts high severity fire spread across the landscape, and because small tree clumps with fewer trees don’t facilitate self-propagating fire behavior (Graham *et al.* 2004; Scholl & Taylor 2010). In forests with relatively intact fire regimes and heterogeneous stand conditions such as in the Jeffrey pine/mixed-conifer forests of the Sierra San Pedro Mártir in Baja, California, there tends to be reduced vegetation mortality after wildfires compared to fire-suppressed forests (Stephens *et al.* 2008). Thus, forests with heterogeneous structure are predicted to persist due to their resistance to inevitable wildfire disturbance (Graham *et al.* 2004; Moritz *et al.* 2005; Stephens *et al.* 2008). However, it is unclear whether this is true at broad spatial extents, nor is it resolved at what scale variability in forest structure is meaningful for resilience (Kotliar & Wiens 1990).

We use Landsat satellite data and a new image processing approach to calculate wildfire severity for all Sierra Nevada wildfires since 1984 that burned in yellow pine/mixed-conifer forest and covered more than 4 hectares. We then pair this comprehensive, system-wide severity data with image texture analysis to ask: does spatial variability in forest structure increase the resilience of California yellow pine/mixed-conifer forests by reducing the severity of wildfires? Further, we ask whether the influence of structural heterogeneity on fire severity depends on topographic, fire weather, or other fuel conditions.

Methods

This work occurred in two phases. First, we developed a new approach to calculating wildfire severity across broad spatial and temporal scales and calibrated our measurements to those from the field. We applied this approach to all known fire perimeters greater than 4 hectares in the Sierra Nevada region between 1984 and 2017 as defined by the Fire and Resource Assessment Program (FRAP, <http://frap.fire.ca.gov/>

projects/fire_data/fire_perimeters_index), which is the most comprehensive digital record of fire occurrence in California. Second, we used texture analysis of remotely-sensed imagery bounded by the perimeters in the FRAP database to develop a measure of vegetation heterogeneity and modeled how that heterogeneity affected wildfire severity, accounting for other key drivers of wildfire behavior.

Study area

Our study assesses the effect of vegetation structure on wildfire severity in the Sierra Nevada mountain range of California in yellow pine/mixed-conifer forests between 1984 and 2016 (Fig. 1). Forests in our study area are dominated by a mixture of conifer species including ponderosa pine (*Pinus ponderosa*), sugar pine (*Pinus lambertiana*), incense-cedar (*Calocedrus decurrens*), Douglas-fir (*Pseudotsuga menziesii*), white fir (*Abies concolor*), and red fir (*Abies magnifica*) (Stephens & Collins 2004; Collins *et al.* 2015). Tree density in the early 20th century was relatively low, with about 25-79 trees/ha and about 8-30 m²/ha of live basal area (Collins *et al.* 2015, @Safford2017). Since this time, canopy cover has increased by 25-49%, overall tree density has increased by >75%, and white fir (*Abies concolor*) makes up a greater percentage of basal area compared to forests in the early 20th century (Stephens *et al.* 2015). The change in tree density is concurrent with a shift in size distribution: modern mixed-conifer forests have 2.5 times as many trees between 30.4 and 61.0cm diameter at breast height (dbh) per hectare (103.9 versus 41.0 trees/ha) and half as many trees greater than 91.4cm dbh per hectare (8.7 versus 16.7 trees/ha) compared to forests in 1911 (Stephens *et al.* 2015).

Mixed-conifer forests in the Sierra Nevada burned every 11 years on average for several centuries prior to Euro-American settlement (Steel *et al.* 2015). These relatively frequent burns prevented the accumulation of fuel on the ground, and limited the intensity of the next fire (Steel *et al.* 2015).

A new approach to remotely sensing wildfire severity

Wildfire severity can be reliably detected remotely by comparing pre- and postfire imagery from sensors aboard the Landsat series of satellites (Eidenshink *et al.* 2007; Miller & Thode 2007): the Thematic Mapper (TM; Landsat 4 and 5), Enhanced Thematic Mapper Plus (ETM+; Landsat 7), and Operational Land Imager (OLI; Landsat 8). Recent advances in radiometric correction post-processing can compensate for various atmospheric distortions and generate more accurate measurements of surface reflectance in narrow wavelength bands spanning the electromagnetic spectrum (Masek *et al.* 2006; Vermote *et al.* 2016; USGS 2017b, a). Landsat satellites image the entire Earth approximately every 16-days and repeat images of the same area




Figure 1: Geographic setting of the study. A) Location of yellow pine/mixed-conifer forests as designated by the Fire Return Interval Departure (FRID) product which, among other things, describes the potential vegetation in an area based on the pre-Euroamerican settlement fire regime. (Data are available for download from <https://www.fs.usda.gov/detail/r5/landmanagement/gis/?cid=STELPRDB5327836>). This image represents a rasterized version of polygons from the FRID database at a 100 m x 100 m pixel resolution, but analyses were performed using a rasterized version at 30 m x 30 m resolution. B) Locations of all fires covering greater than 4 hectares that burned in yellow pine/mixed-conifer forest between 1984 and 2017 in the Sierra Nevada mountain range of California according to the State of California Fire Resource and Assessment Program database, the most comprehensive database of fire perimeters of its kind. (Data are available for download from http://frap.fire.ca.gov/data/frapgisdata-sw-fireperimeters_download). Image represents a rasterized version of polygons from the FRAP database at a 100 m x 100 m pixel resolution. Colors indicate how many fire perimeters overlapped a given pixel within the study time period. C) (red) Locations of composite burn index (CBI) ground plots used to calibrate the remotely sensed measures of severity. (black) Locations of random samples drawn from 972 unique fires depicted in panel B that were in yellow pine/mixed-conifer forest as depicted in panel A, and which were designated as “burned” by exceeding a threshold relative burn ratio (RBR) determined by calibrating the algorithm presented in this study with ground based CBI measurements.

are geometrically coregistered such that overlapping pixels correspond to the same area on the ground. We used Google Earth Engine, a cloud-based geographic information system and image hosting platform, for all image collation and processing in order to leverage the centralized availability of the latest processed satellite images and integrated image processing tools for broad-scale analyses (Gorelick *et al.* 2017).

The base assumption of our new approach to calculating wildfire severity is that each fire’s geographic data and associated attributes are represented by a self-contained “feature”. Many fire datasets (e.g., FRAP, USFS Region 5 Fire Perimeter Data, CBI field plot locations from Zhu *et al.* (2006) and Sikkink *et al.* (2013)) already meet this criteria. In order to achieve a programmatic, automatic assessment of wildfire severity, severity-calculating algorithms must be able to use only the information within each feature. Time efficiencies and data compatibility benefits are attained when those algorithms are applied across an entire feature collection, performing their operation on each feature in turn. At a minimum, our algorithm requires that each feature contain some geographic information (e.g., a fire perimeter or a CBI plot location) and a fire start date (i.e., an “alarm date”).

Fetching and processing pre- and postfire imagery

All Landsat imagery was fetched by “scene”—the atomic unit of image data in the Landsat collection representing an area on the Earth’s surface approximately 170 km long by 183 km wide. For each feature, a collection of Landsat scenes was fetched both before and after the fire by defining a date range to search for imagery. The date range for prefire imagery started one day before each feature’s alarm date and extended backward in time by a user-defined time window. The date range for postfire imagery was exactly one year after the date range for the prefire search (i.e., one year after the day before the fire, extending backward in time by the same time window). We tested 4 time windows: 16, 32, 48, or 64 days which were chosen to ensure that at least 1, 2, 3, or 4 Landsat images, taken on a 16-day interval, were captured by the date ranges (Fig. 2).

The Landsat archive was filtered to generate a prefire image collection comprising only the Landsat scenes depicting some part of the feature geometry and within the prefire date range. A postfire image collection was similarly generated by filtering the Landsat archive by the postfire date range and the feature geometry. The Landsat archive we filtered included imagery from Landsat 4, 5, 7, and 8, so each pre- and postfire image collection may contain a mix of scenes from different satellite sources to enhance coverage.

For each image in the pre- and postfire image collections, we masked pixels that were not clear (i.e., clouds, cloud shadows, snow, and water) and calculated standard indices that capture vegetation cover and fire effects




Figure 2: Schematic for how Landsat imagery was assembled in order to make comparisons between pre- and post-fire conditions. This schematic depicts a 64-day window of image collation prior to the fire which comprise the pre-fire image collection. A similar, 64-day window collection of imagery is assembled one year after the pre-fire image collection.

such as charring: normalized difference vegetation index (NDVI; Eq. 1; Rouse *et al.* (1973)), normalized difference moisture index (NDMI; Eq. 2; Gao (1996)), normalized burn ratio (NBR; Eq. 3; Key & Benson (2006); USGS (2017a); USGS (2017b)), and normalized burn ratio version 2 (NBR2; Eq. 4; USGS (2017a); USGS (2017b); Hawbaker *et al.* (2017)).

$$(1) \text{ ndvi} = (\text{nir} - \text{red}) / (\text{nir} + \text{red})$$

$$(2) \text{ ndmi} = (\text{nir} - \text{swir1}) / (\text{nir} + \text{swir1})$$

$$(3) \text{ nbr} = (\text{nir} - \text{swir2}) / (\text{nir} + \text{swir2})$$

$$(4) \text{ nbr2} = (\text{swir1} - \text{swir2}) / (\text{swir1} + \text{swir2})$$

Where *nir* is the near infrared band (band 4 on Landsat 4, 5, and 7; band 5 on Landsat 8) and *red* is the red band (band 3 on Landsat 4, 5, and 7; band 4 on Landsat 8), *swir1* is the first short wave infrared band (band 5 on Landsat 4, 5, and 7; band 4 on Landsat 8), *swir2* is the second short wave infrared band (band 7 on Landsat 4, 5, 7, and 8)

We summarized each prefire image collection into a single prefire image using a median reducer, which calculated the median of the unmasked values on a per-pixel basis across the stack of images in the prefire collection. We similarly summarized the postfire image collection into a single postfire image.

Calculating wildfire severity

We calculated remotely-sensed wildfire severity using the relative burn ratio (RBR) (Parks *et al.* 2014), the delta normalized burn ratio (dNBR) (Eidenshink *et al.* 2007; Miller & Thode 2007), the relative delta

normalized burn ratio (RdNBR) (Miller & Thode 2007), the delta normalized burn ratio 2 (dNBR2) (Hawbaker *et al.* 2017), the relative delta normalized burn ratio 2 (RdNBR2), and the delta normalized difference vegetation index (dNDVI) (Eidenshink *et al.* 2007). Following the success of the RdNBR metric in other studies, we also calculate an analogous metric using NDVI– the relative delta normalized difference vegetation index (RdNDVI).

We calculated the delta severity indices (dNBR, dNBR2, dNDVI) by subtracting the respective postfire indices from the prefire indices (NBR, NBR2, and NDVI) without multiplying by a rescaling constant (e.g., we did not multiply the result by 1000 as in Miller & Thode (2007); Eq. 5). Following Reilly *et al.* (2017), we chose not to correct the delta indices using a phenological offset value (typically calculated as the delta index in homogenous forest patch outside of the fire perimeter), as our approach implicitly accounts for phenology by incorporating multiple cloud-free images across the same time window both before the fire and one year later.

$$(5) \ dI = I_{\text{prefire}} - I_{\text{postfire}}$$

We calculated the relative delta severity indices, RdNBR and RdNDVI, by scaling the respective delta indices (dNBR and dNDVI) from Eq. 6 by a square root transformation of the absolute value of the prefire index:

$$(6) \ RdI = \frac{dI}{\sqrt{|I_{\text{prefire}}|}}$$

We calculated the relative burn ratio (RBR) following Parks *et al.* (2014) using Eq. 7:

$$(7) \ RBR = \frac{dNBR}{NBR_{\text{prefire}} + 1.001}$$

Example algorithm outputs are shown in Fig. 3.

Calibrating remotely-sensed wildfire severity with field-measured wildfire severity

We calibrated our remotely-sensed measure of wildfire severity with 208 field measures of overstory tree mortality from two previously published studies (Zhu *et al.* 2006; Sikkink *et al.* 2013) (Fig. 4). The Composite Burn Index (CBI) is a metric of change in vegetation across several vertical strata (Key & Benson 2006) and has a long history of use in calibrating remotely-sensed severity data (Miller & Thode 2007; Miller *et al.* 2009; Cansler & McKenzie 2012; Parks *et al.* 2014; Prichard & Kennedy 2014). Following Miller & Thode (2007), Miller *et al.* (2009), and Parks *et al.* (2014), we fit a non-linear model to each remotely-sensed severity metric of the following form:

$$(8) \ \text{remote_severity} = \beta_0 + \beta_1 e^{\beta_2 \text{cbi_overstory}}$$

We fit the model in Eq. 8 for all 7 of our remotely-sensed severity metrics (RBR, dNBR, RdNBR, dNBR2,




Figure 3: Example algorithm outputs for the Hamm Fire of 1987 (top row) and the American Fire of 2013 (bottom row) showing: prefire true color image (left column), postfire true color image (center column), relative burn ratio (RBR) calculation using a 48-day image collation window before the fire and one year later. For visualization purposes, these algorithm outputs have been resampled to a resolution of 100m x 100m from their original resolution of 30m x 30m. Data used for analyses were sampled from the outputs at the original resolution.

RdNBR2, dNDVI, RdNDVI) using 4 different time windows from which to collate satellite imagery (16, 32, 48, and 64 days). Following Cansler & McKenzie (2012) and Parks *et al.* (2014), we used interpolation to extract remotely-sensed severity at the locations of the CBI field plots to better align remote and field measures of severity. We extracted remotely-sensed severity values using both bilinear interpolation, which returns a severity value weighted by the 4 pixel values nearest to the CBI plot location, and bicubic interpolation, which returns a severity value weighted by the 16 pixel values nearest to the CBI plot location. In total, we fit 56 models (7 severity measures, 4 time windows, 2 interpolation methods) and performed five-fold cross validation using the `modelr` and `purrr` packages. To compare goodness of model fits with Miller & Thode (2007), Miller *et al.* (2009), and Parks *et al.* (2014), we report the average R^2 value from the five folds for each of the 56 models but note that R^2 for non-linear regressions do not have the same interpretation that they do for linear regression (i.e., R^2 can be greater than 1 for non-linear regression, so it can't be interpreted as the proportion of variation explained by the model). We used the Relative Burn Ratio (RBR) calculated using bicubic interpolation within a 48-day window as our response variable for analyses of vegetation heterogeneity, as it showed the best correspondence to field severity data measured as average R^2 across the five folds.

Remote sensing other conditions

Variability of vegetation

We used texture analysis to calculate a first order, remotely-sensed measure of local forest variability (Haralick *et al.* 1973; Tuanmu & Jetz 2015). Within a moving square neighborhood window with sides of 90m, 150m, 210m, and 270m (corresponding to a moving neighborhood window of 0.81 ha, 2.25 ha, 4.41 ha, and 7.29 ha), we calculated forest variability for each pixel as the standard deviation of the NDVI values of its neighbors (not including itself) (See Fig. 5).

Topographic conditions

Elevation data were sourced from the Shuttle Radar Topography Mission (Farr *et al.* 2007), a 1-arc second digital elevation model. Slope and aspect were extracted from the digital elevation model. Per-pixel topographic roughness was calculated as the standard deviation of elevation values within a the same kernel sizes as those used for vegetation heterogeneity (approximately 90m, 150m, 210m, and 270m on a side and not including the central pixel). Some work has shown that terrain ruggedness (Holden *et al.* 2009), and particularly coarser-scale terrain ruggedness (Dillon *et al.* 2011), is an important predictor of wildfire severity.




Figure 4: Example of homogenous forest (top row) and heterogenous forest (bottom row) with the same mean NDVI values (~ 0.6). Each column represents heterogeneity measured using a different neighborhood size.

We used the digital elevation model to calculate the potential annual heat load (Eq. 9 at each pixel, which is an integrated measure of latitude, slope, and a folding transformation of aspect about the northeast-southwest line, such that northeast becomes 0 radians and southwest becomes π radians (McCune & Keon (2002) with correction in McCune (2007)):

$$\begin{aligned}
 aspect_{folded} &= \left| \pi - \left| aspect - \frac{5\pi}{4} \right| \right| \\
 &\quad - 1.467 + \\
 &\quad 1.582 * \cos(latitude) \cos(slope) - \\
 \log(pahl) &= 1.5 * \cos(aspect_{folded}) \sin(slope) \sin(latitude) - \\
 &\quad 0.262 * \sin(lat) \sin(slope) + \\
 &\quad 0.607 * \sin(aspect_{folded}) \sin(slope)
 \end{aligned}
 \tag{9}$$

Where $pahl$ is the potential annual heat load, $aspect_{folded}$ is a transformation of aspect in radians, and both $latitude$ and $slope$ are extracted from a digital elevation model with units of radians.

Fire weather conditions

The 100-hour fuel moisture data were sourced from the Gridmet product (Abatzoglou 2013) and were calculated as the median 100-hour fuel moisture for the 3 days prior to the fire.

Modeling the effect of forest variability on severity

We scaled all continuous predictor variables, and treated each individual fire as having a random intercept effect using the mixed effects logistic regression model described in Eq. 10.

$$severity_{i,j} \sim \text{Bern}(\phi_{i,j})$$

$$\beta_0 +$$

$$\beta_{\text{neighborhood_stdev_NDVI}} * \text{neighborhood_stdev_NDVI}_i +$$

$$\beta_{\text{prefire_NDVI}} * \text{prefire_NDVI}_i +$$

$$\beta_{\text{neighborhood_mean_NDVI}} * \text{neighborhood_mean_NDVI}_i +$$

$$\beta_{\text{fm100}} * \text{fm100}_i +$$

$$(10) \quad \text{logit}(\phi_{i,j}) = \beta_{\text{pahl}} * \text{pahl}_i +$$

$$\beta_{\text{topographic_roughness}} * \text{topographic_roughness}_i +$$

$$\beta_{\text{neighborhood_stdev_NDVI} * \text{fm100}} * \text{neighborhood_stdev_NDVI}_i * \text{fm100}_i +$$

$$\beta_{\text{neighborhood_stdev_NDVI} * \text{prefire_NDVI}} * \text{neighborhood_stdev_NDVI}_i * \text{prefire_NDVI}_i +$$

$$\beta_{\text{neighborhood_mean_NDVI} * \text{prefire_NDVI}} * \text{neighborhood_mean_NDVI}_i * \text{prefire_NDVI}_i +$$

$$\gamma_j$$

$$\gamma_j \sim \mathcal{N}(0, \sigma_{\text{fire}})$$

Each neighborhood size (90 m x 90 m, 150 m x 150 m, 210 m x 210 m, and 270 m x 270 m) was substituted in turn for the neighborhood standard deviation of NDVI, neighborhood mean NDVI, and terrain ruggedness covariates to generate a candidate set of 4 models.

To assess the effect of variability of forest structure on the probability of a high-severity wildfire,

To assess the scale at which the forest structure variability effect manifests, we compared the 4 candidate models based on different neighborhood sizes using leave-one-out cross validation (???)

Statistical software and data availability

We used R for all statistical analyses (R Core Team 2018). We used the `brms` package to fit mixed effects models (Bürkner 2017). We used a No U-Turn Sampler (NUTS) with 4 chains and 3000 samples per chain (1500 warmup samples and 1500 posterior samples). Chain convergence was assessed for each estimated parameter by ensuring Rhat values were less than or equal to 1.01.




Figure 5: Three top performing remotely-sensed severity metrics based on 5-fold cross validation (relative burn ratio, 48-day window, bicubic interpolation; relative delta normalized burn ratio, 32-day window, bilinear interpolation; and relative delta normalized difference vegetation index, 48-day window, bilinear interpolation) calculated using new automated image collation algorithms, calibrated to 208 field measures of fire severity (composite burn index). See Supplemental Table 1 for performance of all tested models.

Data are available via the Open Science Framework.

Results

A new approach to remotely sensing wildfire severity

We found that the remotely sensed relative burn ratio (RBR) metric of wildfire severity measured across a 48 day interval prior to the wildfire alarm date correlated best with ground based composite burn index (CBI) measurements of severity (5-fold cross validation $R^2 = 0.82$; Tab. 1). Our method to calculate remotely sensed severity using automated Landsat image fetching performs as well or better than most other reported methods that use hand-curation of Landsat imagery (Edwards *et al.* 2018). Further, several combinations of remotely sensed severity metrics, time windows, and interpolation methods validate well with the ground based severity metrics, including those based on NDVI which is calculated using reflectance in shorter wavelengths than those typically used for measuring severity (Tab. 1). The top three models are depicted in Fig. 6.

Table 1: Comparison of models used to validate and calibrate remotely sensed wildfire severity with ground based composite burn index (CBI) severity sorted in descending order by the R^2 value from a 5-fold cross validation. A total of 56 models were tested representing all possible combinations of 7 different measures of wildfire severity (RBR, dNBR, dNBR2, RdNBR, RdNBR2, dNDVI, and RdNDVI), 4 different time windows in which Landsat imagery was acquired and summarized with a median reducer on a pixel-by-pixel basis (16 days, 32 days, 48 days, and 64 days), and two different interpolation methods (bilinear and bicubic). The three parameters (β_0 , β_1 , and β_2) from the nonlinear model fit described in Eq. 8 are reported. For each model, the value of the remotely sensed wildfire severity measurement corresponding to the lower bounds of 3 commonly used categories of severity are reported ('low' corresponds to a CBI value of 0.1, 'mod' corresponds to a CBI value of 1.25, and 'high' corresponds to a CBI value of 2.25)

Rank	Severity measure	Time window	Interpolation	k-fold R^2	β_0	β_1	β_2	low	mod	high
1	RBR	48	bicubic	0.820	0.014	0.028	1.001	0.045	0.113	0.282
2	RdNBR	32	bilinear	0.813	-0.483	3.061	0.857	2.852	8.450	20.559
3	RdNDVI	48	bilinear	0.809	-2.144	3.273	0.609	1.335	4.867	10.753
4	RBR	32	bilinear	0.807	0.014	0.029	0.985	0.046	0.113	0.280
5	RdNDVI	64	bicubic	0.805	-2.524	3.570	0.590	1.263	4.936	10.929

Based on these model comparisons, we used the relative burn ratio (RBR) calculated using a 48-day time window before the fire and bicubic interpolation as our metric of severity. We created the boolean response variable representing whether the sampled point burned at high severity or not by determining whether the RBR exceeded 0.282, the threshold for high severity derived using the non-linear relationship in Eq. 8 (Tab. 1).

Prefire vegetation density, annual heat load, and topographic roughness effects on wildfire severity

We found that the strongest influence on the probability of a forested area burning at high severity is the density of the vegetation, as measured by the prefire NDVI ($\beta_{\text{prefire_ndvi}}$ = on the log-odds scale; Fig. 7).

For all 4 models using different neighborhood sizes for the heterogeneity and topographic roughness predictors, a greater prefire NDVI led to a greater probability of high severity fire. Potential annual heat load, which integrates aspect, slope, and latitude, also had a strong positive relationship with the probability of a high severity fire (β_{pahl} = Fig. 7).

Areas that were located on southwest facing slopes at lower latitudes tended to be more likely to burn at




Figure 6: Main effects of significant covariates.

high severity. We found no effect of local topographic roughness on wildfire severity at any neighborhood size ($\beta_{\text{topographic_roughness}}$ = Fig. 7).

100 hour fuel moisture effect on wildfire severity

We found a non-linear effect of 100 hour fuel moisture on wildfire severity such that, under normal fuel moisture conditions (100 hour fuel moisture greater than 20th percentile), increasing fuel moisture had a strong negative effect of the probability of a high severity wildfire (β_{fm100} = Fig. 7).

Heterogeneity in vegetation structure effect on wildfire severity

We found strong evidence for an effect of heterogeneity of vegetation structure on the probability of a high severity wildfire.

An increasing heterogeneity of vegetation structure greatly reduced the probability of a high severity wildfire accounting for other variables

Neighborhood size window

Leave-one-out cross validation. Local phenomenon.

Discussion

We developed a new approach to calculating wildfire severity using remotely sensed images from the Landsat series of satellites using a minimal amount of user input— a geometry (i.e., a point location or a perimeter polygon) and a fire start date. We found that the relative burn ratio (RBR) calculated using prefire Landsat images collected over a 48 day period prior to the fire and postfire Landsat images collected over a 48 day period one year after the prefire images validated the best with ground based severity measurements (composite burn index; CBI). We also found that several other remotely sensed measures of severity validated nearly as well with CBI data.

We echo the conclusion of Zhu *et al.* (2006) that the validation of differences between pre- and postfire NDVI to field measured severity data, which uses near infrared reflectance, is comparable to validation using more commonly used severity metrics (e.g., RdNBR and dNBR) that rely on short wave infrared reflectance. One immediately operational implication of this is that the increasing availability of low-cost small unhumanned

aerial systems (sUAS a.k.a. drones) and near infrared detecting imagers (e.g., those used for agriculture monitoring) may be used to measure wildfire severity at very high spatial resolutions.

We used our new approach to calculate wildfire severity for 972 fires that burned in the Sierra Nevada yellow pine/mixed-conifer forest between 1984 and 2016. We additionally calculated 100 hour fuel moisture, local topographic roughness, potential annual heat load, prefire vegetation density, and the local heterogeneity of prefire vegetation density at 4 neighborhood sizes ranging from 0.81 hectares to 7.29 hectares. We modeled the effect of these variables on wildfire severity and found a strong positive relationship with both prefire vegetation density and potential annual heat load. We found no effect of topographic roughness on wildfire severity. We found a negative effect of 100 hour fuel moisture on severity, but only during normal fuel moisture conditions (100 hour fuel moisture greater than 20th percentile). We found a strong negative effect of heterogeneity of vegetation structure on wildfire severity in normal fuel moisture conditions, and a strong positive effect of heterogeneity on severity in extreme fuel moisture conditions.

For similar vegetation density in a given local neighborhood, greater heterogeneity implies more of a mixture of dense patches and sparsely vegetated patches (see Fig. 5).

Caveats

Our method should work best in denser vegetation such as forests, as the signal of a wildfire in other systems can be invisible in a matter of weeks (Goodwin & Collett 2014). This method would also require calibration with field data in other systems, as some severity metrics (such as RBR and RdNBR) have found limited success in other regions (Fernández-García *et al.* 2018).

We have captured a coarse measure of heterogeneity. While we did find that this coarse measure does strongly relate to fire severity, it does not account for fire behavior or spatial pattern forming processes at the individual tree scale. This may still be possible using remotely sensed data at a finer spatial resolution, but with a cost in temporal resolution and time series depth (e.g. NAIP imagery at 1m resolution but with only 3 total images starting in 2008) (Dickinson *et al.* 2016). Additional metrics of heterogeneity such as vegetation patch size distributions or non-vegetated gap size distributions (Malone *et al.* 2018), may also be more tractable using the finer spatial resolution of NAIP imagery.

Translating resistance to long term persistence

Texture analysis has been used to measure habitat heterogeneity in ecology, but has only recently gained recognition for its potential to quantify system resilience (Kéfi *et al.* 2014). For instance, increases in the second angular momentum of the configuration of vegetation patches may represent early warning signs of a catastrophic shift in a system whereby it converts to a vegetation-less desert. The change in this particular texture serves as an indicator of system precariousness because it reflects the spatial process by which the system stabilizes. In the case of desertification as a result of increasing grazing pressure, facilitation is the process driving vegetation patch configurations and the increase in spatial variation of those configurations indicates a breakdown in the process itself as the system moves nearer to a bifurcation point between a vegetated and a non-vegetated state. In our case, we measure heterogeneity as a spatial feature that is part of the feedback loop between disturbance and forest spatial structure, so we gain insight into longer-term patterns by measuring a signature of the pattern forming process itself. More work is needed to verify the degree to which the spatial features of mixed-conifer forests– or the spatial features of the disturbances that affect them– capture the

Conclusions

We encourage researchers and managers to make their ground based severity data available with site location (including datum) and the alarm data for the fire the field data is measuring. Cloud-based GIS, central image hosting, and integration with powerful classification tools are sure to advance our ability to measure wildfire severity remotely, automatically, consistently, and at broad spatial scales. While our contribution here demonstrates that satisfactory validation with ground based measurements is possible using simple and well known calculations, we believe that truly groundbreaking abilities to classify wildfire severity would be possible with more open data sharing of ground based severity measures.

While the severity of a wildfire in any given place may be idiosyncratic and controlled by many variables, it is clear that heterogeneous forest structure generally makes mixed-conifer forest in the Sierra Nevada more resistant to this inevitable disturbance under normal fuel moisture conditions. Because a resistant forest is a resilient forest, heterogeneity in forest structure may increase the probability of long-term forest persistence. Given the opposite effect of heterogeneity in extreme fuel moisture conditions and the normalization of what were once considered extreme fuel moisture conditions in a warming world,

1.

Abatzoglou, J.T. (2013). Development of gridded surface meteorological data for ecological applications and modelling. *International Journal of Climatology*, 33, 121–131.

2.

Ackerly, D.D., Loarie, S.R., Cornwell, W.K., Weiss, S.B., Hamilton, H. & Branciforte, R. *et al.* (2010). The geography of climate change: Implications for conservation biogeography. *Diversity and Distributions*, 16, 476–487.

3.

Agashe, D. (2009). The stabilizing effect of intraspecific genetic variation on population dynamics in novel and ancestral habitats. *The American Naturalist*, 174, 255–67.

4.

Asner, G.P., Brodrick, P.G., Anderson, C.B., Vaughn, N., Knapp, D.E. & Martin, R.E. (2015). Progressive forest canopy water loss during the 2012–2015 California drought. *Proceedings of the National Academy of Sciences*, 2015, 201523397.

5.

Asner, G.P., Martin, R.E., Knapp, D.E., Tupayachi, R., Anderson, C.B. & Sinca, F. *et al.* (2017). Airborne laser-guided imaging spectroscopy to map forest trait diversity and guide conservation. *Science*, 355, 385–389.

6.

Baskett, M.L., Gaines, S.D. & Nisbet, R.M. (2009). Symbiont diversity may help coral reefs survive moderate climate change. *Ecological Applications*, 19, 3–17.

7.

Bastarrika, A., Chuvieco, E. & Martín, M.P. (2011). Mapping burned areas from landsat TM/ETM+ data with a two-phase algorithm: Balancing omission and commission errors. *Remote Sensing of Environment*, 115, 1003–1012.

8.

Boschetti, L., Roy, D.P., Justice, C.O. & Humber, M.L. (2015). MODIS-Landsat fusion for large area 30m burned area mapping. *Remote Sensing of Environment*, 161, 27–42.

9.

Bürkner, P.-C. (2017). brms : An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, 80.

10.

424 Cadotte, M., Albert, C.H. & Walker, S.C. (2013). The ecology of differences: Assessing community assembly
425 with trait and evolutionary distances. *Ecology Letters*, 16, 1234–1244.

426 11.

427 Cansler, C.A. & McKenzie, D. (2012). How robust are burn severity indices when applied in a new region?
428 Evaluation of alternate field-based and remote-sensing methods. *Remote Sensing*, 4, 456–483.

429 12.

430 Cansler, C.A. & McKenzie, D. (2014). Climate, fire size, and biophysical setting control fire severity and
431 spatial pattern in the northern Cascade Range, USA. *Ecological Applications*, 24, 1037–1056.

432 13.

433 Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual Review of Ecology and Systematics*,
434 31, 343–366.

435 14.

436 Clyatt, K.A., Crotteau, J.S., Schaedel, M.S., Wiggins, H.L., Kelley, H. & Churchill, D.J. *et al.* (2016).
437 Historical spatial patterns and contemporary tree mortality in dry mixed-conifer forests. *Forest Ecology and*
438 *Management*, 361, 23–37.

439 15.

440 Collins, B.M., Lydersen, J.M., Everett, R.G., Fry, D.L. & Stephens, S.L. (2015). Novel characterization of
441 landscape-level variability in historical vegetation structure. *Ecological Applications*, 25, 1167–1174.

442 16.

443 Collins, B.M. & Stephens, S.L. (2010). Stand-replacing patches within a 'mixed severity' fire regime:
444 Quantitative characterization using recent fires in a long-established natural fire area. *Landscape Ecology*, 25,
445 927–939.

446 17.

447 Connors, R.W., Trivedi, M.M. & Harlow, C.A. (1984). Segmentation of a high-resolution urban scene using
448 texture operators. *Computer Vision, Graphics, and Image Processing*, 25, 273–310.

449 18.

450 Crowther, T.W., Glick, H.B., Covey, K.R., Bettigole, C., Maynard, D.S. & Thomas, S.M. *et al.* (2015).
451 Mapping tree density at a global scale. *Nature*, 525, 201–205.

452 19.

453 Culbert, P.D., Radeloff, V.C., St-Louis, V., Flather, C.H., Rittenhouse, C.D. & Albright, T.P. *et al.* (2012).

Modeling broad-scale patterns of avian species richness across the Midwestern United States with measures of satellite image texture. *Remote Sensing of Environment*, 118, 140–150.

20.

De Frenne, P., Rodríguez-Sánchez, F., Coomes, D.A., Baeten, L., Verstraeten, G. & Vellend, M. *et al.* (2013). Microclimate moderates plant responses to macroclimate warming. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 18561–5.

21.

De Santis, A., Asner, G.P., Vaughan, P.J. & Knapp, D.E. (2010). Mapping burn severity and burning efficiency in California using simulation models and Landsat imagery. *Remote Sensing of Environment*, 114, 1535–1545.

22.

Dickinson, Y., Pelz, K., Giles, E. & Howie, J. (2016). Have we been successful? Monitoring horizontal forest complexity for forest restoration projects. *Restoration Ecology*, 24, 8–17.

23.

Dillon, G.K., Holden, Z.A., Morgan, P., Crimmins, M.A., Heyerdahl, E.K. & Luce, C.H. (2011). Both topography and climate affected forest and woodland burn severity in two regions of the western US, 1984 to 2006. *Ecosphere*, 2, art130.

24.

Edwards, A.C., Russell-Smith, J. & Maier, S.W. (2018). A comparison and validation of satellite-derived fire severity mapping techniques in fire prone north Australian savannas: Extreme fires and tree stem mortality. *Remote Sensing of Environment*, 206, 287–299.

25.

Eidenshink, J., Schwind, B., Brewer, K., Zhu, Z.-l., Quayle, B. & Howard, S. (2007). A project for monitoring trends in burn severity. *Fire Ecology*, 3, 3–21.

26.

Farr, T., Rosen, P., Caro, E., Crippen, R., Duren, R. & Hensley, S. *et al.* (2007). The shuttle radar topography mission. *Reviews of Geophysics*, 45, 1–33.

27.

Fernández-García, V., Santamarta, M., Fernández-Manso, A., Quintano, C., Marcos, E. & Calvo, L. (2018). Burn severity metrics in fire-prone pine ecosystems along a climatic gradient using Landsat imagery. *Remote*

484 *Sensing of Environment*, 206, 205–217.

485 28.

486 Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T. & Gunderson, L. *et al.* (2004). Regime shifts,
 487 resilience, and biodiversity in ecosystem management. *Annual Review of Ecology, Evolution, and Systematics*,
 488 35, 557–581.

489 29.

490 Ford, K.R., Ettinger, A.K., Lundquist, J.D., Raleigh, M.S. & Hille Ris Lambers, J. (2013). Spatial
 491 heterogeneity in ecologically important climate variables at coarse and fine scales in a high-snow mountain
 492 landscape. *PLoS ONE*, 8, e65008.

493 30.

494 Gao, B.C. (1996). NDWI - A normalized difference water index for remote sensing of vegetation liquid water
 495 from space. *Remote Sensing of Environment*, 58, 257–266.

496 31.

497 Gazol, A. & Camarero, J.J. (2016). Functional diversity enhances silver fir growth resilience to an extreme
 498 drought. *Journal of Ecology*.

499 32.

500 Goodwin, N.R. & Collett, L.J. (2014). Development of an automated method for mapping fire history captured
 501 in Landsat TM and ETM+ time series across Queensland, Australia. *Remote Sensing of Environment*, 148,
 502 206–221.

503 33.

504 Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D. & Moore, R. (2017). Remote Sensing of
 505 Environment Google Earth Engine : Planetary-scale geospatial analysis for everyone. *Remote Sensing of*
 506 *Environment*, 202, 18–27.

507 34.

508 Graham, R.T., McCaffrey, S. & Jain, T.B. (2004). *Science basis for changing forest structure to modify*
 509 *wildfire behavior and severity* (No. April). US Department of Agriculture, Forest Service, Rocky Mountain
 510 Research Station, Fort Collins, CO.

511 35.

512 Gunderson, L.H. (2000). Ecological resilience– in theory and application. *Annual Review of Ecology and*
 513 *Systematics*, 31, 425–439.

36.

Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A. & Tyukavina, A. (2013). High-resolution global maps of 21st-century forest cover change. *Science*, 342, 850–853.

37.

Haralick, R.M., Shanmugam, K. & Dinstein, I. (1973). Textural Features for Image Classification. *IEEE Transactions on Systems, Man, and Cybernetics*, SMC-3, 610–621.

38.

Harvey, B.J., Donato, D.C. & Turner, M.G. (2016). Drivers and trends in landscape patterns of stand-replacing fire in forests of the US Northern Rocky Mountains (1984–2010). *Landscape Ecology*, 31, 2367–2383.

39.

Hawbaker, T.J., Vanderhoof, M.K., Beal, Y.J., Takacs, J.D., Schmidt, G.L. & Falgout, J.T. *et al.* (2017). Mapping burned areas using dense time-series of Landsat data. *Remote Sensing of Environment*, 198, 504–522.

40.

Holden, Z.A., Morgan, P. & Evans, J.S. (2009). A predictive model of burn severity based on 20-year satellite-inferred burn severity data in a large southwestern US wilderness area. *Forest Ecology and Management*, 258, 2399–2406.

41.

Holling, C.S. (1973). Resilience and Stability of Ecological Systems. *Annual Review of Ecology and Systematics*, 4, 1–23.

42.

Huang, Q., Swatantran, A., Dubayah, R. & Goetz, S.J. (2014). The influence of vegetation height heterogeneity on forest and woodland bird species richness across the United States. *PLoS ONE*, 9.

43.

Key, C.H. & Benson, N.C. (2006). Landscape assessment: Sampling and analysis methods. *USDA Forest Service General Technical Report RMRS-GTR-164-CD*, 1–55.

44.

Kéfi, S., Guttal, V., Brock, W.A., Carpenter, S.R., Ellison, A.M. & Livina, V.N. *et al.* (2014). Early warning signals of ecological transitions: Methods for spatial patterns. *PLoS ONE*, 9, 10–13.

45.

Kolden, C.A., Smith, A.M.S. & Abatzoglou, J.T. (2015). Limitations and utilisation of Monitoring Trends in

Burn Severity products for assessing wildfire severity in the USA. *International Journal of Wildland Fire*, 24, 1023–1028.

46.

Kotliar, N.B. & Wiens, J. a. (1990). Multiple Scales of Patchiness and Patch Structure: A Hierarchical Framework for the Study of Heterogeneity. *Oikos*, 59, 253–260.

47.

Larson, A.J. & Churchill, D. (2012). Tree spatial patterns in fire-frequent forests of western North America, including mechanisms of pattern formation and implications for designing fuel reduction and restoration treatments. *Forest Ecology and Management*, 267, 74–92.

48.

Lenoir, J., Graae, B.J., Aarrestad, P.A., Alsos, I.G., Armbruster, W.S. & Austrheim, G. *et al.* (2013). Local temperatures inferred from plant communities suggest strong spatial buffering of climate warming across Northern Europe. *Global Change Biology*, 19, 1470–1481.

49.

Malone, S.L., Fornwalt, P.J., Battaglia, M.A., Chambers, M.E., Iniguez, J.M. & Sieg, C.H. (2018). Mixed-severity fire fosters heterogeneous spatial patterns of conifer regeneration in a dry conifer forest. *Forests*, 9.

50.

Masek, J.G., Vermote, E.F., Saleous, N.E., Wolfe, R., Hall, F.G. & Huemmrich, K.F. *et al.* (2006). A Landsat Surface Reflectance Dataset. *IEEE Geoscience and Remote Sensing Letters*, 3, 68–72.

51.

McCune, B. (2007). Improved estimates of incident radiation and heat load using non-parametric regression against topographic variables. *Journal of Vegetation Science*, 18, 751–754.

52.

McCune, B. & Keon, D. (2002). Equations for potential annual direct incident radiation and heat load. *Journal of Vegetation Science*, 13, 603–606.

53.

Millar, C.I. & Stephenson, N.L. (2015). Temperate forest health in an era of emerging megadisturbance. *Science*, 349, 823–826.

54.

Miller, J.D., Knapp, E.E., Key, C.H., Skinner, C.N., Isbell, C.J. & Creasy, R.M. *et al.* (2009). Calibration and validation of the relative differenced Normalized Burn Ratio (RdNBR) to three measures of fire severity in the Sierra Nevada and Klamath Mountains, California, USA. *Remote Sensing of Environment*, 113, 645–656.

55.

Miller, J.D. & Thode, A.E. (2007). Quantifying burn severity in a heterogeneous landscape with a relative version of the delta Normalized Burn Ratio (dNBR). *Remote Sensing of Environment*, 109, 66–80.

56.

Moritz, M.A., Morais, M.E., Summerell, L.A., Carlson, J.M. & Doyle, J. (2005). Wildfires, complexity, and highly optimized tolerance. *Proceedings of the National Academy of Sciences*, 102, 17912–7.

57.

Näsi, R., Honkavaara, E., Lyytikäinen-Saarenmaa, P., Blomqvist, M., Litkey, P. & Hakala, T. *et al.* (2015). Using UAV-based photogrammetry and hyperspectral imaging for mapping bark beetle damage at tree-level. *Remote Sensing*, 7, 15467–15493.

58.

North, M.P., Stephens, S.L., Collins, B.M., Agee, J.K., Aplet, G. & Franklin, J.F. *et al.* (2015). Reform forest fire management. *Science*, 349, 1280–1281.

59.

North, M., Stine, P., Hara, K.O., Zielinski, W. & Stephens, S. (2009). An Ecosystem Management Strategy for Sierran Mixed- Conifer Forests. *General Technical Report PSW-GTR-220*, 1–49.

60.

Park Williams, A., Allen, C.D., Macalady, A.K., Griffin, D., Woodhouse, C.A. & Meko, D.M. *et al.* (2012). Temperature as a potent driver of regional forest drought stress and tree mortality. *Nature Climate Change*, 3, 292–297.

61.

Parks, S.A., Dillon, G.K. & Miller, C. (2014). A new metric for quantifying burn severity: The relativized burn ratio. *Remote Sensing*, 6, 1827–1844.

62.

Prichard, S.J. & Kennedy, M.C. (2014). Fuel treatments and landform modify landscape patterns of burn severity in an extreme fire event. *Ecological Applications*, 24, 571–590.

63.

604 Questad, E.J. & Foster, B.L. (2008). Coexistence through spatio-temporal heterogeneity and species sorting
 605 in grassland plant communities. *Ecology Letters*, 11, 717–726.
 606 64.
 607 R Core Team. (2018). *R: A language and environment for statistical computing*. <http://www.r-project.org/>.
 608 R Foundation for Statistical Computing, Vienna, Austria.
 609 65.
 610 Raffa, K.F., Aukema, B., Bentz, B.J., Carroll, A., Erbilgin, N. & Herms, D.A. *et al.* (2009). A literal use of
 611 'forest health' safeguards against misuse and misapplication. *Journal of Forestry*, 276–277.
 612 66.
 613 Raffa, K.F., Aukema, B.H., Bentz, B.J., Carroll, A.L., Hicke, J.A. & Turner, M.G. *et al.* (2008). Cross-scale
 614 drivers of natural disturbances prone to anthropogenic amplification: The dynamics of bark beetle eruptions.
 615 *BioScience*, 58, 501.
 616 67.
 617 Reilly, M.J., Dunn, C.J., Meigs, G.W., Spies, T.A., Kennedy, R.E. & Bailey, J.D. *et al.* (2017). Contemporary
 618 patterns of fire extent and severity in forests of the Pacific Northwest, USA (1985-2010). *Ecosphere*, 8.
 619 68.
 620 Reusch, T.B.H., Ehlers, A., Hämmerli, A. & Worm, B. (2005). Ecosystem recovery after climatic extremes
 621 enhanced by genotypic diversity. *Proceedings of the National Academy of Sciences*, 102, 2826–2831.
 622 69.
 623 Rouse, J.W., Hass, R.H., Schell, J. & Deering, D. (1973). Monitoring vegetation systems in the great plains
 624 with ERTS. *Third Earth Resources Technology Satellite (ERTS) symposium*, 1, 309–317.
 625 70.
 626 Scholl, A.E. & Taylor, A.H. (2010). Fire regimes, forest change, and self-organization in an old-growth
 627 mixed-conifer forest, Yosemite National Park, USA. *Ecological Applications*, 20, 362–380.
 628 71.
 629 Sikkink, P.G., Dillon, G.K., Keane, R.E., Morgan, P., Karau, E.C. & Holden, Z.A. *et al.* (2013). *Composite*
 630 *Burn Index (CBI) data and field photos collected for the FIRESEV project, western United States*. Forest
 631 Service Research Data Archive, Fort Collins, CO.
 632 72.
 633 Steel, Z.L., Safford, H.D. & Viers, J.H. (2015). The fire frequency-severity relationship and the legacy of fire

634 suppression in California forests. *Ecosphere*, 6, 1–23.

635 73.

636 Stein, A., Gerstner, K. & Kreft, H. (2014). Environmental heterogeneity as a universal driver of species
637 richness across taxa, biomes and spatial scales. *Ecology Letters*, 17, 866–880.

638 74.

639 Stephens, S.L. & Collins, B.M. (2004). Fire regimes of mixed conifer forests in the North-Central Sierra
640 Nevada at multiple scales. *Northwest Science*, 78, 12–23.

641 75.

642 Stephens, S.L., Fry, D.L. & Franco-Vizcaíno, E. (2008). Wildfire and spatial patterns in forests in northwestern
643 Mexico: The United States wishes it had similar fire problems. *Ecology and Society*.

644 76.

645 Stephens, S.L., Lydersen, J.M., Collins, B.M., Fry, D.L. & Meyer, M.D. (2015). Historical and current
646 landscape-scale ponderosa pine and mixed conifer forest structure in the Southern Sierra Nevada. *Ecosphere*,
647 6, 1–63.

648 77.

649 Sugihara, N.G. & Barbour, M.G. (2006). Fire and California vegetation. In: *Fire in california's ecosystems*
650 (eds. Sugihara, N.G., Van Wagtenonk, J.W., Shaffer, K.E., Fites-Kaufman, J. & Thode, A.E.). University
651 of California Press, Berkeley; Los Angeles, CA, USA, pp. 1–9.

652 78.

653 Tilman, D. (1994). Competition and biodiversity in spatially structured habitats. *Ecology*, 75, 2–16.

654 79.

655 Trumbore, S., Brando, P. & Hartmann, H. (2015). Forest health and global change. *Science*, 349.

656 80.

657 Tuanmu, M.-N. & Jetz, W. (2015). A global, remote sensing-based characterization of terrestrial habitat
658 heterogeneity for biodiversity and ecosystem modelling. *Global Ecology and Biogeography*, n/a–n/a.

659 81.

660 USGS. (2017a). Product Guide: Landat 8 Surface Reflectance Code (LaSRC) Product. *USGS Professional*
661 *Paper*, 4.2.

662 82.

663 USGS. (2017b). Product Guide: Landsat 4-7 Surface Reflectance (LEDAPS) Product. *USGS Professional*

664 *Paper*, 8, 38.

665 83.

666 Veraverbeke, S. & Hook, S.J. (2013). Evaluating spectral indices and spectral mixture analysis for assessing
667 fire severity, combustion completeness and carbon emissions. *International Journal of Wildland Fire*, 22,
668 707–720.

669 84.

670 Vermote, E., Justice, C., Claverie, M. & Franch, B. (2016). Preliminary analysis of the performance of the
671 Landsat 8/OLI land surface reflectance product. *Remote Sensing of Environment*, 185, 46–56.

672 85.

673 Virah-Sawmy, M., Willis, K.J. & Gillson, L. (2009). Threshold response of Madagascar’s littoral forest to
674 sea-level rise. *Global Ecology and Biogeography*, 18, 98–110.

675 86.

676 Walker, B., Holling, C.S., Carpenter, S.R. & Kinzig, A. (2004). Resilience, adaptability, and transformability
677 in social-ecological systems. *Ecology and Society*, 9, 5.

678 87.

679 Westerling, A.L., Hidalgo, H.G., Cayan, D.R. & Swetnam, T.W. (2006). Warming and earlier spring increase
680 western U.S. forest wildfire activity. *Science*, 313, 940–943.

681 88.

682 Wood, E.M., Pidgeon, A.M., Radeloff, V.C. & Keuler, N.S. (2012). Image texture as a remotely sensed
683 measure of vegetation structure. *Remote Sensing of Environment*, 121, 516–526.

684 89.

685 Young, D.J.N., Stevens, J.T., Earles, J.M., Moore, J., Ellis, A. & Jirka, A.L. *et al.* (2017). Long-term climate
686 and competition explain forest mortality patterns under extreme drought. *Ecology Letters*, 20, 78–86.

687 90.

688 Zhu, Z., Key, C., Ohlen, D. & Benson, N. (2006). Evaluate Sensitivities of Burn-Severity Mapping Algorithms
689 for Different Ecosystems and Fire Histories in the United States. *Final Report to the Joint Fire Science*
690 *Program, Project JFSP 01-1-4-12*, 1–35.