

Invariant Geometric Structures and Chern numbers of G_2 Flag Manifolds

Daniël Thung

LMU München

September 28, 2017

Outline

- 1 Generalized flag manifolds
- 2 G_2 and its flag manifolds
- 3 The twistor space
- 4 The quadric

Generalized flag manifolds

Definition

A *flag* is a strictly increasing sequence

$$\{0\} = V_0 \subset V_1 \subset V_2 \subset \cdots \subset V_k = V \quad k \leq n$$

of subspaces of an n -dimensional vector space V (e.g. \mathbb{C}^n). Set $d_i = \dim V_i$; then (d_1, \dots, d_{k-1}) is the signature of the flag. If $d_i = i$, we call the flag *complete* and otherwise we have a *partial* flag.

Generalized flag manifolds

Example (Flag manifolds of flags in \mathbb{C}^n)

a) The Grassmannian of k -planes parametrizes flags of signature (k) :

$$\mathrm{Gr}_k(\mathbb{C}^n) \cong \frac{U(n)}{U(k) \times U(n-k)} \cong \frac{SU(n)}{S(U(k) \times U(n-k))}$$

Generalized flag manifolds

Example (Flag manifolds of flags in \mathbb{C}^n)

a) The Grassmannian of k -planes parametrizes flags of signature (k) :

$$\mathrm{Gr}_k(\mathbb{C}^n) \cong \frac{U(n)}{U(k) \times U(n-k)} \cong \frac{SU(n)}{S(U(k) \times U(n-k))}$$

b) The *complete* flag manifold is $U(n)/T^n \cong SU(n)/T^{n-1}$.

Generalized flag manifolds

In general, any *flag manifold* of flags in \mathbb{C}^n is of the form

$$\frac{SU(n)}{S(U(r_1) \times \dots \times U(r_k))}$$

where $\{r_1, \dots, r_k\}$ is an ordered partition of n . The isotropy subgroup is the centralizer of a torus $T^{k-1} \subset SU(n)$.

Generalized flag manifolds

In general, any *flag manifold* of flags in \mathbb{C}^n is of the form

$$\frac{SU(n)}{S(U(r_1) \times \dots \times U(r_k))}$$

where $\{r_1, \dots, r_k\}$ is an ordered partition of n . The isotropy subgroup is the centralizer of a torus $T^{k-1} \subset SU(n)$.

This naturally generalizes to:

Definition

A *generalized flag manifold* is a homogeneous space of the form $G/C(T)$, where G is a compact, connected and semisimple Lie group, and $C(T)$ is the centralizer of a torus $T \subset G$.

Generalized flag manifolds

Generalized flag manifolds carry interesting invariant complex-geometric structures:

Generalized flag manifolds

Generalized flag manifolds carry interesting invariant complex-geometric structures:

Theorem (Borel [1], Koszul [2], Matsushima [3])

Every generalized flag manifold admits an invariant complex structure and a unique (up to scaling) compatible invariant Kähler-Einstein metric, which has positive scalar curvature. Conversely, any compact, simply connected homogeneous space that admits an invariant Kähler metric is isomorphic to a generalized flag manifold.

Generalized flag manifolds

Generalized flag manifolds carry interesting invariant complex-geometric structures:

Theorem (Borel [1], Koszul [2], Matsushima [3])

Every generalized flag manifold admits an invariant complex structure and a unique (up to scaling) compatible invariant Kähler-Einstein metric, which has positive scalar curvature. Conversely, any compact, simply connected homogeneous space that admits an invariant Kähler metric is isomorphic to a generalized flag manifold.

Furthermore, certain examples are known to carry multiple invariant almost complex structures with distinct Chern numbers (cf. Borel & Hirzebruch [4]).

Generalized flag manifolds

Invariant structures are traditionally studied by means of Lie theory, obscuring the geometric meaning of many results.

Generalized flag manifolds

Invariant structures are traditionally studied by means of Lie theory, obscuring the geometric meaning of many results.

Objective

Study generalized flag manifolds *geometrically* and give a concrete interpretation of their invariant geometric structures.

Generalized flag manifolds

Invariant structures are traditionally studied by means of Lie theory, obscuring the geometric meaning of many results.

Objective

Study generalized flag manifolds *geometrically* and give a concrete interpretation of their invariant geometric structures.

This program was carried out for $SU(n)/S(U(n-2) \times U(1) \times U(1))$ by Kotschick & Terzić [6].

G_2 and its flag manifolds

The exceptional Lie group G_2 is the automorphism group of the octonions

- ①. Its generalized flag manifolds fit into the following diagram:

G_2 and its flag manifolds

The exceptional Lie group G_2 is the automorphism group of the octonions

①. Its generalized flag manifolds fit into the following diagram:

$$G_2/T^2$$

G_2 and its flag manifolds

The exceptional Lie group G_2 is the automorphism group of the octonions

①. Its generalized flag manifolds fit into the following diagram:

$$\begin{array}{ccc} & & G_2/T^2 \\ & \swarrow & \\ \mathbb{CP}^1 & & \\ & \searrow & \\ & & G_2/U(2)_- \end{array}$$


G_2 and its flag manifolds

The exceptional Lie group G_2 is the automorphism group of the octonions
①. Its generalized flag manifolds fit into the following diagram:

$$\begin{array}{ccc} & & G_2/T^2 \\ & \swarrow \mathbb{CP}^1 & \\ & & G_2/U(2)_- \\ & \swarrow \mathbb{CP}^2 & \\ G_2/SU(3) \cong S^6 & & \end{array}$$


G_2 and its flag manifolds

The exceptional Lie group G_2 is the automorphism group of the octonions
①. Its generalized flag manifolds fit into the following diagram:




G_2 and its flag manifolds

The exceptional Lie group G_2 is the automorphism group of the octonions
①. Its generalized flag manifolds fit into the following diagram:



G_2 and its flag manifolds

The exceptional Lie group G_2 is the automorphism group of the octonions
①. Its generalized flag manifolds fit into the following diagram:



The twistor space

$\pi : Z \rightarrow G_2/SO(4)$ exhibits Z as the *twistor space* over $G_2/SO(4)$, which is a quaternionic Kähler manifold.

The twistor space

$\pi : Z \rightarrow G_2/SO(4)$ exhibits Z as the *twistor space* over $G_2/SO(4)$, which is a quaternionic Kähler manifold.

Definition

A quaternionic Kähler (QK) manifold is an (oriented) Riemannian manifold of dimension $4n$, $n \geq 2$, whose holonomy group is contained in the subgroup $Sp(n) \cdot Sp(1)$ of $SO(4n)$.

The twistor space

$\pi : Z \rightarrow G_2/SO(4)$ exhibits Z as the *twistor space* over $G_2/SO(4)$, which is a quaternionic Kähler manifold.

Definition

A quaternionic Kähler (QK) manifold is an (oriented) Riemannian manifold of dimension $4n$, $n \geq 2$, whose holonomy group is contained in the subgroup $Sp(n) \cdot Sp(1)$ of $SO(4n)$.

Reduced holonomy implies curvature restrictions: QK manifolds are Einstein.

The twistor space

Any QK manifold M carries a rank 3 subbundle E of $\text{End } TM$, locally spanned by a quaternionic triple $\{I, J, IJ = K\}$.

The twistor space

Any QK manifold M carries a rank 3 subbundle E of $\text{End } TM$, locally spanned by a quaternionic triple $\{I, J, IJ = K\}$.

Definition (Salamon [7])

The total space $S(E)$ of the sphere bundle of E is called the *twistor space* over M .

The twistor space

Any QK manifold M carries a rank 3 subbundle E of $\text{End } TM$, locally spanned by a quaternionic triple $\{I, J, IJ = K\}$.

Definition (Salamon [7])

The total space $S(E)$ of the sphere bundle of E is called the *twistor space* over M .

A point $z = \alpha I + \beta J + \gamma K \in S(E)$ corresponds to an orthogonal complex structure on $T_{\pi(z)}M$.

The twistor space

Theorem (Salamon [7], Bérard-Bergery [8])

- (i) $S(E)$ admits a natural (integrable) complex structure.
- (ii) If M has positive scalar curvature, then $S(E)$ admits a Kähler-Einstein metric with positive scalar curvature.

The twistor space

Theorem (Salamon [7], Bérard-Bergery [8])

- (i) $S(E)$ admits a natural (integrable) complex structure.
- (ii) If M has positive scalar curvature, then $S(E)$ admits a Kähler-Einstein metric with positive scalar curvature.

Proof of (i).

$TS(E) \cong \mathcal{V} \oplus \mathcal{H}$, where $\mathcal{H}_z \cong T_{\pi(z)}M$; we define a complex structure $J = J_v \oplus J_h$ as follows: $J_v = J_{\mathbb{CP}^1}^{\text{std}}$, while $z \in S(E)$ is the image of J_h under the identification $\mathcal{H}_z \cong T_{\pi(z)}M$. The Nijenhuis tensor is explicitly shown to vanish. □

The twistor space: Rigidity of the Kähler structure

Using the Gysin sequence we determine the cohomology ring and Pontryagin classes of $Z = G_2/U(2)_+$. Additively, the cohomology is that of \mathbb{CP}^5 , but the ring structure is different.

The twistor space: Rigidity of the Kähler structure

Using the Gysin sequence we determine the cohomology ring and Pontryagin classes of $Z = G_2/U(2)_+$. Additively, the cohomology is that of \mathbb{CP}^5 , but the ring structure is different.

Theorem

Let X be a Kähler manifold. If it is homeomorphic to Z , then it is biholomorphic to Z

The twistor space: Rigidity of the Kähler structure

Using the Gysin sequence we determine the cohomology ring and Pontryagin classes of $Z = G_2/U(2)_+$. Additively, the cohomology is that of \mathbb{CP}^5 , but the ring structure is different.

Theorem

Let X be a Kähler manifold. If it is homeomorphic to Z , then it is biholomorphic to Z

This is the exact analog of rigidity results of Hirzebruch & Kodaira [9] (and Yau [10]) for \mathbb{CP}^n , and Brieskorn [11] for Q_n ($n \geq 3$).

The twistor space: Rigidity of the Kähler structure

Sketch of Proof.

We have $c_1(X) = d \cdot g_2$, where g_2 is the positive generator of $H^2(X; \mathbb{Z})$. The Chern number $c_1 c_4[X]$ is fixed by $h^{p,q}(X) = h^{p,q}(\mathbb{CP}^5)$, hence $c_1 c_4[X] = 90$. Thus, d divides 90. Every possibility except $d = 3$ is ruled out case-by-case, using the Pontryagin classes as well as the Todd genus, which is also determined by the Hodge numbers.

The twistor space: Rigidity of the Kähler structure

Sketch of Proof.

We have $c_1(X) = d \cdot g_2$, where g_2 is the positive generator of $H^2(X; \mathbb{Z})$. The Chern number $c_1 c_4[X]$ is fixed by $h^{p,q}(X) = h^{p,q}(\mathbb{CP}^5)$, hence $c_1 c_4[X] = 90$. Thus, d divides 90. Every possibility except $d = 3$ is ruled out case-by-case, using the Pontryagin classes as well as the Todd genus, which is also determined by the Hodge numbers.

Once $c_1(X)$ is fixed, Mukai's classification of Fano manifolds of coindex 3 finishes the proof. □

The twistor space: Invariant almost complex structures

The twistor space: Invariant almost complex structures

The second invariant almost complex structure is obtained by “flipping the fiber”: $J' = (-J_v) \oplus J_h$. The Chern numbers are (compare GNO [12]):

The twistor space: Invariant almost complex structures


The second invariant almost complex structure is obtained by “flipping the fiber”: $J' = (-J_v) \oplus J_h$. The Chern numbers are (compare GNO [12]):

Proposition

The Chern numbers of the two invariant almost complex structures on the twistor space are:

<i>Chern Number</i>	<i>Z</i>	<i>N</i>
c_5	6	6
c_1^5	4374	-18
$c_1^3 c_2$	2106	-6
$c_1^2 c_3$	594	18
$c_1 c_4$	90	18
$c_1 c_2^2$	1014	-2
$c_2 c_3$	286	6

Reminder: G_2 flag manifolds



The quadric

$G_2/U(2)_- = Q$ is the space of oriented 2-planes in $\text{Im } \mathbb{O} \cong \mathbb{R}^7$.

The quadric

$G_2/U(2)_- = Q$ is the space of oriented 2-planes in $\operatorname{Im} \mathbb{O} \cong \mathbb{R}^7$.

Let $\{e_1, e_2\}$ be a positive, orthonormal basis for $P \in Q$ (unique up to $U(1)$ -transformation). \mathbb{C} -linearly extending the standard inner product $(-, -)_{\mathbb{R}^7}$ to \mathbb{C}^7 , set $f(z) = (z, z) = \sum z_j^2$. Then $f(e_1 + ie_2) = 0$.

The quadric

$G_2/U(2)_- = Q$ is the space of oriented 2-planes in $\operatorname{Im} \mathbb{O} \cong \mathbb{R}^7$.

Let $\{e_1, e_2\}$ be a positive, orthonormal basis for $P \in Q$ (unique up to $U(1)$ -transformation). \mathbb{C} -linearly extending the standard inner product $(-, -)_{\mathbb{R}^7}$ to \mathbb{C}^7 , set $f(z) = (z, z) = \sum z_j^2$. Then $f(e_1 + ie_2) = 0$.

This yields a diffeomorphism

$$Q \cong \{f(z) = 0\} \subset \mathbb{CP}^6$$

The quadric

$G_2/U(2)_- = Q$ is the space of oriented 2-planes in $\operatorname{Im} \mathbb{O} \cong \mathbb{R}^7$.

Let $\{e_1, e_2\}$ be a positive, orthonormal basis for $P \in Q$ (unique up to $U(1)$ -transformation). \mathbb{C} -linearly extending the standard inner product $(-, -)_{\mathbb{R}^7}$ to \mathbb{C}^7 , set $f(z) = (z, z) = \sum z_j^2$. Then $f(e_1 + ie_2) = 0$.

This yields a diffeomorphism

$$Q \cong \{f(z) = 0\} \subset \mathbb{CP}^6$$

The invariant complex structure and Kähler-Einstein metric are inherited from \mathbb{CP}^6 : The restriction of the Fubini-Study metric is Kähler-Einstein, and even $SO(7)$ -invariant.

The quadric: Invariant almost complex structures

Equip S^6 with its G_2 -invariant almost complex structure, given by $J_x(v) = x \cdot v$. Then $\mathbb{P}(TS^6)$ inherits an invariant almost complex structure.

The quadric: Invariant almost complex structures

Equip S^6 with its G_2 -invariant almost complex structure, given by $J_x(v) = x \cdot v$. Then $\mathbb{P}(TS^6)$ inherits an invariant almost complex structure. The identification

complex line \longleftrightarrow oriented 2-plane

induces a diffeomorphism $\mathbb{P}(TS^6) \cong Q \cong \mathbb{P}(T^*S^6)$.

The quadric: Invariant almost complex structures

Equip S^6 with its G_2 -invariant almost complex structure, given by $J_x(v) = x \cdot v$. Then $\mathbb{P}(TS^6)$ inherits an invariant almost complex structure. The identification

complex line \longleftrightarrow oriented 2-plane

induces a diffeomorphism $\mathbb{P}(TS^6) \cong Q \cong \mathbb{P}(T^*S^6)$.

Remark

This shows that a complex structure on S^6 gives rise to (at least two) non-standard complex structures on the quadric $Q \subset \mathbb{CP}^6$.


The quadric: Invariant almost complex structures

We have now found three invariant almost complex structures on $G_2/U(2)_-$, out of a total of four.

The quadric: Invariant almost complex structures

We have now found three invariant almost complex structures on $G_2/U(2)_-$, out of a total of four.

The fourth is obtained from the quadric by flipping the fiber over S^6 . In fact, all four invariant almost complex structures are related by flips:



The quadric: Chern numbers

The invariant almost complex structures are distinguished by their Chern numbers (compare GNO [12]):

Proposition

The Chern numbers of the four invariant almost complex structures on $G_2/U(2)_-$ are:

<i>Chern Number</i>	Q	$\mathbb{P}(TS^6)$	$\mathbb{P}(T^*S^6)$	X
c_5	6	6	6	6
c_1^5	6250	-486	486	-2
$c_1^3 c_2$	2750	-162	162	2
$c_1^2 c_3$	650	18	18	2
$c_1 c_4$	90	18	18	-6
$c_1 c_2^2$	1210	-54	54	-2
$c_2 c_3$	286	6	6	-2

References I

- [1] A. Borel. “Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts”. *Ann. of Math.* (2) 57 (1953), pp. 115–207.
- [2] J. L. Koszul. “Sur la forme hermitienne canonique des espaces homogènes complexes”. *Canad. J. Math.* 7 (1955), pp. 562–576.
- [3] Y. Matsushima. “Sur les espaces homogènes kählériens d’un groupe de Lie réductif”. *Nagoya Math. J.* 11 (1957), pp. 53–60.
- [4] A. Borel and F. Hirzebruch. “Characteristic Classes and Homogeneous Spaces. I–III”. *Amer. J. Math.* 80–82 (1958–1960), pp. 458–538, 315–382, 491–504.
- [5] F. Hirzebruch. “The Projective Tangent Bundles of a Complex Three-Fold”. *Pure Appl. Math. Q.* 1 (2005), pp. 441–448.

References II

- [6] D. Kotschick and S. Terzić. “Chern Numbers and the Geometry of Partial Flag Manifolds”. *Comment. Math. Helv.* 84 (2009), pp. 587–616.
- [7] S. Salamon. “Quaternionic Kähler Manifolds”. *Invent. Math.* 67 (1982), pp. 143–171.
- [8] A. L. Besse. *Einstein Manifolds*. Springer-Verlag, Berlin, 1987.
- [9] F. Hirzebruch and K. Kodaira. “On the Complex Projective Spaces”. *J. Math. Pures Appl.* (9) 36 (1957), pp. 201–216.
- [10] S. T. Yau. “Calabi’s Conjecture and Some New Results in Algebraic Geometry”. *Proc. Nat. Acad. Sci. USA* 74 (1977), pp. 1798–1799.
- [11] E. Brieskorn. “Ein Satz über die komplexen Quadriken”. *Math. Ann.* 155 (1964), pp. 184–193.

- [12] L. Grama, C. J. C. Negreiros, and A. R. Oliveira. “Invariant Almost Complex Geometry on Flag Manifolds: Geometric Formality and Chern Numbers”. *Ann. Mat. Pura Appl. (4)* 196 (Feb. 1, 2017), pp. 165–200.