

# Test 3 Notes

Daniel Palma

November 14, 2023

## Contents

<b>1</b>	<b>Change of Variables (Jacobians)</b>	<b>3</b>
1.1	Jacobian . . . . .	3
1.2	Change of Variables in a Double Integrals . . . . .	4
1.2.1	Triple Integrals . . . . .	4
<b>2</b>	<b>Vector Fields</b>	<b>5</b>
2.1	Gradient Fields . . . . .	5
<b>3</b>	<b>Line Integrals</b>	<b>6</b>
<b>4</b>	<b>The Fundamental Theorem of Line Integrals</b>	<b>7</b>
<b>5</b>	<b>Green's Theorem</b>	<b>8</b>
<b>6</b>	<b>Curl and Divergence</b>	<b>9</b>
<b>7</b>	<b>Parametric Surfaces</b>	<b>10</b>
<b>8</b>	<b>Surface Integrals</b>	<b>11</b>

# 1 Change of Variables (Jacobians)

In one-dimensional calculus we often use a change of variable to simplify an integrals.

A change of variables can also be useful in double integrals. We have already seen one example of this: conversion to polar coordinates. The new variables  $r$  and  $\theta$  are related to the old variables  $x$  and  $y$  by the equations

$$x = r \cos \theta \quad y = r \sin \theta$$

and the change of variables formula can be written as

$$\iint_R f(x, y) dA = \iint_S f(r \cos \theta, r \sin \theta) r \, dr \, d\theta$$

Where  $S$  is the region in the  $r\theta$ -plane that corresponds to the region  $R$  in the  $xy$ -plane

More generally, we consider a change of variables that is given by a **transformation**  $T$  from the  $uv$ -plane to the  $xy$ -plane:

$$x = g(u, v) \quad y = h(u, v)$$

or, as we sometimes write,

$$x = x(u, v) \quad y = y(u, v)$$

We usually assume that  $T$  is a  **$\mathbf{C}^1$  transformation**, which means that  $g$  and  $h$  have continuous first-order partial derivatives.

A transformation  $T$  is really just a function whose domain and range are both subsets of  $\mathbb{R}^2$ . If  $T(u_1, v_1) = (x_1, y_1)$ , then the point  $(x_1, y_1)$  is called the **image** of the point  $(u_1, v_1)$ . If no two points have the same image,  $T$  is called **one-to-one**.

If  $T$  is a one-to-one transformation, then it has an **inverse transformation**  $T^{-1}$  from the  $xy$ -plane to the  $uv$ -plane and it may be possible to solve

$$x = x(u, v) \quad y = y(u, v)$$

for  $u$  and  $v$  in terms of  $x$  and  $y$ :

$$u = G(x, y) \quad v = H(x, y)$$

## 1.1 Jacobian

The **Jacobian** of the transformation  $T$  is given by  $x = g(u, v)$  and  $y = h(u, v)$  is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

## 1.2 Change of Variables in a Double Integrals

Suppose that  $T$  is a  $C^1$  transformation whose Jacobian is nonzero and that  $T$  maps a region  $S$  in the  $uv$ - plane onto a region  $R$  in the  $xy$ -plane. Suppose that  $f$  is continuous on  $R$  and that  $R$  and  $S$  are type I or type II plane regions. Suppose also that  $T$  is one-to-one, except perhaps on the boundary of  $S$ . Then

$$\iint_R f(x, y) \, dA = \iint_S f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv$$

### 1.2.1 Triple Integrals

Lets use the definition of the Jacobian, extend it to three dimensions and find the formula for a triple integral and use it to derive the formula for spherical coordinates.

The Jacobian of  $T$  is the following  $3 \times 3$  determinant:

$$\frac{\partial(x, y, z)}{\partial(u, v, w)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix}$$

this gives us the MASSIVE formula lol:

$$\iiint_R f(x, y, z) \, dV = \iiint_S f(x(u, v, w), y(u, v, w), z(u, v, w)) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| \, du \, dv \, dw$$

now lets use this to find the formula for triple integration in spherical coordinates!!!! (i'm losing my fucking mind)

$$x = \rho \sin \phi \cos \theta \quad y = \rho \sin \phi \sin \theta \quad z = \rho \cos \phi$$

lets compute this absolute unit of a jacobian

$$\begin{aligned} \frac{\partial(x, y, z)}{\partial(\rho, \theta, \phi)} &= \begin{vmatrix} \sin \phi \cos \theta & -\rho \sin \phi \sin \theta & \rho \cos \phi \cos \theta \\ \sin \phi \sin \theta & \rho \sin \phi \cos \theta & \rho \cos \phi \sin \theta \\ \cos \phi & 0 & -\rho \sin \phi \end{vmatrix} \\ &= \cos \phi (-\rho^2 \sin \phi \cos \phi \sin^2 \theta - \rho^2 \sin \phi \cos \phi \cos^2 \theta) - \rho \sin \phi (\rho \sin^2 \phi \cos^2 \theta + \rho \sin^2 \phi \sin^2 \theta) \\ &= (\text{This reduces all the way to}) \rho^2 \sin \phi \quad (\text{lol}) \end{aligned}$$

anyways, putting this back into our equation would give us

$$\iiint_R f(x, y, z) \, dV = \iiint_S f(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \rho^2 \sin \phi \, d\rho \, d\theta \, d\phi$$

lets goo!!!!!!

## 2 Vector Fields

In general, a vector field is a function whose domain is a set of points in  $\mathbb{R}^2$  (or  $\mathbb{R}^3$  in three dimensions) is a function  $\mathbf{F}$  that assigns to each point  $(x, y)$  in  $D$  a two-dimensional vector  $\mathbf{F}(x, y)$ .

The best way to picture a vector field is to draw the arrow representing the vector  $\mathbf{F}(x, y)$  starting at the point  $(x, y)$ . Of course it's impossible to do this for all points  $(x, y)$ , but we can gain a reasonable impression of  $\mathbf{F}$  by doing it for a few representative points in  $D$ . since  $\mathbf{F}(x, y)$  is a two-dimensional vector, we can write it in terms of its **component functions**  $P$  and  $Q$  as follows:

$$\mathbf{F}(x, y) = P(x, y)\hat{i} + Q(x, y)\hat{j} = \langle P(x, y), Q(x, y) \rangle$$

or, for short,  $\mathbf{F} = P \mathbf{i} + Q \mathbf{j}$

Notice that  $P$  and  $Q$  are scalar functions of two variables and are sometimes called **scalar fields** to distinguish them from vector fields.

Let  $E$  be a subset of  $\mathbb{R}^3$ . a **vector field on  $\mathbb{R}^3$**  is a function  $\mathbf{F}$  that assigns each point  $(x, y, z)$  in  $E$  a three-dimensional vector  $\mathbf{F}(x, y, z)$ .

### 2.1 Gradient Fields

if  $f$  is a scalar function of two variables, recall that  $\nabla f$  is defined by  $\nabla f(x, y) = f_x(x, y)\mathbf{i} + f_y(x, y)\mathbf{j}$

Therefore,  $\nabla f$  is really a vector field on  $\mathbb{R}^2$  and is called a **gradient vector field**. Likewise, if  $f$  is a scalar function of three variables (it extends but im too lazy to type this out).

The length of the gradient vector is the value of the directional derivative of  $f$  and closely spaced level curves indicate a steep graph.

A vector field  $\mathbf{F}$  is called a **conservative vector field** if it is the gradient of some scalar function, that is, if there exists a function  $f$  such that  $\mathbf{F} = \nabla f$ . in this situation  $f$  is called a **potential function** for  $\mathbf{F}$ .

Not all vector fields are conservative though!!

### 3 Line Integrals

## 4 The Fundamental Theorem of Line Integrals

## 5 Green's Theorem



## 6 Curl and Divergence

## 7 Parametric Surfaces

## 8 Surface Integrals