

**11<sup>th</sup> IMACSSS International Conference**  
5<sup>th</sup> World Scientific Congress of  
Combat Sports and Martial Arts  
October 13–15, 2022, Rzeszów, Poland

# ABSTRACT BOOK



Wojciech Jan Cynarski, Przemysław Pawelec,  
Paweł Świder and Jerzy Kulasa [Eds.]

Rzeszów 2022

### **Scientific Committee**

Prof. Dr hab. Wojciech J. Cynarski – President \* Prof. Dr Maicon Albuquerque (Brazil)  
Prof. Dr hab. Tadeusz Ambroży (Poland) \* Prof. Dr Pavol Bartik (Slovakia)  
Prof. Dr hab. Zbigniew Borysiuk (Poland) \* Dr Michel Calmet (France)  
Prof. Dr Raquel Escobar Molina (Spain) \* Prof. Dr Abel A. Figueiredo (Portugal)  
Prof. Dr Thomas A. Green (USA) \* Prof. Dr Zhang Guodong (China)  
Prof. Dr Carlos Gutiérrez-García (Spain) \* Prof. Dr Kiyoshi Ito (Japan)  
Prof. Dr Rudolf Jakhel (Slovenia) \* Dr John A. Johnson (USA)  
Dr Mohsen Kazemi (Canada) \* Prof. Dr Georgiy Korobeynikov (Ukraine)  
Dr James Lee-Barron (UK) \* Doc. Dr Irena Martinkova (Czech Republic)  
Prof. Dr Bianca Miarka (Brazil) \* Dr Udo Moenig (Germany)  
Prof. Dr Robert J. Parkes (Australia) \* Dr Mikel Pérez-Gutiérrez (Spain)  
Prof. Dr Leonid Podrigalo (Ukraine) \* Dr Sergio Raimondo (Italy)  
Doc. Dr Zdenko Reguli (Czech Republic) \* Doc. Dr Pavel Ružbarský (Slovakia)  
Dr Mohamad Nizam Mohamed Shapie (Malaysia) \* Prof. Dr Jozef Simenko (UK)  
Dr Kentaro Tai (Japan) \* Prof. Dr Antonio Vences de Brito (Portugal)  
Doc. Dr Michal Vit (Czech Republic) \* Dr hab. Jacek Wąsik (Poland)  
Prof. Dr hab. Kurt Weis (Germany) \* Dr Jong-Hoon Yu (USA) \* Dr Howard Zeng (USA)

\* \* \*

### **Congress Organisers**

University of Rzeszów \* Institute of Physical Culture Studies, UR  
\* Chair of Social and Humanistic Foundations of Physical Culture  
\* **IMACSSS – International Martial Arts and Combat Sport Scientific Society**  
\* IPA - Idokan Poland Association, Committee of Scientific Research

\* \* \*

### **In cooperation with**

\* EMAC – European Martial Arts Committee  
\* IMAS – the Institute of Martial Arts and Sciences – Lancashire, UK  
\* JAB – Japanese Academy of Budo – Tokyo

\* \* \*

### **Sponsor**

\* President of the City of Rzeszów

\* \* \*

### **Under the patronage of**

\* President of the City of Rzeszów \* Rector of the University of Rzeszów  
\* “Ido Movement for Culture. Journal of Martial Arts Anthropology”

\* \* \*

Proofreader: dr Renata Grzywacz

**Dariusz Mosler**<sup>1(ABCDEF)</sup>, **Vladena Pasko**<sup>2(BCDE)</sup>, **Jacek Wąsik**<sup>1(A,DEFG)</sup>

<sup>1</sup> Jan Długosz University in Czeŝochowa, Department of Kinesiology and Health Prevention (Poland)

<sup>2</sup> Kharkiv State Academy of Physical Culture, Kharkiv (Ukraine)

A-Study Design, B-Data Collection, C-Statistical Analysis, D-Data Interpretation, E-Manuscript Preparation, F-Literature Search

## COMPARISON OF ELECTROMYOGRAPHIC ACTIVITY OF RECTUS FEMORIS BETWEEN ISOKINETIC KNEE EXTENSION TRAINING AND ROUNDHOUSE KICKS - CASE STUDY

**Key words:** biomechanics; martial arts; strength training

**Introduction.** Roundhouse kick is one of the most commonly used leg technique in many martial arts [Gavagan, Sayers 2017]. Its main power is generated by knee extension at final stage of technique execution [Thibordee, Prasartwuth 2014]. Proper power training needs to imitate electromyographic activity of muscles generated during the roundhouse kick execution to reduce need of motor learning transfer [Moreira *et al.* 2021]. The aim is to analyse the contribution of the data collected with BIODEX isokinetic device in better understanding the execution of the roundhouse kick.

**Methods.** 33 years old karateka with 13 years of experience performed 10 roundhouse kicks to a shield and 10 repetition of knee extension in isokinetic test protocol on BIODEX SYSTEM 4 PRO with maximum speed of 500 deg/s, both with right leg. EMG electrode was attached to rectus femoris accordingly to SENIAM protocol. NORAXON Ultium IMU sensors was used to measure EMG activity of right rectus femoris muscle, angular velocity of knee extension movement and foot acceleration.

**Results.** For both roundhouse kick and isokinetic knee extension, occurrence of maximum angular velocity was before the occurrence of maximum acceleration with mean time difference was 0.01s and 0.034s, respectively. In both cases peak activation of rectus femoris occurs before reaching maximum angular velocity and acceleration ( $x = 0.138s$  and  $x = 0.064s$  respectively) (Table 1).

**Table 1.** Differences in the time of occurrence for maximal values of studied variables.

	Roundhouse kick				Isokinetic knee extension			
Move	$T_{EMGmax} - T_{amax}$	$T_{EMGmax} - T_{\omega max}$	$T_{\omega max} - T_{amax}$	Peak muscle activity (uV)	$T_{EMGmax} - t_a$	$T_{EMGmax} - T_{\omega max}$	$T_{\omega max} - T_{amax}$	Peak muscle activity (uV)
1	-0.325	-0.123	-0.202	701.89	-0.096	-0.059	-0.037	317.37
2	-0.102	-0.115	0.013	614.92	0.013	0.052	-0.039	455.48
3	-0.125	-0.137	0.012	665.23	-0.183	-0.148	-0.035	508.05
4	-0.112	-0.123	0.011	499.10	-0.053	-0.014	-0.039	610.38
5	-0.111	-0.122	0.011	684.36	-0.022	0.014	-0.036	616.34
6	-0.114	-0.124	0.010	641.31	-0.042	-0.009	-0.033	578.00

7	-0.113	-0.123	0.010	678.01	-0.100	-0.069	-0.030	704.51
8	-0.147	-0.161	0.014	549.84	-0.022	0.010	-0.032	876.78
9	-0.122	-0.126	0.004	692.72	-0.092	-0.063	-0.030	538.48
10	-0.111	-0.126	0.015	719.44	-0.042	-0.008	-0.034	724.84
<b>mean</b>	<b>-0.138</b>	<b>-0.128</b>	<b>-0.010</b>	<b>644.68</b>	<b>-0.064</b>	<b>-0.029</b>	<b>-0.034</b>	<b>593.02</b>

$T_{EMGmax}$  – time of reaching maximal electromyographic activity;  $T_{amax}$  – time of reaching maximal acceleration;  $T_{\omega max}$  – time of reaching maximal angular velocity

For roundhouse kick, mean maximal electromyographic activity of rectus femoris was  $644.68 \text{ uV} \pm 67.35$ . For isokinetic knee extension mean maximal electromyographic activity of rectus femoris was  $593.02 \text{ uV} \pm 147.13 \text{ uV}$  (Table 1).

**Discussion and conclusion.** Angular velocity of knee extension during roundhouse kick (over 2000 degs) exceeds the possibility of isokinetic machine adjustment, making impossible to imitate the speed of fast kicking techniques on BIODEX (fixed on 500 deg/s). Contrary to previous studies, no muscle fatigue was observed as peak muscle activity varies without decreasing tendency through this experiment [Quinzi *et al.* 2016]. Similar level of rectus femoris activity during the execution of both moves in 5 trials suggest that trained martial artist can maintain muscle control of knee extension in different circumstances, even if physical target of a kick is not present [Kim *et al.* 2011]. Shorter time between maximum muscle activity and maximum movement speed justifies use of isolated isokinetic training for shaping power of roundhouse kick, which is concurrent with other study, where frontal kick was tested [Quinzi *et al.* 2013]. As karate type of roundhouse kick have faster onset of peak angular velocities than Taekwondo athletes [Barnamehei *et al.* 2020], but lower maximal velocity of a kicks [Gavagan, Sayers 2017], isokinetic training could be beneficial as training transfer tool in changing motor pattern of technique execution, depending on training goals. High speed isokinetic training could be beneficial for rehabilitation after lower limb injuries to regain similar muscle control before going back to martial arts practice.

## References

1. Barnamehei H., Khazaei F., Safaei M., Jabari H., Golfeshan N., Barnamehei M., Rezaei A., Kharazi M., Naghavi N. (2020), *Motor learning and training strategy effect on motor control; Comparison between Taekwondo and Karate front kick (Ap Chagi and Mae Geri)*, "International Journal of Martial Arts", vol. 6, pp. 48-65.
2. Gavagan C.J., Sayers M.G.L. (2017), *A biomechanical analysis of the roundhouse kicking technique of expert practitioners: A comparison between the martial arts disciplines of Muay Thai, Karate, and Taekwondo*, "PLOS ONE", vol. 12, no. 8, pp. 1-15; doi: 10.1371/journal.pone.0182645.
3. Kim Y.K., Kim Y.H., Im S.J. (2011), *Inter-joint coordination in producing kicking velocity of Taekwondo kicks*, "Journal of Sports Science and Medicine", vol. 10, no. 1, pp. 31-38.
4. Moreira P.V.S., Falco C., Menegaldo L.L., Goethel M.F., de Paula L.V., Gonçalves M. (2021), *Are isokinetic leg torques and kick velocity reliable predictors of competitive level in taekwondo athletes?*, "PLOS ONE", vol. 16, no. 6, pp. e0235582; doi: 10.1371/journal.pone.0235582.
5. Quinzi F., Camomilla V., Felici F., Di Mario A., Sbriccoli P. (2013), *Differences in neuromuscular control between impact and no impact roundhouse kick in athletes of*

*different skill levels*, "Journal of Electromyography and Kinesiology", vol. 23, no. 1, pp. 140-150; doi: 10.1016/j.jelekin.2012.09.006.

6. Quinzi F., Camomilla V., Di Mario A., Sbriccoli P. (2016), *Repeated kicking actions in karate: Effect on technical execution in elite practitioners*, "International Journal of Sports Physiology and Performance", vol. 11, no. 3, pp. 363-369; doi: 10.1123/ijsp.2015-0162.

7. Thibordee S., Prasartwuth O. (2014), *Effectiveness of roundhouse kick in elite Taekwondo athletes*, "Journal of Electromyography and Kinesiology", vol. 24, no. 3, pp. 353-358; doi: 10.1016/j.jelekin.2014.02.002.