

Part III - Algebraic Geometry

Lectured by Dhruv Ranganathan

Artur Avameri

Contents

0	Introduction	2
1	Beyond algebraic varieties	2
1.1	Summary of classical algebraic geometry	2
1.2	Limitations of classical algebraic geometry	4
1.3	The spectrum of a ring	4
1.4	A topology on $\text{Spec } A$	5
1.5	Functions on opens	5

0 Introduction

6 Oct 2022,
Lecture 1

The course consists of four parts.

- (1) Basics of sheaves on topological spaces.
- (2) Definition of schemes and morphisms.
- (3) Properties of schemes (e.g. the algebraic geometry notion of compactness and other properties).
- (4) A rapid introduction to the cohomology of schemes.

The main reference for the course is Hartshorne's *Algebraic Geometry*.

1 Beyond algebraic varieties

08 Oct 2022,
Lecture 2

1.1 Summary of classical algebraic geometry

We let $k = \bar{k}$ be an algebraically closed field and consider $\mathbb{A}_k^n = \mathbb{A}^n = k^n$ as a set.

Definition 1.1. An **affine variety** is a subset $V \subset \mathbb{A}^n$ of the form $\mathbb{V}(S)$ with $S \subset k[x_1, \dots, x_n]$, where \mathbb{V} is the common vanishing locus.

Note that $\mathbb{V}(S) = \mathbb{V}(I(S))$ (the ideal generated by S). By Hilbert Basis Theorem (since $k[x_1, \dots, x_n]$ is noetherian), $\mathbb{V}(I(S)) = \mathbb{V}(S')$ for some finite set $S' \subset k[x_1, \dots, x_n]$.

In fact, $\mathbb{V}(I) = \mathbb{V}(\sqrt{I})$, where

$$\sqrt{I} = \{f \in k[x_1, \dots, x_n] \mid f^m \in I \text{ for some } m \geq 0\}$$

is the **radical** of I . For example, in $k[x]$, if $I = (x^2)$, then $\sqrt{I} = (x)$.

Definition 1.2. Given varieties $V \subset \mathbb{A}^n$ and $W \subset \mathbb{A}^m$, a **morphism** is a (set-theoretic) map $\phi : V \rightarrow W \subset \mathbb{A}_k^m$ such that if $\phi = (f_1, \dots, f_m)$, then each f_i is the restriction of a polynomial in $\{x_1, \dots, x_n\}$.

An **isomorphism** is a morphism with a two-sided inverse.

Our basic correspondence is

$$\begin{array}{c} \{\text{Affine varieties over } k\} / \text{up to isomorphism} \\ \leftrightarrow \\ \{\text{finitely generated } k\text{-algebras } A \text{ without nilpotent elements}\} \end{array}$$

A finitely generated k -algebra is just a quotient of a polynomial ring in finitely many variables. A nilpotent element is such that some power of it is zero. For example, in $k[x]/(x^2)$, the element x is nilpotent.

How does this correspondence work? Given a variety V (representing an isomorphism class), we write $V = \mathbb{V}(I)$ for $I \subset k[x_1, \dots, x_n]$ a radical ideal¹, and map $V \mapsto k[x_1, \dots, x_n]/I$.

For the reverse, if A is a finitely generated nilpotent free algebra, then $A \cong k[y_1, \dots, y_m]/J$ where we can choose J to be radical (exercise: why?).

We have to check that this is independent of our choice on both sides (exercise: think through this, it should be clear).

Definition 1.3. The algebra associated to V is classically denoted $k[V]$ and called the **coordinate ring of V** .

We have the compatibility of morphisms with our basic correspondence: there is a bijection between

$$\text{Morphisms}(V, W) \leftrightarrow \text{Ring homomorphisms}_k(k[W], k[V])$$

(here RingHom_k means that our homomorphisms preserve k).

We can now make our set into a topological space:

Definition 1.4. Let $V = \mathbb{V}(I) \subset \mathbb{A}^n$ be a variety with coordinate ring $k[V]$. The **Zariski topology** on V is defined such that the closed sets are $\mathbb{V}(S)$, where $S \subset k[V]$.

If $V \cong W$, then the Zariski topological spaces are homeomorphic as varieties (exercise).

Theorem 1.1 (Nullstellensatz). Fix V a variety and let $k[V]$ be its coordinate ring. Given $p \in V$, we can produce a homomorphism $\text{ev}_p : k[V] \rightarrow k$ by sending $f \mapsto f(p)$. Note that ev_p is surjective (since we have constant functions), hence $\ker(\text{ev}_p) = m_p$ is a maximal ideal, giving us a map

$$\{\text{points of } V\} \rightarrow \{\text{maximal ideals in } k[V]\}.$$

Nullstellensatz says that this is actually a bijection. For the converse map, given $m \subset k[V]$, we get a quotient $k[V] \rightarrow k[V]/m = k$ (Nullstellensatz says this extension is finite, hence must be k). So using/choosing a representation for V in $k[x_1, \dots, x_n]$ gives a surjective homomorphism onto k and specifies a bunch of points.

¹A radical ideal is an ideal equal to its radical.

1.2 Limitations of classical algebraic geometry

Question. What is an abstract variety, i.e. "some "space" X such that locally as a cover $\{U_i\}$, each U_i is an affine variety, compatible with overlaps".

Example 1.1 (non-algebraically closed fields). Take $I = (x^2 + y^2 + 1) \subset \mathbb{R}[x, y]$. Then $\mathbb{V}(I) = \emptyset \subset \mathbb{R}^2$, but I is prime, so radical, so nullstellensatz fails.

Question. On what topological space is $\mathbb{R}[x, y]/(x^2 + y^2 + 1)$ "naturally" the set of functions? (or \mathbb{Z} , or $\mathbb{Z}[x]$).

Example 1.2 (Why restrict to radical ideals?). Take $C = \mathbb{V}(y - x^2) \subset \mathbb{A}_k^2$ and $D = \mathbb{V}(x, y)$, so $C \cap D = \mathbb{V}(y, y - x^2) = \mathbb{V}(x, y) = \{(0, 0)\}$. This is a single point, but if $D_\delta = \mathbb{V}(y + \delta)$ for some $\delta \in k$, then $C \cap D_\delta = \{\pm\sqrt{\delta}\}$, which is 2 points for all $\delta \neq 0$. In other words, intersections of varieties don't want to be varieties.

1.3 The spectrum of a ring

Let A be a commutative ring with identity. We will define a topological space on which A is the ring of functions.

11 Oct 2022,
Lecture 3

Definition 1.5. The **Zariski spectrum** of A is

$$\text{Spec } A = \{\mathfrak{p} \subset A \mid \mathfrak{p} \text{ is a prime ideal}\}.$$

A ring homomorphism $\phi : A \rightarrow B$ induces a map $\phi^{-1} : \text{Spec } B \rightarrow \text{Spec } A$ by $q \mapsto \phi^{-1}(q)$. In general, the preimage of a prime ideal is a prime ideal.

Warning. This would fail if we only considered maximal ideals, since the preimage of a maximal ideal need not be maximal.

Given $f \in A$ and $\mathfrak{p} \in \text{Spec}(A)$, we have an induced $\bar{f} \in A/\mathfrak{p}$ obtained via a quotient. Informally, we can evaluate any $f \in A$ at points $\mathfrak{p} \in \text{Spec}(A)$ with the caveat that the codomain of this evaluation depends on \mathfrak{p} .

Example 1.3. Take $A = \mathbb{Z}$. Then $\text{Spec } A = \text{Spec } (\mathbb{Z}) = \{p \mid p \text{ is prime}\} \cup \{(0)\}$. Let's pick an element in \mathbb{Z} , say $132 \in \mathbb{Z}$. Given a prime p , we can look at $132 \pmod{p} \in \mathbb{Z}/p$. The takeaway here is that

$$\begin{aligned} \text{Spec } \mathbb{Z} &\rightarrow \text{Space} \\ 132 \in \mathbb{Z} &\rightarrow \text{a function} \\ 132 \pmod{p} &\rightarrow \text{value of that function at } p. \end{aligned}$$

Note that based on the value of p , our codomain changes from point to point.

Example 1.4. Take $A = \mathbb{R}[x]$, then $\text{Spec } \mathbb{R}[x] = \mathbb{C}/\text{complex conjugation} \cup \{(0)\}$.

Exercise. Draw $\text{Spec } \mathbb{Z}[x]$ and $\text{Spec } k[x]$ for k any field (i.e. describe all prime ideals and their containment). This is on example sheet 1.

Example 1.5. If $A = \mathbb{C}[x]$, then $\text{Spec } A = \mathbb{C} \cup \{(0)\}$, where given $a \in \mathbb{C}$, we send it to the maximal ideal $\langle x - a \rangle$.

1.4 A topology on Spec A

Fix $f \in A$. Then $\mathbb{V}(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \equiv 0 \pmod{\mathfrak{p}}\} \subset \text{Spec } A$. (Note that $f \equiv 0 \pmod{\mathfrak{p}}$ is the same as $f \in \mathfrak{p}$).

Similarly for $J \subset A$ an ideal, $\mathbb{V}(J) = \{\mathfrak{p} \in \text{Spec } A \mid f \in \mathfrak{p} \forall f \in J\}$.

Proposition 1.2. The sets $\mathbb{V}(J) \subset \text{Spec } A$ ranging over all ideals J form the closed sets of a topology on $\text{Spec } A$. This topology is called the **Zariski topology**.

Proof. Easy fact: \emptyset and $\text{Spec } A$ are closed, since we have functions 1 (vanishing nowhere) and 0 (vanishing everywhere). Since $\mathbb{V}(\sum_{\alpha} I_{\alpha}) = \bigcap_{\alpha} \mathbb{V}(I_{\alpha})$ (this is because $I_1 + I_2$ is the smallest ideal containing $I_1 \cup I_2$), arbitrary intersections are closed.

Finally, we claim $\mathbb{V}(I_1) \cup \mathbb{V}(I_2) = \mathbb{V}(I_1 \cap I_2)$. The containment \subset is clear: if a prime ideal contains I_1 or I_2 , it contains $I_1 \cap I_2$. Conversely, $I_1 I_2 \subset I_1 \cap I_2$, so if $I_1 I_2 \subset I_1 \cap I_2 \subset \mathfrak{p}$, then by primality $I_1 \subset \mathfrak{p}$ or $I_2 \subset \mathfrak{p}$. \square

Example 1.6. Let $k = \mathbb{C}$ and consider $\text{Spec } \mathbb{C}[x, y]$. We make a few observations:

- The point $(0) \in \text{Spec } \mathbb{C}[x, y]$ is dense in the Zariski topology, i.e. $\overline{\{(0)\}} = \text{Spec } \mathbb{C}[x, y]$ because every prime ideal contains (0) (because we are in an integral domain).
- Consider the prime ideal $(y^2 - x^3)$ (which is prime since the quotient is an integral domain). Consider a maximal ideal $\mathfrak{m}_{a,b} = (x - a, y - b)$. We can ask: when is $\mathfrak{m}_{a,b} \in \overline{\{(y^2 - x^3)\}}$? The answer: if and only if $b^2 = a^3$, e.g. $(1, 1)$ (see example sheet 1). The lesson here is that points are not closed in the Zariski topology.

1.5 Functions on opens

Definition 1.6. Let $f \in A$. Define the **distinguished open** corresponding to f to be

$$\mathcal{U}_f = (\text{Spec}(A)) / \mathbb{V}(f).$$

Example 1.7. • Let $A = \mathbb{C}[x]$, so $\text{Spec } A = \mathbb{C} \cup \{(0)\}$ (with the Zariski topology). Take $f = x$ and consider \mathcal{U}_x . Recall the bijection $\text{Spec } \mathbb{C} \leftrightarrow \mathbb{C} \cup \{(0)\}$ by $(x - a) \leftrightarrow a \in \mathbb{C}$ and $(0) \leftrightarrow (0)$. Then $\mathbb{V}(x) = \{\mathfrak{p} \in \text{Spec } A \mid x \in \mathfrak{p}\} = \{(x)\}$, so $\mathcal{U}_f = \text{Spec } A \setminus \{(x)\}$.

- More generally, suppose we fix $a_1, \dots, a_r \in \mathbb{C}$, then $\text{Spec } A \setminus \{(x-a_i)\}_{i=1}^r = \mathcal{U}$ and $\mathcal{U} = \mathcal{U}_f$, where $f = \prod_{i=1}^r (x - a_i)$.

Lemma 1.3. The distinguished opens \mathcal{U}_f taken over all $f \in A$ form a basis for the Zariski topology on $\text{Spec } A$.

Proof. Left as an exercise on example sheet 1. □

A bit of commutative algebra:

Definition 1.7. Given $f \in A$, the **localization of A at f** is $A_f = A[x]/(xf-1)$, which we can informally think of as $A_f = A[\frac{1}{f}]$.

Lemma 1.4. The distinguished open $\mathcal{U}_f \subset \text{Spec } A$ is naturally homeomorphic to $\text{Spec } A_f$.