

# Mechanism of catalysis by retaining $\beta$ -glycosyl hydrolases

André White\* and David R Rose†

Recent structural studies provide a fresh look at the catalytic mechanism of polysaccharide hydrolysis by retaining  $\beta$ -glycosyl hydrolases. Highlights include insights into saccharide ring distortion, both upon binding and during the course of catalysis, and evidence for the regulation of the  $pK_a$  of key catalytic residues.

## Addresses

\*Rosenstiel Basic Medical Sciences Research Center, Brandeis University, 315 South Street, Waltham, MA 02254-9110, USA; e-mail: [awhite@auriga.rose.brandeis.edu](mailto:awhite@auriga.rose.brandeis.edu)

†Ontario Cancer Institute, and Department of Medical Biophysics, University of Toronto, 610 University Avenue, Toronto, Ontario, Canada, M5G 2M9; e-mail: [drose@oci.utoronto.ca](mailto:drose@oci.utoronto.ca)

Current Opinion in Structural Biology 1997, 7:645–651

<http://biomednet.com/elecref/0959440X00700645>

© Current Biology Ltd ISSN 0959-440X

## Abbreviations

**Gal** galactose  
**GalNAc** *N*-acetyl-galactosamine  
**GlcNAc** *N*-acetyl-glucosamine  
**MurNAc** *N*-acetyl-muramic acid

## Introduction

Retaining  $\beta$ -glycosyl hydrolases release their saccharide products with retention of configuration at the newly exposed anomeric center [1]. Although these enzymes may adopt a wide variety of folding motifs, their catalytic function is similar [2]. The catalytic mechanism of these enzymes was proposed some years ago to involve either a covalent glycosyl–enzyme intermediate [3] or an oxocarbenium-ion-like intermediate [4]. Evidence suggests that both states are involved, such that a covalent glycosyl–enzyme intermediate is formed and then released via oxocarbenium-ion-like transition states [5]. The catalytic nucleophile is a carboxylate sidechain, except for one example of a histidine residue [6]. Typically, a second carboxylate sidechain provides acid/base catalytic assistance, first by donating a proton to the linking oxygen of the scissile bond, and then by abstracting a proton from an incoming water molecule; this water molecule reacts with the transition-state species to form the product. Substrate cleavage is defined to occur between binding subsites –1 and +1 [7].

Recent studies from X-ray crystallography and other techniques provide structural insights into events in the catalytic mechanism of retaining  $\beta$ -glycosyl hydrolases. We review results that suggest that, during catalysis, the saccharide unit binding at subsite –1 adopts a ‘twisted-boat’ conformation, and deforms to a ‘half-chair’ at the transition state and then to a ‘chair’ at the catalytic

intermediate. The final steps of the mechanism involve a reversal of this series of conformations.

## Saccharide conformation on binding

Catalysis by retaining  $\beta$ -glycosyl hydrolases is believed to involve saccharide deformation at the enzyme’s catalytic center. For lysozyme, a ‘half-chair’ distortion was originally proposed on the basis of early crystallographic results [4,8,9]. Recent crystallographic analyses have provided evidence of other saccharide conformations during binding and formation of the catalytic intermediate in this mechanism.

Two high-resolution crystallographic studies of hen egg-white lysozyme reveal saccharide distortion towards a ‘half-chair’ form, which approaches the transition-state conformation. In one study, the reaction product MurNAc-GlcNAc-MurNAc binds to the active site of lysozyme [10], whereas in a second study oligosaccharides of GlcNAc are cleaved by a mutant lysozyme [11]. In this latter report, careful refinement has revealed an inverted  $\alpha$ -saccharide product. These results support the ‘half-chair’ model, although in both examples the saccharide ligand does not span the scissile linkage. Additionally, crystallographic results of the Thr26→Glu mutant of T4 lysozyme with a covalent substrate [12] suggest a ‘half-chair’ conformation for the catalytic intermediate. Because of its inherent non-natural modification, however, this system must be interpreted with care in the analysis of the natural catalytic intermediate.

Alternatively, the saccharide at the –1 site in retaining  $\beta$ -glycosyl hydrolases has been proposed to initially adopt a boat conformation on the basis of stereochemical considerations [13], although this argument is not yet generally accepted [5]. Until recently, the available experimental evidence could neither support nor exclude the possibility of a boat conformation.

There is a precedent for induced saccharide distortion on binding to proteins. For example, an unusual boat form has been observed for sialic acid (*N*-acetyl-neuraminic acid) in the influenza virus neuraminidase [14]. The carbohydrate distortion may be attributed to the carboxylic substituent at C1, which forms stabilizing electrostatic interactions with a triarginyl cluster (residues 118, 292, and 371) and advantageous packing with Tyr106. More recently, the 2.1 Å resolution crystal structure of the heat-labile enterotoxin from *Escherichia coli* complexed to the tumour marker disaccharide D-Gal- $\beta$ -1,3-GalNAc reveals one sugar unit as a chair and the other as a boat conformation [15]. The geometric rearrangement of the *N*-acetylglactosyl residue allows the attached T-antigen

protein moiety to be directed toward the outside of the binding pocket of this bacterial toxin.

Recent crystallographic evidence from a number of enzymes suggests that the saccharide at position -1 may indeed be distorted upon binding to retaining  $\beta$ -glycosyl hydrolases. Crystallographic analysis of the endocellulase E1 with cellotetraose suggests that the saccharide may not be bound in a relaxed chair form at subsite -1 [16<sup>••</sup>], although the nature of this deformation has not been identified. From the high-resolution crystal structure of the endoglucanase CelA from *Clostridium thermocellum* complexed to oligosaccharides, it is inferred that the saccharide unit at subsite -1 (site 'C') must undergo deformation to connect to the observed saccharide units [17<sup>•</sup>].

An abortive complex of *Serratia marcescens* chitobiase with di-*N*-acetyl-glucosamine (di-GlcNAc) has been obtained by a short ten minute incubation with its substrate [18<sup>••</sup>]. In addition to suggesting substrate assistance in catalysis by the substituent at C2 of the ligand, this crystal structure, refined to 1.9 Å resolution, shows a nonhydrolyzed disaccharide spanning the scissile bond, in which the sugar unit at position -1 forms a twisted-boat conformation. Interestingly, the presumed mechanism for this chitobiase is similar to the reverse of that proposed for the transglycosylase SLT70 [19], even though these two enzymes have different enzymatic products. A twisted-boat conformation is also observed in the crystal structure of the retaining  $\beta$ -glycosyl hydrolase EGI in complex with a thio-substituted nonhydrolyzable trisaccharide inhibitor [20<sup>••</sup>]. Although the resolution of this crystal structure is only 2.7 Å, the saccharide unit at position -1 is clearly constrained to adopt a twisted-boat form. The consequence of this deformation for the catalytic mechanism is that it positions the glycosidic oxygen in the proximity of the acid/base catalyst, alleviates steric hindrance between C1 and the catalytic nucleophile, and presents the aglycon leaving group in an axial position [21]. This latter aspect may have favourable stereoelectronic implications as previously predicted [13].

In contrast to  $\beta$ -glucose, the linking oxygen of  $\alpha$ -glucose saccharides is inherently positioned axial in the relaxed sugar form in solution. A saccharide close to a chair conformation is observed for the nonhydrolyzable D-glucodihydroacarbose and acarbose inhibitors in a glucoamylase [22<sup>•</sup>]. This conformation is consistent with stereoelectronic implications for an elimination reaction [13] without requiring a deformation towards a twisted-boat conformation.

### Acid/base catalyst

For a double-displacement catalytic mechanism, a side-chain provides acid/base catalytic assistance. Typically, the sidechain of this acid/base catalyst is a carboxylate located ~5.5 Å away from the catalytic nucleophile for retaining enzymes and ~9.5 Å for inverting enzymes [1,23].

In endoglucanase CelC from *Clostridium thermocellum*, the acid/base catalyst is located in a loop that is believed to fold back on the active site upon binding of the substrate [24]. This example indicates an induced-fit activation mechanism.

Because of the dual role for a general acid/base catalyst, progress through catalysis requires modulation of the ionization state of its sidechain. For example, kinetic studies with *Agrobacterium faecalis*  $\beta$ -glucosidase show that the  $pK_a$  of a key active-site residue changes its value by nearly three units of pH during catalysis [25]. While  $pK_a$  is a useful indicator of the ionization state of amino acids, its measurement for key catalytic residues is restricted by difficulties in distinguishing specific titration values within a protein. One means of addressing this problem is the use of 2D NMR experiments in titrating specific sidechains. In an elegant study combining 2-deoxy-2-fluoro-saccharides and <sup>13</sup>C-NMR spectroscopy on a 20 kDa xylanase from *Bacillus circulans*, the group of McIntosh [26<sup>••</sup>,27] have monitored chemical shifts of specific enzyme residue sidechains that are pH dependent. Their study reveals  $pK_a$  values of 4.6 for the catalytic nucleophile Glu78, and 6.7 for Glu172. Interestingly, the  $pK_a$  of Glu172 decreases to 4.2 in a covalent glycosyl intermediate with Glu78, consistent with a dual role of Glu172 as an acid/base catalyst. When the charge on Glu78 is abolished by forming a Gln78 mutant, a lower  $pK_a$  of 4.2 for Glu172 is observed, even in the absence of substrate. This suggests that the ionization state of the catalytic nucleophile (Glu78 for this xylanase) modulates the  $pK_a$  value of the acid/base catalyst. Alternatively, a close approach of the two key catalytic residues may explain the large shift in  $pK_a$  for the acid/base catalyst during catalysis, as proposed on the basis of the crystal structure of endocellulase E1 [16<sup>••</sup>]. Such a large decrease in the  $pK_a$  value of Glu172 is likely to be common for retaining  $\beta$ -glycosyl hydrolases.

### Catalytic glycosyl-enzyme intermediate

Evidence from retaining  $\beta$ -glycosyl hydrolases other than lysozyme has generally favoured a covalent intermediate for this mechanism [5].  $\alpha$ -secondary deuterium kinetic isotope effects have been measured to be greater than 1.0 for two catalytic steps [5,28], which is indicative of a tetrahedral geometry at the saccharide C1 in the intermediate. In addition, Brønsted plot analyses, using a variety of substrates with different  $pK_a$  values, indicate a large degree of bond cleavage at the transition state, consistent with a covalent intermediate [25,28,29,30<sup>•</sup>]. Finally, such a covalent intermediate has been trapped by the use of 2-fluoro-glycosides with relatively reactive leaving groups [31–33].

Typically, the sidechain of either an aspartate or a glutamate fulfils the role of the catalytic nucleophile in retaining  $\beta$ -glycosyl hydrolases [5]. A comparison of many similar enzymes, even those with low amino acid sequence identity, may help to identify key catalytic residues, such

as for hevamine—a plant chitinase/lysozyme [34]. Frequently, more than one carboxylate sidechain is conserved for homologous  $\beta$ -glycosyl hydrolases; therefore, amino acid sequence analyses alone are not sufficient to identify the catalytic nucleophile. Recently, a direct strategy for identifying the catalytic nucleophile has been proposed [30\*] on the basis of the formation of azide–glycoside products by mutated enzymes. In a different report, the combination of kinetic and crystallographic studies for cellobiohydrolase I from *Trichoderma reesei* and its complex with saccharides provides a useful setting to assess the importance of a catalytic trio of carboxylates [35].

In an attempt to locate the key active-site catalytic nucleophiles of retaining  $\beta$ -glycosyl hydrolases and their specificity of interaction, saccharide-derived epoxide compounds have often been used as labelling agents. Such epoxide compounds contain two parts: a reactive epoxide group, which forms an irreversible covalent attachment to a nucleophilic sidechain of the targeted enzyme; and a saccharide moiety, which provides specificity for binding. X-ray crystallographic studies of complexes between  $\beta$ -glycosyl hydrolases and such epoxide-saccharide irreversible inhibitors reveal a covalent attachment to active-site carboxylates (Table 1). Such epoxide compounds offer the promise of localizing an enzyme's active site but may not provide information about the enzyme's catalytic mechanism for three reasons. First, additional atoms mediate the attachment of the residual saccharide to the enzyme compared with the natural substrate; given the restricted space in the active site of an enzyme, these inherent additional atoms may complicate any possible conclusions on catalytic mechanism drawn from the structural interpretation. Second, the

site of covalent attachment may vary depending on the epoxide compound used because of the high reactivity of these compounds. In *T. reesei* endo-1,4-xylanase II, either propyl- or pentyl-epoxide compounds bind to the putative catalytic nucleophile Glu86, whereas the butyl-derived ligands bind to the putative acid/base catalyst Glu177 [36]. This makes the assignment of particular functionalities to sidechains problematic. Third, the covalent binding of epoxides leads to irreversible complexes. Hence, the reaction mechanism followed for forming the complex is not catalytically representative for an enzyme.

In contrast to epoxide compounds, 2-fluoro-saccharide compounds can be used to trap the catalytic nucleophile in a state much closer structurally and functionally to the natural mechanism [1,33]. Upon cleavage by a retaining  $\beta$ -glycosidase, such a minimally modified saccharide forms a covalent species specifically with the enzyme's catalytic nucleophile; the complex formed has been shown to be catalytically competent as it slowly turns over to release the reaction product and the free enzyme [31,33,37,38\*]. This strategy has been used for the  $\beta$ -glucanase cex-cd, which has also been extensively studied both structurally and kinetically [23,39].

### Conformation of the catalytic intermediate

A glycosyl–enzyme catalytic intermediate has recently been observed in the crystal structure of the  $\beta$ -glucanase cex-cd with a covalently bound 2-fluoro-cellobioside substrate (Figure 1) [40\*\*]. In this crystal structure, determined at a resolution of 1.8 Å, the saccharide in subsite –1 binds with an inverted  $\alpha$  configuration at the anomeric carbon C1. The configuration agrees with experiments in which the intermediate is trapped

**Table 1**


**Crystal structure of  $\beta$ -glycosidases in complex with epoxide irreversible inhibitor affinity labels.**

Epoxide-saccharide	Labelled enzyme	Labelled residue	Reference
2',3'-epoxypropyl- $\beta$ -glycoside of <i>N</i> -acetyl-D-glucosamine	Hen egg-white lysozyme*	Asp52	[49]
3,4-epoxybutyl- $\beta$ -D-cellubioside	Hybrid <i>Bacillus</i> $\beta$ -1,3-1,4-glucanase	Glu105	[50]
2',3'-epoxypropyl- $\beta$ -glycoside of <i>N,N'</i> -diacetylchitobiose	Human lysozyme	Asp53	[51]
2',3'-epoxypropyl- $\beta$ -glycoside of <i>N</i> -acetylactosamine	Human lysozyme	Asp 53	[51]
3,4-epoxybutyl- $\beta$ -D-cellobioside	<i>Fusarium oxysporum</i> EGI	Glu197	[21]
3,4-epoxypropyl- $\beta$ -D-xyloside	<i>Trichoderma reesei</i> endo-1,4-xylanase II	Glu86	[36]
3,4-epoxybutyl- $\beta$ -D-xyloside	<i>Trichoderma reesei</i> endo-1,4-xylanase II	Glu177	[36]
3,4-epoxypentyl- $\beta$ -D-xyloside	<i>Trichoderma reesei</i> endo-1,4-xylanase II	Glu86	[36]

\*The structure of this complex has not been refined.

using 2-fluoro-saccharides and then analyzed using 1D  $^{19}\text{F}$ -NMR studies [28,41]. The structure also shows that the covalently bound saccharide adopts a 'full chair' cyclic (glucopyranosyl) conformation. This latter observation is in contrast to a previously proposed controversial mechanism [42,43], in which, on the basis of stereoelectronic considerations, the retaining  $\beta$ -glycosyl hydrolase lysozyme would hydrolyze its substrate via an endocyclic opening of the binding saccharide.

**Figure 1**



Structural characteristics of the catalytic intermediate of the retaining  $\beta$ -1,4-glycanase cex-cd. The saccharide is shown in a full chair conformation, forming a covalent link in the inverted ( $\alpha$ ) configuration with the nucleophilic sidechain of the enzyme (lower right). This sidechain, the anomeric carbon C1, and the cyclic oxygen are nearly coplanar (shaded plane); a lone pair of the oxygen is also inferred to be coplanar. The acid/base catalyst (upper right) is positioned to activate a water molecule (not shown) for the subsequent step of the mechanism. The hydroxyl group at the C2 position of the sugar ring (shown below the ring) is positioned to accept a hydrogen bond from a conserved asparagine sidechain and the carbonyl oxygen of the nucleophilic sidechain. Remaining saccharide units are abbreviated by R'.

In the covalent complex of cex-cd [40••], the saccharide at subsite -1 binds at the *syn* position of the (now) esterified sidechain of the nucleophile Glu233. By similarity to carboxylates in solution, this *syn* position is preferred to the *anti* position in chemical reactions [44]. The orientation of the attached saccharide in cex-cd is such that its cyclic oxygen O5 is found to be nearly coplanar with the nucleophile Glu233 sidechain. As a consequence, one lone pair of electrons of O5 is inherently coplanar with the ester group of the sidechain of Glu233. The significance of this coplanarity is not known, although it may have long-range stereoelectronic implications.

In the cex-cd complex, the acid/base catalyst Glu127 is located 4.5 Å from the saccharide C1 and 5.5 Å from the nucleophile Glu233, in agreement with other retaining

$\beta$ -glycosyl hydrolases [1,23]. The distance of 4.5 Å is presumed to allow the positioning of a water molecule between the acid/base catalyst and the anomeric C1 in the deglycosylation step of the mechanism. In a xylanase from *Bacillus circulans*, either shortening or lengthening of the catalytic nucleophile leads to a large decrease in catalytic activity, which confirms the importance of precise positioning of the catalytic nucleophile at the active site [45]. The geometrical disposition of the acid/base catalyst relative to the nucleophile could also explain the observations on the  $pK_a$  regulation discussed above.

Previous kinetic data from other  $\beta$ -glycosyl hydrolases with a variety of substrates have shown that the substituent at the C2 position of the saccharide binding at subsite -1 contributes at least 8 kcal mol $^{-1}$  of binding energy [46–48]. Inspection of the crystal structure of the cex-cd complex reveals that the fluorine substituent at C2 forms two interactions with the enzyme: it accepts a hydrogen bond from Asn126, which is a highly conserved residue among families of  $\beta$ -glycosyl hydrolases [24]; and it is in close proximity with the nucleophile Glu233 O $\epsilon$ 2. Thus, a similarly positioned hydroxyl group at C2 is inferred to form a hydrogen bond with the enzyme's catalytic nucleophile, providing an explanation for the high binding energy. Indeed, this interaction between the noncovalently bonded nucleophile carbonyl and the C2 hydroxyl is seen to be 2.3 Å in a catalytically inactive mutant cex-cd complexed with a natural substrate (V Monem, RAJ Warren, SG Withers, DR Rose, unpublished data).


It is reasonable to speculate that the above structural characteristics of the catalytic intermediate observed with cex-cd are conserved for many retaining  $\beta$ -glycosyl hydrolases. A comparison with other reversible catalytic intermediates will better address this issue.

### Conclusions: mechanism of catalysis by retaining $\beta$ -glycosyl hydrolases

The recent evidence that we have reviewed permits us to expand the description of the catalytic mechanism as follows (Figure 2):

1. Saccharide binding at subsite -1 induces a 'twisted-boat' conformation. A general acid catalyst donates a proton to the scissile  $\beta$ -1,4 oxygen.
2. The  $\beta$ -1,4 linkage is broken, leading to the formation of a transition state with build-up of a positive charge at the anomeric carbon; typically, such a transition state develops a substantial oxocarbenium-ion-like character and resembles a 'half-chair' conformation.
3. A close approach of this ionic species to the catalytic nucleophile leads to the formation of a covalent intermediate of the inverted  $\alpha$  configuration in a 'chair' conformation. The formation of this covalent species promotes a drop of about 2–3 units of pH in the  $pK_a$  of the acid catalyst,

Figure 2



Proposed catalytic mechanism for retaining  $\beta$ -1,4-glycosyl hydrolases. A covalent  $\alpha$ -glycosyl-enzyme intermediate is formed and then hydrolyzed via oxocarbenium-ion-like transition states. For stereoelectronic considerations, the binding saccharide conformation is presumed to undergo conformational dynamics. **1.** The saccharide binds as a twisted-boat. **2.** Structure deforms to a 'half-chair' at the first transition state. **3.** Structure relaxes to a chair at the covalent intermediate. **4.** Structure deforms again to a half-chair at the second transition state. **5.** The product is a twisted-boat conformation.

which becomes a base catalyst. The aglycon is released and a water molecule diffuses into the vicinity of this base catalyst.

4. The covalent species reactivates to pass through an oxocarbenium ion-like transition state similar to the event 2 above. The base catalyst abstracts a proton from the

incoming water molecule, which in turn performs a nucleophilic attack on the C1 of the residual saccharide. We expect that the formation of the product will restore the initial high  $pK_a$  of the acid/base catalyst.

5. This catalysis leads to retention of anomeric configuration at C1.

Results in the near future should confirm or modify aspects of this mechanism, including the generalization of the covalent intermediate observations to other enzyme systems and a further characterization of the transition state species.

## Acknowledgements

Work in the authors' laboratory is supported by the Protein Engineering Network of Centres of Excellence (PENCE) and the National Cancer Institute of Canada. We are grateful to S Withers, RAJ Warren, D Kilburn and N Gilkes and their colleagues for a continuing collaboration on this work and to both A Hadfield and A Mulholland for fruitful discussion.

## References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. McCarter JD, Withers SG: **Mechanisms of enzymatic glycoside hydrolysis.** *Curr Opin Struct Biol* 1994, 4:885-892.
  2. Davies G, Henriissat B: **Structures and mechanisms of glycosyl hydrolases.** *Structure* 1995, 3:853-859.
  3. Koshland DEJ: **Stereochemistry and the mechanism of enzymatic reactions.** *Biol Rev* 1953, 28:416-436.
  4. Imoto T, Johnson LN, North ACT, Phillips DC, Rupley JA: **Vertebrate lysozymes.** In *The Enzymes*. Edited by Boyer PD. New York: Academic Press; 1972:666-868.
  5. Sinnott ML: **Catalytic mechanisms of enzymic glycosyl transfer.** *Chem Rev* 1990, 90:1171-1202.
  6. Kuroki R, Weaver LW, Matthews BW: **Structure-based design of a lysozyme with altered catalytic activity.** *Nat Struct Biol* 1995, 2:1007-1011.
  7. Davies GJ, Wilson KS, Henriissat B: **Nomenclature for sugar-binding subsites in glycosyl hydrolases.** *Biochem J* 1997, 321:557-559.
  8. Johnson LN, Phillips DC: **Structure of some crystalline lysozyme-inhibitor complexes determined by X-ray analysis at 6 Å resolution.** *Nature* 1965, 206:761-763.
  9. Ford LO, Johnson LN, Machin PA, Phillips DC, Tjian R: **Crystal structure of a lysozyme-tetrasaccharide lactone complex.** *J Mol Biol* 1974, 88:349-371.
  10. Strynadka NCJ, James MNG: **Lysozyme revisited: crystallographic evidence for distortion of an N-acetylmuramic acid residue bound in site D.** *J Mol Biol* 1991, 220:401-424.
  11. Hadfield AT, Harvey DJ, Archer DB, MacKenzie DA, Jeenes DJ, Radford SE, Lowe G, Dobson CM, Johnson LN: **Crystal structure of the mutant D52S hen egg white lysozyme with an oligosaccharide product.** *J Mol Biol* 1994, 243:846-872.
  12. Kuroki R, Weaver LH, Matthews BW: **A covalent enzyme-substrate intermediate with saccharide distortion in a mutant T4 lysozyme.** *Science* 1993, 262:2030-2033.
  13. Deslongchamps P: **Stereoelectronic Effects in Organic Chemistry.** Oxford: Pergamon Press; 1983.
  14. Varghese JN, McKimm-Breschkin JL, Caldwell JB, Kortt AA, Colman PM: **The structure of the complex between influenza virus neuraminidase and sialic acid, the rival receptor.** *Proteins* 1992, 14:327-332.
  15. Van den Akker F, Steensma E, Hol WGJ: **Tumor marker disaccharide D-Gal-β1,3-GalNAc complexed to heat-labile enterotoxin from *Escherichia coli*.** *Protein Sci* 1996, 5:1184-1188.
  16. Sakon J, Adney WS, Himmel ME, Thomas SR, Karplus PA: **Crystal structure of thermostable Family 5 endoglucanase E1 from *Acidothermus cellulolyticus* in complex with cellotetraose.** *Biochemistry* 1996, 35:10648-10660.
  - This report presents the crystal structure of the thermostable endoglucanase E1 determined in the presence of cellobiose. Interestingly, the enzyme in the crystal condenses two disaccharides by transglycosylation, thus producing a cellotetraose which stays bound in the enzyme's active site. Although the geometry of the saccharide binding at subsite -1 is constrained to a relaxed 'chair' conformation in the model, it is recognized that this saccharide unit may have a strained conformation away from a chair. This report also gives an scholarly account of the structural context of conserved residues for the Family 5 glycosyl hydrolases; the discussion is then extended to the 4/7 superfamily of enzymes. These conserved residues are incorporated with the catalytic mechanism for a better understanding of catalysis by retaining β-glycosidyl hydrolases. A close approach of the two key catalytic glutamate residues is also proposed to account for the large shift in the pK<sub>a</sub> of the acid/base catalyst during catalysis.
  17. Alzari PM, Souchon H, Dominguez R: **The crystal structure of endoglucanase CelA, a Family 8 glycosyl hydrolase from *Clostridium thermocellum*.** *Structure* 1996, 4:265-275.
  - The high-resolution crystal structure of CelA, a Family 8 glycosyl hydrolase, is described. On the basis of binding of cellobiosaccharides and modified cellobioside, five saccharide subsites are identified. A further comparison of CelA with Family 9 endoglucanase CelD and with Family 15 glucoamylase-I enzymes reflects the stability of their common (α/α)<sub>6</sub> barrel folding motif and helps to better understand the evolutionary relationship between different families of β-glycosyl hydrolases.
  18. Tews I, Perrakis A, Oppenheim A, Dauter Z, Wilson KS, Vorgias CE: **Bacterial chitinase structure provides insights into catalytic mechanism and the basis of Tay-Sachs disease.** *Nat Struct Biol* 1996, 3:638-648.
  - This crystal structure of an abortive complex of a di-N-acetyl-glucosamine bound to a chitinase clearly shows one distorted saccharide at the binding subsite -1 in a 'twisted-boat' conformation, whereas the other sugar is in a typical chair form. The enzyme is believed to 'dock' the substrate into its active-site pocket. The proposed mechanism of catalysis involves substrate assistance by the N-acetyl substituent at C2.
  19. Thunnissen A-MWH, Rozeboom HJ, Kalk KH, Dijkstra BW: **Structure of the 70 kDa soluble lytic transglycosylase complexed with bulgenin A. Implications for the enzymatic mechanism.** *Biochemistry* 1995, 34:12729-12737.
  20. Sulzenbacher G, Driguez H, Henriissat B, Schülein M, Davies GJ: **Structure of the *Fusarium oxysporum* endoglucanase I with a nonhydrolyzable substrate analogue: substrate distortion gives rise to the preferred axial orientation for the leaving group.** *Biochemistry* 1996, 35:15280-15287.
  - Assessing the conformation of the saccharide at the subsite -1 for retaining β-glycosyl hydrolases has always been difficult. In this paper, a nonhydrolyzable thio-oligosaccharide is successfully used with an endoglucanase I to show the deformation of the saccharide binding at subsite -1. In this crystal structure, the saccharide spans the cleavage centre of the enzyme and shows a 'twisted-boat' conformation. Hence, the scissile bond is axial to this saccharide, which may be important for catalysis.
  21. Sulzenbacher G, Schülein M, Davies GJ: **Structure of the endoglucanase I from *Fusarium oxysporum*: native, cellobio, and 3,4-epoxybutyl β-D-cellobioside-inhibited forms, at 2.3 Å resolution.** *Biochemistry* 1997, 36:5902-5911.
  22. Aleshin AE, Stoffer B, Firsov LM, Svensson B, Honzatko RB: **Crystallographic complexes of glucoamylase with maltooligosaccharide analogs: relationship of stereochemical distortions at the nonreducing end to the catalytic mechanism.** *Biochemistry* 1996, 35:8319-8328.
  - High-resolution crystal structures of the inverting enzyme glucoamylase complexed with D-glucosyl-dihydroacarbonyl and acarbonyl show saccharide distortion at the catalytic center toward a half-chair conformation. This study also reports an analysis of the extent of such distortion as a function of pH.
  23. White A, Withers SG, Gilkes NR, Rose DR: **Crystal structure of the catalytic domain of the β-1,4-glycanase Cex from *Cellulomonas fimi*.** *Biochemistry* 1994, 33:12546-12552.
  24. Dominguez R, Souchon H, Lascombe M-B, Alzari PM: **The crystal structure of a Family 5 endoglucanase mutant in complexed and uncomplexed forms reveals an induced fit activation mechanism.** *J Mol Biol* 1996, 257:1042-1051.

25. Kempton JB, Withers SG: **Mechanism of *Agrobacterium*  $\beta$ -glucosidase: kinetic studies.** *Biochemistry* 1992, **31**:9961-9969.
26. McIntosh LP, Hand G, Johnson PE, Joshi MD, Körner M, Plesniak LA, Ziser L, Wakarchuk WW, Withers SG: **The  $pK_a$  of the general acid/base carboxyl group of a glycosidase cycles during catalysis: a  $^{13}\text{C}$ -NMR study of *Bacillus circulans* xylanase.** *Biochemistry* 1996, **35**:9958-9966.
- This  $^{13}\text{C}$ -NMR study reveals the  $pK_a$  of the acid/base catalyst Glu172 in the context of a covalent intermediate both for the catalytic nucleophile and for a mutant of that nucleophile. In these two cases, a large  $pK_a$  shift is observed for the acid/base catalyst Glu78, compared with the free enzyme. This phenomenon is presumed to be a common feature for glycosidases.
27. Plesniak LA, Wakarchuk WW, McIntosh LP: **Secondary structure and NMR assignments of *Bacillus circulans* xylanase.** *Protein Sci* 1996, **5**:1118-1135.
28. Tull D, Withers SG: **Mechanisms of cellulases and xylanases: a detailed kinetic study of the exo- $\beta$ -1,4-glycanase from *Cellulomonas fimi*.** *Biochemistry* 1994, **33**:6363-6370.
29. Richard JP, Westerfeld JG, Lin S: **Structure/reactivity relationships for  $\beta$ -galactosidase (*Escherichia coli*, *lac Z*). 1. Brønsted parameters for cleavage of alkyl  $\beta$ -D-galactopyranosides.** *Biochemistry* 1995, **34**:11703-11712.
30. MacLeod AM, Tull D, Rupitz K, Warren RAJ, Withers SG: **Mechanistic consequences of mutation of active site carboxylates in a retaining  $\beta$ -1,4-glycanase from *Cellulomonas fimi*.** *Biochemistry* 1996, **35**:13165-13172.
- This report presents a new method to identify the catalytic nucleophile of retaining  $\beta$ -glycosyl hydrolases, without the use of active-site labelling agents. Results of activity assays for enzyme mutants at conserved aspartate/glutamate positions are indicative of the possible catalytic function of the mutated residue. As a test case, results obtained with three sidechains of the  $\beta$ -1,4-glycanase cex-cd correlate with their expected role. This method is likely to be applicable to other retaining  $\beta$ -glycosyl hydrolases.
31. Withers SG, Street IP, Bird P, Dolphin DH: **2-Deoxy-2-fluoroglucosides: a novel class of mechanism-based glucosidase inhibitors.** *J Am Chem Soc* 1987, **109**:7530-7531.
32. Withers SG, Rupitz K, Street IP: **2-Deoxy-2-fluoro-D-glycosyl fluorides. A new class of specific mechanism-based glycosidase inhibitors.** *J Biol Chem* 1988, **263**:7929-7932.
33. Withers SG, Aebersold R: **Approaches to labeling and identification of active site residues in glycosidases.** *Protein Sci* 1995, **4**:361-372.
34. Terwisscha van Scheltinga AC, Hennig M, Dijkstra BW: **The 1.8 Å resolution structure of hevamine, a plant chitinase/lysozyme, and analysis of the conserved sequence and structure motifs of glycosyl hydrolase Family 18.** *J Mol Biol* 1996, **262**:243-257.
35. Ståhlberg J, Divne C, Koivula A, Piens K, Claeysens M, Teeri TT, Jones TA: **Activity studies and crystal structures of catalytically deficient mutants of cellobiohydrolase I from *Trichoderma reesei*.** *J Mol Biol* 1996, **264**:337-349.
36. Havukainen R, Törrönen A, Laitinen T, Rouvinen J: **Covalent binding of three epoxyalkyl xylosides to the active site of endo-1,4-xylanase II from *Trichoderma reesei*.** *Biochemistry* 1996, **35**:9617-9624.
37. Withers SG, Warren RAJ, Street IP, Rupitz K, Kempton JB, Aebersold R: **Unequivocal demonstration of the involvement of a glutamate residue as a nucleophile in the mechanism of a 'retaining' glycosidase.** *J Am Chem Soc* 1990, **112**:5887-5889.
38. Mackenzie LF, Davies GJ, Schülein M, Withers SG: **Identification of the catalytic nucleophile of endoglucanase I from *Fusarium oxysporum* by mass spectrometry.** *Biochemistry* 1997, **36**:5893-5901.
- 2-fluoro-saccharides are used to inactivate endoglucanase from *Fusarium oxysporum* and thus to identify the catalytic nucleophile. The saccharide-enzyme complex is shown to be catalytically competent, undergoing transglycosylation with D-cellobiose.
39. Bray MR, Creagh AL, Damude HG, Gilkes NR, Haynes CA, Jervis E, Kilburn DG, MacLeod AM, Meinke A, Miller RC Jr et al.:  **$\beta$ -1,4-glycanases of *Cellulomonas fimi*: families, mechanisms and kinetics.** In *Enzymes for Pulp and Paper Processing*, Symposium series no 655. Edited by Jeffries TW, Viikari L. Washington: American Chemical Society; 1997:in press.
40. White A, Tull D, Johns K, Withers SG, Rose DR: **Crystallographic observation of a covalent intermediate in a  $\beta$ -glycosidase.** *Nat Struct Biol* 1996, **3**:149-154.
- A minimally modified slow substrate is used to observe the structural features of the catalytic intermediate in a retaining  $\beta$ -glycosyl hydrolase. Upon cleavage of the disaccharide substrate by the enzyme in the crystal, a reversible covalent glycosyl-enzyme complex is formed.
41. Withers SG, Street IP: **Identification of a covalent  $\alpha$ -D-glucopyranosyl enzyme intermediate formed on a  $\beta$ -glucosidase.** *J Am Chem Soc* 1988, **110**:8551-8553.
42. Post CB, Karplus M: **Does lysozyme follow the lysozyme pathway? An alternative based on dynamics, structural, and stereoelectronic considerations.** *J Am Chem Soc* 1986, **108**:1317-1319.
43. Sinnott ML: **There is no experimental evidence for endocyclic cleavage in the action of erne (retaining  $\beta$ ) glycopyranosidases, and much against it.** *Bioorg Chem* 1993, **21**:34-40.
44. Gandour RD: **On the importance of orientation in general base catalysis by carboxylate.** *Bioorg Chem* 1981, **10**:169-176.
45. Lawson SL, Wakarchuk WW, Withers SG: **Effects of shortening and lengthening the active site nucleophile of *Bacillus circulans* xylanase on catalytic activity.** *Biochemistry* 1996, **35**:10110-10118.
46. Roeser K-R, Legler G: **Role of sugar hydroxyl groups in glycoside hydrolysis – cleavage mechanism of deoxyglucosides and related substrates by  $\beta$ -glucosidase  $A_3$  from *Aspergillus wentii*.** *Biochim Biophys Acta* 1981, **657**:321-333.
47. Wolfenden R, Kati WM: **Testing the limits of protein-ligand binding discrimination with transition-state analogue inhibitors.** *Acc Chem Res* 1991, **24**:209-215.
48. McCarter JD, Adams MJ, Withers SG: **Binding energy and catalysis: fluorinated and deoxygenated glycosides as mechanistic probes of *Escherichia coli* (*lac Z*)  $\beta$ -galactosidase.** *Biochem J* 1992, **286**:721-727.
49. Moults J, Eshdat YE, Sharon N: **The identification by X-ray crystallography of the site of attachment of an affinity label to hen egg-white lysozyme.** *J Mol Biol* 1973, **75**:1-4.
50. Keitel T, Simon O, Borriss R, Heinemann U: **Molecular and active-site structure of a *Bacillus* 1,3-1,4- $\beta$ -glucanase.** *Proc Natl Acad Sci USA* 1993, **90**:5287-5291.
51. Muraki M, Harata K, Sugita N, Sato K-I: **Origin of carbohydrate recognition specificity of human lysozyme revealed by affinity labeling.** *Biochemistry* 1996, **35**:13562-13567.