

DARKSTAR DYNAMICAL MODEL — v7

Scott Hunter Hughes “DarkStar Aurora Shockwave”

October 1, 2026

DARKSTAR DYNAMICAL MODEL — v7

Objective: construct the smallest mathematical model capable of distinguishing ordinary black-hole transformation and feedback from the additional DarkStar hypothesis of cosmic recycling. The model is intentionally conservative: every new DarkStar term is isolated so it can be tested against zero.

1. State Variables

Let $M_{BH}(t)$ be black-hole mass; $M_g(t)$ the surrounding gas reservoir; $E_{rad}(t)$ the radiative energy stored in the selected control volume; $E_{out}(t)$ the cumulative mechanical/radiative outflow energy; $S(t)$ the generalized entropy budget; and $a(t)$ the cosmological scale factor when the model is extended to cosmological scales.

Introduce $f_{rec}(t)$, a dimensionless recycling fraction. It is not assumed to be nonzero. It represents only a channel that survives after known accretion-disk, jet, wind, stellar, and environmental processes have been modeled.

2. Black-Hole Growth Equation

$$dM_{BH}/dt = (1-\epsilon) \dot{M}_{acc} - \dot{M}_{DH}.$$

Here $\dot{M}_{acc}$ is the accretion rate reaching the black-hole system, ϵ is the fraction of accretion rest-mass energy emerging through modeled radiative/mechanical channels, and $\dot{M}_{DH}$ is a placeholder for any genuinely new DarkStar mass-transfer term. In the null model, $\dot{M}_{DH} = 0$.

This equation prevents the word “recycling” from silently becoming extra mass. Any proposed DarkStar channel must appear explicitly in the ledger.

3. Energy Ledger

$$P_{acc} = \dot{M}_{acc} c^2; P_{out} = P_{rad} + P_{jet} + P_{wind} + P_{other}.$$

$$d(M_{BH} c^2 + E_{env})/dt = P_{in} - P_{out} + P_{ext}.$$

E_{env} is the energy stored in the defined environment and P_{ext} represents energy crossing the outer control boundary. The model is valid only after the boundary is explicitly defined.

4. Recycling Channel

$$\dot{M}_{rec} = f_{rec} \dot{M}_{in}.$$

For a first test, compare two models: H0: $f_{rec} = 0$ after established channels are included; H1: f_{rec} is nonzero and follows a specified functional dependence, such as $f_{rec} = f(M_{BH}, \dot{M}_{acc}, \text{spin}, \text{environment})$. H1 becomes scientifically meaningful only if it produces a statistically significant residual and survives independent modeling of known physics.

5. Composition and Spectral Test

Let X_i be the mass fraction of particle or elemental species i in the outflow. DarkStar predicts no particular new composition until a physical transformation mechanism is specified. Therefore the first useful test is comparative:

does the measured vector X_i , together with spectra and momentum flux, differ from the distribution predicted by standard accretion, stellar, nuclear, and jet/wind models?

A DarkStar-specific transformation mechanism must then predict a distinctive ΔX_i , spectral feature, particle population, or correlation. Without such a signature, observed outflows remain compatible with established astrophysical feedback.

6. Entropy Constraint

$$dS_{\text{gen}}/dt \geq 0.$$

The generalized entropy S_{gen} must include the relevant matter/radiation entropy and black-hole entropy. For a semiclassical black hole, $S_{\text{BH}} = k_B c^3 A/(4G\hbar)$. DarkStar cannot obtain a cyclic “cleaning” process by simply deleting entropy from the ledger. Any local decrease must be compensated by entropy production elsewhere so that the generalized second law is respected.

7. Horizon Boundary Condition

In classical general relativity, the event horizon is a causal boundary: an ordinary future-directed causal trajectory that crosses it does not return to the same exterior region. NASA describes the horizon as the point of no return and states that what happens after crossing is not accessible to an outside observer. [NASA sources]

Therefore an interior-to-exterior DarkStar channel cannot be inserted as an ordinary outflow term without changing the spacetime model. It requires at least one of the following: a nonclassical causal structure, a quantum-gravity mechanism, a topology-changing/bounce construction, or another explicitly defined modification.

8. Minimal Interior-to-Exterior Ansatz

If DarkStar is to pursue the “other side” idea mathematically, introduce two regions, an exterior E and a proposed daughter/exit region D , joined by a transition surface Σ . The model must specify a metric $g_{\mu\nu}$ on both regions and matching conditions on Σ .

$$G_{\mu\nu}(E) = (8\pi G/c^4)T_{\mu\nu}(E), \quad G_{\mu\nu}(D) = (8\pi G/c^4)T_{\mu\nu}(D).$$

The transition must preserve the relevant constraint equations and specify whether Σ is timelike, null, or spacelike. It must also state whether the exit is in the same universe, a causally disconnected region, or a new expanding region. Without those choices, “other side” is a metaphor rather than a mathematical object.

9. Why This Is a Hard Test

The event-horizon problem is not merely an engineering obstacle. General relativity predicts a one-way causal structure, while quantum gravity is expected to be needed in the regime where the classical singularity appears. NASA notes that general relativity is widely expected to be incomplete because it lacks a quantum foundation. [NASA sources]

Modern work on the black-hole information problem demonstrates that the relationship between interior information and exterior radiation is itself an active theoretical problem. That does not establish a DarkStar exit channel, but it means the interior should be treated as an open research frontier rather than as experimentally observed plumbing. [research literature]

10. Cosmological Extension

$$H^2 = (8\pi G/3)\rho - kc^2/a^2 + \Lambda c^2/3.$$

For standard components, $\rho_r \propto a^{-4}$ and $\rho_m \propto a^{-3}$. A DarkStar cycle requires a modified dynamical term or component capable of producing a turnaround or bounce. Define an effective DarkStar density $\rho_{\text{DS}}(a)$ and pressure $p_{\text{DS}}(a)$, but do not assign them arbitrary values. They must arise from an underlying mechanism.

$$\ddot{a}/a = -(4\pi G/3)(\rho + 3p/c^2) + \Lambda c^2/3.$$

The decisive mathematical question is whether there exists a physically allowed state with $H = 0$ followed by $H > 0$, while maintaining nonnegative generalized entropy production and consistency with observations.

11. The Critical Null-Test Strategy

The model should first be run with every DarkStar-specific term set to zero. This is H_0 , the established-physics baseline. Then introduce one new term at a time. If the new term improves an observable fit but is mathematically redundant with an established parameter, it is not a distinct DarkStar prediction.

The strongest result would be a robust observable that H_0 cannot reproduce within uncertainties but H_1 predicts before the observation is used for fitting.

12. Current Mathematical Conclusion

The dynamical model does not yet prove an interior recycling mechanism. It does something more valuable at this stage: it identifies the exact mathematical doorway through which that claim must pass.

DarkStar now has a falsifiable hierarchy: ordinary astrophysical recycling → measurable residual recycling fraction → specified transformation signature → explicit interior/exit spacetime model → cosmological-cycle dynamics. Failure at any layer constrains the next layer rather than being hidden by metaphor.

Author and Concept Nomenclature

“DarkStar” denotes the scientific hypothesis. The author is Scott Hunter Hughes “DarkStar Aurora Shockwave.” The author's personal philosophy is separate from this mathematical model and is not presented as scientific evidence.