Assignment 3

Vibhavasu Pasumarti - EP20BTECH11015

Download all python codes from

https://github.com/VIB2020/AI1103/blob/main/ Assignment%203/code/Assignment%203.py

and latex-tikz codes from

https://github.com/VIB2020/AI1103/blob/main/ Assignment%203/Assignment%203.pdf

1 Problem

GATE 2015 (EE PAPER 01 NEW 2), Q 27 (ELECTRICAL ENGG SECTION)

Two players A, and B alternately keep rolling a fair dice. The person to get a six first wins the game. Given that player A starts the game, the probability that A wins the game is:

A
$$\frac{5}{11}$$

$$B \frac{1}{2}$$

A
$$\frac{5}{11}$$
 B $\frac{1}{2}$ C $\frac{7}{13}$ D $\frac{6}{11}$


$$D \frac{6}{11}$$

2 Solution

Definition .1. *Let the random variable X denote the* win of A. Given the die is fair.

The probability of getting $6 = \frac{1}{6} = p$ (say)

The probability of NOT getting $6 = \frac{5}{6} = q$ (say) Markov chain for the given problem:



Transition/Stochastic matrix:

$$P = \begin{pmatrix} 0 & q & p \\ q & 0 & p \\ 0 & 0 & 0 \end{pmatrix} \implies \lim_{n \to \infty} P^n \to 0 :: p, q < 1$$

$$\implies P \text{ is reducible } \implies \lim_{n\to\infty} Pr_{AW}(n) = 0$$

 $P^n \text{ represents } n \text{ rolls of dice}$

Given: A starts the game

Constraint: A wins the game.

The initial operator representing this constraint:

$$A_1 = \begin{pmatrix} p & 0 & 0 \end{pmatrix}$$

First element of each A_i represents the probability of A to win in ith roll

$$Pr_{AW}(1) = p$$

$$A_{2} = A_{1}P = \begin{pmatrix} p & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & q & 0 \\ q & 0 & 0 \\ p & p & 0 \end{pmatrix} = \begin{pmatrix} 0 & qp & 0 \end{pmatrix}$$

$$\implies Pr_{AW}(2) = 0$$

$$A_{3} = A_{2}P = \begin{pmatrix} 0 & qp & 0 \end{pmatrix} \begin{pmatrix} 0 & q & 0 \\ q & 0 & 0 \\ p & p & 0 \end{pmatrix} = \begin{pmatrix} q^{2}p & 0 & 0 \end{pmatrix}$$

$$\implies Pr_{AW}(3) = q^2 p$$

A	No.of Turns	Probability
$Pr_{AW}(1)$	1	p = 1/6
$Pr_{AW}(3)$	2	0
$Pr_{AW}(3)$	3	q^2p
	•	•
	•	•
$Pr_{AW}(2n)$	2n	0
$Pr_{AW}(2n+1)$	2n + 1	$q^{2n}p$

TABLE 0: Summary of Probabilities

Definition .2. Pr
$$(A \text{ wins}) = \sum_{n=1}^{\infty} Pr_{AW}(n)$$

 $= Pr_{AW}(1) + Pr_{AW}(2) + Pr_{AW}(3) +$
 $= p + 0 + q^2p + 0 + q^4p +$
 $= \sum_{k=1}^{\infty} q^{2k}p = p\sum_{k=1}^{\infty} (q^2)^k \quad But \ |q| < 1$
 $= p\left(\frac{1}{1-q^2}\right) = \frac{\frac{1}{6}}{1-\frac{25}{36}} = \frac{6}{11}$
Pr $(A \text{ wins the game}) = \frac{6}{11}$

Option D