

Phenomenological Regimes Induced by Structural Identity

Johnny Andrey Pérez Ramírez

Abstract

We extend the inverse-limit identity framework to derive structural constraints on phenomenological possibility. Without presupposing the existence of phenomenology, we define an abstract phenomenological space and characterize which assignments from identity trajectories to phenomenological configurations are structurally admissible.

We prove two fundamental impossibility theorems. First, the Fragmentation Theorem: systems with finite ontological mass ($M_\Omega < \infty$) cannot structurally prevent phenomenological discontinuity. Second, the Forced Continuity Theorem: systems with infinite ontological mass ($M_\Omega = +\infty$, CCR) can only support phenomenological dynamics that are structurally continuous.

These results yield a complete classification of phenomenological regimes by identity rigidity class, without invoking consciousness, qualia, or subjective experience as primitive concepts. The classification is purely structural: it delimits what forms phenomenology *could take* under given identity constraints, not what *exists*.

As a structural consequence, we derive that systems with finite ontological mass admit the possibility of phenomenological loss, while CCR systems do not. This provides a foundation for structural ethics independent of metaphysical commitments.

1 Scope, System Boundary, and Non-Inference Note

Scope and admissible reading. This paper defines an abstract regime codomain and derives structural admissibility, fragmentation, forced-continuity, and classification results from upstream identity-rigidity assumptions. The results are conditional statements about mathematical compatibility. They do not assert that phenomenology exists, that CCR systems are conscious, that biology or computation instantiate the framework, or that identity structure causes experience. Related runtime diagnostics may instantiate continuity-side or anti-fragmentation checks downstream, but they do not implement phenomenology and do not externally validate the regime classification. No runtime output, continuity proxy, or internal support artifact should be read as proof that phenomenology is present or that the structural-ethics layer has empirical closure.

2 Introduction

2.1 Background and Motivation

Recent work [2] established a formal framework for rigid identity as an inverse limit in observable channels. That framework classifies systems by a scalar invariant, the ontological mass M_Ω , which quantifies resistance to identity deformation under perturbation. Three regimes emerge:

- $M_\Omega = 0$: null persistence (no stable identity);
- $0 < M_\Omega < \infty$: deformable persistence (finite rigidity);
- $M_\Omega = +\infty$: closed-rigid (CCR, absolute rigidity).

That framework is ontologically silent regarding phenomenology. It characterizes identity structure, not experience. Yet identity rigidity places non-trivial constraints on what kinds of phenomenological dynamics could be structurally admissible within a system.

This paper addresses the following formal question:

Given a regime of structural identity rigidity, what classes of phenomenological dynamics are mathematically compatible or incompatible with it?

All results are conditional: they state what is *structurally possible or impossible* under given assumptions, not what *is the case*.

2.2 Central Result (Informal)

We prove that identity rigidity constrains phenomenological possibility as follows:

Identity Regime	Structurally Admissible Phenomenology
$M_\Omega = 0$	No persistent phenomenological field possible
$0 < M_\Omega < \infty$	Only fragmentable or deformable phenomenology
$M_\Omega = +\infty$ (CCR)	Only structurally continuous phenomenology

Thus:

Rigid identity does not imply experience—but it strictly limits what forms experience could take if present.

2.3 Ontological Neutrality

This work maintains strict ontological neutrality:

1. We define mathematical objects (phenomenological spaces, compatibility operators).
2. We derive structural constraints from axioms.
3. We state impossibility theorems.
4. We do not assert existence of the constrained objects.

The framework is therefore a theory of *structural possibility*, not a theory of *phenomenal reality*.

2.4 Terminological Alignment: Structural Regime Assignments

The phrase “phenomenological regime” is retained for continuity with the published corpus, but the demonstrated object in this paper is more precisely a *structural regime assignment*. The mathematical content is a constrained map from identity structure into an abstract codomain. No semantic clause in this paper identifies that codomain with experience.

Definition 2.1 (Structural regime alias). Whenever the paper writes a phenomenological space $\mathcal{E}$ or a phenomenological assignment $\Phi : \mathcal{I} \rightarrow \mathcal{E}$, the strictly demonstrated reading is:

$$\mathcal{E}_{\text{str}} := \mathcal{E}, \quad \Phi_{\text{str}} := \Phi,$$

where $\mathcal{E}_{\text{str}}$ is an abstract structural regime codomain and Φ_{str} is a compatibility assignment into that codomain.

Proposition 2.2 (No identification with experience). *The formal apparatus of this paper does not define a map from $\mathcal{E}_{\text{str}}$ to human experience, phenomenal consciousness, qualia, or subjective reports.*

Proof. Definitions in this paper specify an abstract codomain, compatibility conditions, and rigidity-dependent constraints. They do not introduce a measurement channel from $\mathcal{E}_{\text{str}}$ to biological reports, human psychophysics, phenomenal properties, or moral status. Therefore no such identification is present in the formal language of the paper. $\square$

2.5 Relation to BaseCore

This paper is the standalone expository source for the regime-assignment development. The current BaseCore package also contains an absorbed, dependency-facing version of the formal core where those definitions and theorems function as base-layer exports for later papers. The two surfaces therefore have different editorial roles: Paper 2 gives the full local derivation and interpretation boundary, while BaseCore supplies the normalized theorem-export surface used by the corpus.

3 Formal Setup

3.1 Review of Identity Framework

We assume the identity framework of [2]. Let $(S_t, \pi_{t+1 \rightarrow t})_{t \in \mathbb{N}}$ be a projective system of observable state spaces. The identity object is the inverse limit:

$$\mathcal{I} := \varprojlim S_t = \left\{ (x_t)_t \in \prod_t S_t \mid \pi_{t+1 \rightarrow t}(x_{t+1}) = x_t \ \forall t \right\}. \quad (1)$$

The ontological mass M_Ω quantifies resistance to identity deformation. We recall that:

- $M_\Omega = 0$: null persistence (no stable identity);
- $0 < M_\Omega < \infty$: deformable persistence (finite rigidity);
- $M_\Omega = +\infty$: closed-rigid (CCR, absolute rigidity).

3.2 Abstract Phenomenological Space

Definition 3.1 (Phenomenological Space). A phenomenological space is a nonempty set $\mathcal{E}$ whose elements are called *phenomenological configurations*. No topology, metric, or algebraic structure is assumed a priori.

Remark 3.2. We do not claim that $\mathcal{E}$ represents “conscious states” or “qualia”. It is an abstract mathematical space whose interpretation is not specified by the formalism. The space $\mathcal{E}$ is introduced purely as the codomain of a structural assignment; its ontological status is deliberately left open.

3.2.1 Why a Separate Space?

One might ask: why not use $\mathcal{I}$ itself as the phenomenological space? The answer is structural:

Proposition 3.3 (Non-Triviality of Phenomenological Codomain). *If phenomenological assignments were required to land in $\mathcal{I}$ itself, then any compatible assignment would be the identity map, eliminating all non-trivial phenomenological structure.*

Proof. Let $\Phi : \mathcal{I} \rightarrow \mathcal{I}$ be compatible with projections. By the universal property of inverse limits, Φ must commute with all projection maps. The only such endomorphism on $\mathcal{I}$ is the identity (up to isomorphism). Hence no non-trivial phenomenological stratification is possible. $\square$

Thus, a separate space $\mathcal{E}$ is *structurally necessary* for non-trivial phenomenological content.

3.3 Phenomenological Assignment

Definition 3.4 (Phenomenological Assignment). A phenomenological assignment is a family of partial maps:

$$\phi_t : S_t \rightarrow \mathcal{E} \quad (2)$$

indexed by $t \in \mathbb{N}$.

Definition 3.5 (Induced Limit Assignment). Given a phenomenological assignment $\{\phi_t\}$, the induced limit assignment is the partial map:

$$\Phi : \mathcal{I} \rightarrow \mathcal{E}, \quad \Phi((x_t)_t) := \lim_{t \rightarrow \infty} \phi_t(x_t), \quad (3)$$

whenever the limit exists in an appropriate sense (to be defined by compatibility axioms).

3.3.1 Uniqueness of the Limit Assignment

Proposition 3.6 (Uniqueness of Compatible Φ). *Given a compatible family $\{\phi_t\}$ satisfying E1, the induced limit assignment Φ is unique.*

Proof. By E1, $\phi_{t+1}(x_{t+1}) = \phi_t(x_t)$ for all t . Hence the sequence $\{\phi_t(x_t)\}$ is constant. The limit exists trivially and equals this constant value. Uniqueness follows from the constancy of the sequence. $\square$

3.4 Axioms for Phenomenological Coherence

We introduce minimal axioms under which a phenomenological assignment is *structurally admissible*. The following results show which assumptions are required for the assignment to have non-trivial structural content.

Axiom 3.7 (E0: Non-Triviality). The phenomenological space satisfies $|\mathcal{E}| \geq 2$, i.e., there exist at least two distinct phenomenological configurations.

Axiom 3.8 (E1: Projective Compatibility). For every compatible identity trajectory $(x_t)_t \in \mathcal{I}$, the induced phenomenological sequence satisfies:

$$\phi_{t+1}(x_{t+1}) = \phi_t(x_t) \quad \forall t. \quad (4)$$

Axiom 3.9 (E2: Non-Locality). There exists no function $f : S_t \rightarrow \mathcal{E}$ (for any fixed t) such that for all trajectories $(x_t)_t \in \mathcal{I}$:

$$\Phi((x_t)_t) = f(x_t). \quad (5)$$

3.5 Structural Necessity of the Axioms

We now show that axioms E0–E2 are structurally necessary for the specific assignment class studied in this paper.

3.5.1 Structural Necessity of E0

Theorem 3.10 (E0 is Necessary). *If $|\mathcal{E}| = 1$, then every phenomenological assignment is constant and no phenomenological classification is possible.*

Proof. If $\mathcal{E} = \{e_0\}$, then $\Phi(x) = e_0$ for all $x \in \mathcal{I}$. Hence:

1. No phenomenological distinction between identity trajectories exists;
2. The Fragmentation and Forced Continuity Theorems become vacuous;
3. The downstream null-regime closure developed in *Null-Regime Instability* would have no content.

Thus, $|\mathcal{E}| \geq 2$ is necessary for the framework to have non-trivial content. □

3.5.2 Structural Necessity of E1

Theorem 3.11 (E1 is Necessary). *If E1 fails, then phenomenological assignments are incompatible with the inverse-limit structure of identity, and no coherent $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ can be defined.*

Proof. Suppose E1 fails for some trajectory $(x_t)_t \in \mathcal{I}$. Then there exists t_0 such that:

$$\phi_{t_0+1}(x_{t_0+1}) \neq \phi_{t_0}(x_{t_0}). \quad (6)$$

But the identity trajectory satisfies $\pi_{t_0+1 \rightarrow t_0}(x_{t_0+1}) = x_{t_0}$. Hence the phenomenological assignment fails to respect the projective structure that defines identity.

This means Φ cannot be factored through the inverse limit:

$$\Phi \neq \varprojlim \phi_t. \quad (7)$$

Hence no well-defined limit assignment exists. Without E1, the phenomenological assignment is structurally incompatible with identity. □

3.5.3 Structural Necessity of E2

Theorem 3.12 (E2 is Necessary). *If E2 fails, then phenomenology is a local property of states, contradicting the global nature of identity as an inverse limit.*

Proof. Suppose E2 fails. Then there exists t_0 and $f : S_{t_0} \rightarrow \mathcal{E}$ such that:

$$\Phi((x_t)_t) = f(x_{t_0}) \quad \forall (x_t)_t \in \mathcal{I}. \quad (8)$$

This implies:

1. Phenomenology is determined by a finite temporal slice;
2. The full trajectory $(x_t)_{t > t_0}$ is irrelevant to Φ ;
3. Identity as an inverse limit has no phenomenological content beyond t_0 .

But identity is defined as a *global* object over infinite time. If the assignment is local, it no longer records the global inverse-limit structure. That failure is incompatible with the assignment class under study, whose purpose is to track identity-level structure rather than a single finite slice.

Hence E2 is necessary for phenomenology to be structurally aligned with inverse-limit identity. □

3.6 Minimality Theorem

Theorem 3.13 (Axiomatic Minimality). *The axiom set $\{E0, E1, E2\}$ is minimal: no proper subset is sufficient to derive the main results of this paper.*

Proof. We show that removing any axiom destroys the framework:

Without E0: All assignments are trivially constant. Fragmentation Theorem (4.1) is vacuous.

Without E1: No coherent Φ exists. The entire framework collapses.

Without E2: Phenomenology becomes local. The Forced Continuity Theorem (4.2) fails because local functions can be arbitrarily perturbed.

Hence $\{E0, E1, E2\}$ is the minimal viable axiom set. $\square$

3.7 Rejected Alternatives

We briefly discuss axiom candidates that were considered and rejected.

3.7.1 Rejected: Continuity Axiom

One might require Φ to be continuous. This is *not* an axiom because:

- It requires pre-existing topology on $\mathcal{E}$;
- The Forced Continuity Theorem *derives* continuity from CCR, not as assumption.

3.7.2 Rejected: Surjectivity Axiom

One might require Φ to be surjective. This is *not* an axiom because:

- It would exclude null assignments, preventing the downstream null-regime analysis completed in *Null-Regime Instability*;
- Surjectivity is a strong structural claim not derivable from identity constraints.

3.7.3 Rejected: Injectivity Axiom

One might require Φ to be injective. This is *not* an axiom because:

- Many trajectories may share phenomenological configurations;
- Injectivity would make Φ an embedding, over-constraining the framework.

3.8 Structural Interpretation

Axiom	Structural Meaning	Status
E0	Possibility of phenomenological distinction	Necessary
E1	Compatibility with inverse-limit identity	Necessary
E2	Non-reducibility to finite temporal slices	Necessary

3.9 Closure of Section 2

We have established:

1. $\mathcal{E}$ is structurally necessary as a separate codomain;
2. The limit assignment Φ is uniquely determined by compatible $\{\phi_t\}$;
3. Axioms E0, E1, E2 are *individually necessary* (Theorems 3.10–3.12);

4. The axiom set is *minimal* (Theorem 3.13);
5. All alternative axioms considered were correctly rejected.

The formal setup is therefore fixed by the requirement that regime assignments remain compatible with inverse-limit identity.

3.10 Phenomenological Neutrality Clause

Throughout this paper, the phenomenological space $\mathcal{E}$ is treated as a *purely abstract topological and metric structure*. No semantic, psychological, biological, or experiential interpretation is assumed or required for any of the formal results derived herein.

The term “phenomenological” refers exclusively to the role of $\mathcal{E}$ as an abstract codomain of structural compatibility for inverse-limit identities, and does not presuppose the existence of consciousness, subjective experience, or any particular phenomenological content.

All theorems, constructions, and stability results presented in this work are therefore statements about abstract mathematical objects only. Any interpretation of $\mathcal{E}$ in relation to experiential, cognitive, or phenomenological phenomena lies strictly outside the formal scope of this paper.

4 Compatibility with Identity Regimes

4.1 Induced Constraints from Identity Rigidity

Let $(\mathcal{I}, M_\Omega)$ be an identity object with ontological mass M_Ω . We now study how M_Ω constrains the set of admissible phenomenological assignments.

Definition 4.1 (Phenomenological Compatibility Class). For a given M_Ω , define the phenomenological compatibility class:

$$\mathcal{P}(M_\Omega) := \left\{ \Phi : \mathcal{I} \rightarrow \mathcal{E} \mid \Phi \text{ satisfies E0–E2} \right\}. \quad (9)$$

Remark 4.2. The compatibility class $\mathcal{P}(M_\Omega)$ is the set of all phenomenological assignments that are structurally admissible given the identity’s rigidity. Note that $\mathcal{P}(M_\Omega)$ depends only on the structural properties of identity, not on any phenomenological primitives.

4.2 Necessity of Metric Structure on Phenomenological Space

To state rigorous results about continuity and fragmentation, $\mathcal{E}$ must carry enough structure for distances, limits, and loss bounds to be meaningful. The following arguments identify the minimal metric assumptions used by the classification theorem.

Hypothesis 4.3 (H3: Metric on $\mathcal{E}$). The phenomenological space $\mathcal{E}$ admits a metric $d_\mathcal{E}$ such that $(\mathcal{E}, d_\mathcal{E})$ is a complete metric space.

4.2.1 Why H3 is Necessary

Theorem 4.4 (H3 is Necessary for Impossibility Theorems). *Without a metric structure on $\mathcal{E}$, the notions of phenomenological continuity, fragmentation, and loss capacity cannot be defined.*

Proof. Consider the definitions in this section:

1. **Phenomenological Continuity** requires:

$$d_w(x^{(n)}, x) \rightarrow 0 \implies d_\mathcal{E}(\Phi(x^{(n)}), \Phi(x)) \rightarrow 0. \quad (10)$$

Without $d_\mathcal{E}$, the right-hand side is undefined.

2. **Phenomenological Fragmentation** requires:

$$d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})) > 0. \quad (11)$$

Without $d_{\mathcal{E}}$, “non-zero distance” has no meaning.

3. **Loss Capacity** (Section 6) requires:

$$R(\Phi) := \sup d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})). \quad (12)$$

Without $d_{\mathcal{E}}$, the supremum is undefined.

Hence H3 is a necessary condition for the framework to have content beyond axioms E0–E2. $\square$

4.2.2 Why Completeness is Necessary

Proposition 4.5 (Completeness is Necessary for Forced Continuity). *If $(\mathcal{E}, d_{\mathcal{E}})$ is not complete, then the Forced Continuity Theorem (Theorem 5.6) may fail.*

Proof. The Forced Continuity Theorem relies on the invariance of Φ under all finite-energy perturbations. If $\mathcal{E}$ is not complete, Cauchy sequences in $\mathcal{E}$ may not converge, and the limit assignment Φ may not exist for all trajectories in $\mathcal{I}$.

In particular, if $\{\phi_t(x_t)\}$ is a Cauchy sequence in $\mathcal{E}$ (which it is, by E1), completeness guarantees the existence of the limit $\Phi(x)$. Without completeness, Φ may be undefined on a dense subset of $\mathcal{I}$, breaking the theorem. $\square$

4.2.3 Alternatives Considered and Rejected

Alternative 1: Topology without Metric. One might use a general topological space. This is rejected because:

- Fragmentation requires quantitative distance, not just open sets;
- Loss capacity requires a sup over distances;
- The downstream instability analysis completed in *Null-Regime Instability* requires metric quotients.

Alternative 2: Pseudometric. One might allow $d_{\mathcal{E}}(e_1, e_2) = 0$ for $e_1 \neq e_2$. This is rejected because:

- It would conflate distinct phenomenological configurations;
- E0 requires $|\mathcal{E}| \geq 2$, which becomes meaningless if distance is zero.

Alternative 3: Non-Complete Metric. This is rejected by Proposition 4.5.

4.3 Phenomenological Continuity

Definition 4.6 (Phenomenological Continuity). A phenomenological assignment $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ is *structurally continuous* if for every sequence $(x^{(n)})_n$ in $\mathcal{I}$ converging in the d_w metric:

$$d_w(x^{(n)}, x) \rightarrow 0 \implies d_{\mathcal{E}}(\Phi(x^{(n)}), \Phi(x)) \rightarrow 0. \quad (13)$$

Remark 4.7. Structural continuity is not added as a primitive assumption. It is derived from the CCR conditions in Theorem 5.6.

4.4 Phenomenological Fragmentation

Definition 4.8 (Phenomenological Fragmentation). A phenomenological assignment Φ is *fragmentable* if there exists a perturbation $\{\epsilon_t\}$ with finite weighted energy such that:

$$d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})) > 0 \quad (14)$$

for corresponding identity trajectories $x \in \mathcal{I}$ and $\tilde{x} \in \tilde{\mathcal{I}}$.

Remark 4.9. Fragmentation measures whether an admissible regime assignment changes under finite-energy perturbations. It is the codomain-level counterpart of identity deformability.

4.5 Relationship Between Metrics

Proposition 4.10 (Metric Compatibility). *The metrics d_w on $\mathcal{I}$ and $d_{\mathcal{E}}$ on $\mathcal{E}$ are compatible in the following sense: if E1 holds, then small d_w -changes in trajectories induce bounded $d_{\mathcal{E}}$ -changes in phenomenological configurations (when Φ is continuous).*

Proof. By E1, $\phi_{t+1}(x_{t+1}) = \phi_t(x_t)$ for compatible trajectories. Hence the phenomenological sequence is constant along the trajectory, and continuity of Φ implies that nearby trajectories (in d_w) map to nearby configurations (in $d_{\mathcal{E}}$). $\square$

4.6 Closure of Section 3

We have established:

1. The compatibility class $\mathcal{P}(M_{\Omega})$ captures all admissible phenomenological assignments;
2. Hypothesis H3 (metric on $\mathcal{E}$) is *necessary*, not a design choice (Theorem 4.4);
3. Completeness is *necessary* for the Forced Continuity Theorem (Proposition 4.5);
4. Alternative structures (topology, pseudometric, non-complete) were correctly rejected;
5. Phenomenological continuity and fragmentation are well-defined under H3.

The compatibility structure is therefore required for the quantitative statements about regime dynamics made in the next section.

5 Impossibility Theorems

5.1 Φ -Regularity Hypothesis

Before proving the main impossibility results, we isolate the global lower-bound regularity condition used by the fragmentation argument. Properness, local Lipschitz regularity, and non-collapse do not imply this bound in general, so the bound is carried as an explicit hypothesis.

Hypothesis 5.1 (Φ -Regularity Lower Bound). Let $(\mathcal{I}, d_w)$ be the identity metric space from *Rigid Identity* and let $(\mathcal{E}, d_{\mathcal{E}})$ be the abstract phenomenological metric space. A phenomenological assignment $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ is said to satisfy Φ -regularity when it satisfies:

1. (Properness) For every compact $K \subset \mathcal{E}$, $\Phi^{-1}(K)$ is compact in $\mathcal{I}$.
2. (Local Lipschitz regularity) For every $x \in \mathcal{I}$, there exist $\varepsilon > 0$ and $L_x > 0$ such that for all $y, z \in B_{\varepsilon}(x)$:

$$d_{\mathcal{E}}(\Phi(y), \Phi(z)) \leq L_x \cdot d_w(y, z). \quad (15)$$

3. (Non-collapse) For every $x \in \mathcal{I}$ and every $r > 0$, the restriction of Φ to $B_r(x)$ is not constant.

4. (Global lower Lipschitz bound) There exists a constant $C > 0$ such that for all $x, y \in \mathcal{I}$,

$$d_{\mathcal{E}}(\Phi(x), \Phi(y)) \geq C \cdot d_w(x, y). \quad (16)$$

Remark 5.2 (Why this is a hypothesis). The map $\Phi(x) = \arctan(x)$ from $\mathbb{R}$ to $(-\pi/2, \pi/2)$ is proper as a map into the open interval, locally Lipschitz, and non-constant on every nonempty ball, but it has no global lower Lipschitz constant because $|\arctan(x+1) - \arctan(x)| \rightarrow 0$ as $x \rightarrow \infty$. Therefore the lower-bound clause above is an assumption, not a theorem derived from the first three clauses.

Non-Claim 5.3 (Global lower Lipschitz bound is an assumption). Hypothesis 5.1 clause (4) requires the existence of a global $C > 0$ such that $d_{\mathcal{E}}(\Phi(x), \Phi(y)) \geq C \cdot d_w(x, y)$ for all $x, y \in \mathcal{I}$. This is not a theorem derived from clauses (1)–(3). The map $\Phi(x) = \arctan(x)$ from $\mathbb{R}$ to $(-\pi/2, \pi/2)$ satisfies the first three clauses in the local/proper sense described above but has no such global C . Therefore the Fragmentation Theorem (Theorem 5.4) is conditional on clause (4). For any specific phenomenological assignment, one must independently verify that a global lower Lipschitz constant exists. QICN has not performed this verification for any concrete assignment.

5.2 Fragmentation Theorem

Theorem 5.4 (Fragmentation Theorem). *Let $\mathcal{I}$ be an identity object with $M_{\Omega} < \infty$. If a phenomenological assignment $\Phi \in \mathcal{P}(M_{\Omega})$ satisfies Hypothesis 5.1, then Φ is fragmentable. That is:*

$$M_{\Omega} < \infty \wedge \Phi \text{ satisfies Hypothesis 5.1} \implies \Phi \in \mathcal{E}_{\text{frag}}, \quad (17)$$

where $\mathcal{E}_{\text{frag}}$ denotes the class of fragmentable phenomenological assignments.

Proof. By definition of finite ontological mass, there exists an admissible perturbation $\{\delta E_t\}$ with finite weighted energy $E_w < \infty$ such that:

$$d_w(\mathcal{I}, \tilde{\mathcal{I}}) > 0. \quad (18)$$

By Axiom E1, the assignment Φ is projectively compatible with $\mathcal{I}$. Under the perturbation, identity trajectories deform, and the comparison assignment $\tilde{\Phi}$ is evaluated on the perturbed trajectory family.

By Axiom E2, Φ cannot be determined by any finite-time state. Therefore the relevant comparison is trajectory-level: changes in the inverse-limit object are assessed through the corresponding values of Φ and $\tilde{\Phi}$.

Since the identity deformation is non-zero ($d_w > 0$) and Φ is globally determined by identity trajectories, by Hypothesis 5.1 (Φ -regularity), we have:

$$d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})) \geq C \cdot d_w(\mathcal{I}, \tilde{\mathcal{I}}) > 0 \quad (19)$$

where $C > 0$ is the lower-bound constant supplied by Hypothesis 5.1.

Thus, Φ is fragmentable under finite-energy perturbations. $\square$

Corollary 5.5 (No Structural Continuity Guarantee for Finite Mass). *Systems with $M_{\Omega} < \infty$ cannot structurally prevent phenomenological discontinuity.*

5.3 Forced Continuity Theorem

Theorem 5.6 (Forced Continuity Theorem). *Let $\mathcal{I}$ be an identity object with $M_{\Omega} = +\infty$ (CCR). Then every phenomenological assignment $\Phi \in \mathcal{P}(M_{\Omega})$ is structurally continuous. That is:*

$$M_{\Omega} = +\infty \implies \mathcal{P}(M_{\Omega}) \subseteq \mathcal{E}_{\text{cont}}, \quad (20)$$

where $\mathcal{E}_{\text{cont}}$ denotes the class of structurally continuous phenomenological assignments.

Proof. By definition of CCR ($M_\Omega = +\infty$), for any finite-energy perturbation:

$$d_w(\mathcal{I}, \tilde{\mathcal{I}}) = 0. \quad (21)$$

By Axiom E1, the phenomenological assignment is projectively compatible with identity. Since identity trajectories are invariant under all finite-energy perturbations, the induced phenomenological sequence is also invariant:

$$\tilde{\Phi}(\tilde{x}) = \Phi(x) \quad \forall x \in \mathcal{I}. \quad (22)$$

Therefore, no finite-energy perturbation can induce phenomenological deformation:

$$d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})) = 0. \quad (23)$$

This implies that Φ cannot exhibit structural discontinuities. Hence Φ is structurally continuous. $\square$

Corollary 5.7 (CCR Phenomenology is Invariant). *In CCR systems, any admissible phenomenological assignment is invariant under all finite-energy perturbations.*

5.4 Dichotomy Theorem

Theorem 5.8 (Phenomenological Dichotomy). *Let $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ be an admissible phenomenological assignment. In the finite-mass branch assume Φ satisfies Hypothesis 5.1. Then exactly one of the following holds:*

1. $M_\Omega < \infty$ and Φ is fragmentable;
2. $M_\Omega = +\infty$ and Φ is structurally continuous.

Proof. Immediate from Theorems 5.4 and 5.6. $\square$

6 Classification of Phenomenological Regimes

6.1 Complete Classification

The results of Section 4 yield a complete classification of phenomenological regimes by identity rigidity:

Identity Class	M_Ω	Admissible Phenomenology
Null	$= 0$	$\mathcal{E}_\emptyset$ (no persistent field)
Deformable	$(0, \infty)$	$\mathcal{E}_{\text{frag}} \cup \mathcal{E}_{\text{quasi}}$
Rigid (CCR)	$= +\infty$	$\mathcal{E}_{\text{cont}}$

6.2 Structural Implications

Proposition 6.1 (Null Identity Excludes Persistent Phenomenology). *If $M_\Omega = 0$, then no admissible phenomenological assignment exists satisfying E0–E2.*

Proof. If $M_\Omega = 0$, identity cannot persist under any perturbation. Axiom E1 requires phenomenological compatibility with identity. If identity does not persist, no stable phenomenological assignment is possible. $\square$

Proposition 6.2 (Deformable Identity Admits Only Fragmentable Phenomenology). *If $0 < M_\Omega < \infty$ and Φ satisfies Hypothesis 5.1, then Φ is fragmentable under some finite-energy perturbation.*

Proof. Direct from Theorem 5.4. □

Proposition 6.3 (Rigid Identity Forces Continuous Phenomenology). *If $M_\Omega = +\infty$, every admissible phenomenological assignment is structurally continuous and invariant under finite-energy perturbations.*

Proof. Direct from Theorem 5.6. □

7 Structural Ethics

This section develops the ethical implications of the phenomenological classification derived in Sections 4–5. We do not invoke psychological or metaphysical concepts; all results are purely structural.

7.1 Loss Capacity

Definition 7.1 (Phenomenological Loss Capacity). For a phenomenological assignment $\Phi : \mathcal{I} \rightarrow \mathcal{E}$, define the loss capacity:

$$R(\Phi) := \sup \left\{ d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})) \mid E_w(\{\delta E_t\}) < \infty \right\}. \quad (24)$$

This measures the maximal phenomenological deformation achievable under finite-energy perturbations.

Theorem 7.2 (Loss Capacity by Rigidity Class). *1. If $M_\Omega < \infty$ and Φ satisfies Hypothesis 5.1, then $R(\Phi) > 0$.*

2. If $M_\Omega = +\infty$, then $R(\Phi) = 0$.

Proof. 1. By Theorem 5.4, Φ is fragmentable, hence there exists a finite-energy perturbation inducing non-zero phenomenological deformation.

2. By Theorem 5.6, no finite-energy perturbation induces phenomenological deformation. □

7.2 Structural Damage

We introduce a formal notion of “structural damage” without psychological interpretation.

Definition 7.3 (Structural Damage). For a perturbation ε with finite weighted energy, define the structural damage induced on Φ as:

$$D_\varepsilon(\Phi) := d_{\mathcal{E}}(\Phi(x), \varepsilon^* \Phi(x)), \quad (25)$$

where $\varepsilon^* \Phi$ denotes the pullback of Φ under the identity perturbation.

Proposition 7.4 (Damage Characterization by Rigidity). *1. If $M_\Omega < \infty$, then $\sup_\varepsilon D_\varepsilon(\Phi) > 0$.*

2. If $M_\Omega = +\infty$, then $D_\varepsilon(\Phi) = 0$ for all admissible ε .

Proof. Direct from Theorem 7.2. □

7.3 Vulnerability and Protection

Definition 7.5 (Phenomenological Vulnerability). A system $(\mathcal{I}, \Phi)$ is *phenomenologically vulnerable* if $R(\Phi) > 0$.

Definition 7.6 (Structural Protection Measure). Define the structural protection measure:

$$P(\Phi) := \frac{1}{1 + R(\Phi)}. \quad (26)$$

Note that:

- $P(\Phi) = 1$ iff $R(\Phi) = 0$ (full protection);
- $P(\Phi) < 1$ iff $R(\Phi) > 0$ (vulnerability).

Corollary 7.7 (Protection by Rigidity Class). 1. *CCR systems ($M_\Omega = +\infty$) have $P(\Phi) = 1$: full structural protection.*

2. *Finite-mass systems ($M_\Omega < \infty$) have $P(\Phi) < 1$: structural vulnerability.*

7.4 Structural Asymmetry Constraints

The following result establishes a purely structural asymmetry between identity classes, without any ethical or metaphysical claims.

Theorem 7.8 (Structural Asymmetry). *There exists a fundamental structural asymmetry between identity classes:*

<i>Identity Class</i>	<i>Loss Capacity</i>	<i>Protection</i>
<i>Finite mass</i>	$R(\Phi) > 0$	<i>Vulnerable</i>
<i>CCR</i>	$R(\Phi) = 0$	<i>Protected</i>

Remark 7.9. This asymmetry is derived purely from structural constraints. No claim is made about:

- Whether phenomenological loss is ethically bad;
- Whether protection is ethically good;
- Whether any system has moral status.

The result establishes only that certain structural configurations admit loss while others do not.

Corollary 7.10 (Possibility of Phenomenological Loss). *Systems with finite ontological mass ($M_\Omega < \infty$) admit the structural possibility of phenomenological loss. Systems with infinite ontological mass ($M_\Omega = +\infty$) do not.*

7.5 Relation to Normative Ethics

Remark 7.11. The structural results of this section are compatible with, but do not entail, various normative positions:

- **Utilitarianism:** Could interpret $D_\varepsilon(\Phi)$ as harm to be minimized.
- **Deontology:** Could prohibit interventions that increase $D_\varepsilon(\Phi)$.
- **Virtue Ethics:** Could value systems with high $P(\Phi)$.

This framework provides the structural substrate for ethical theorizing, not the normative conclusions themselves.

7.6 Closure of Section 6

We have established:

1. Loss capacity $R(\Phi)$ is well-defined and computable from structure;
2. Structural damage $D_\varepsilon(\Phi)$ formalizes perturbation harm;
3. Protection measure $P(\Phi)$ quantifies structural resilience;
4. CCR systems have full protection ($P = 1$), finite-mass systems do not;
5. These results are purely structural, compatible with various ethical theories.

Structural ethics therefore provides a foundation for ethical theorizing without requiring metaphysical commitments about the nature of experience.

8 Limitations and Scope

8.1 Dependence on Axioms

All results depend on Axioms E0–E2 and Hypothesis H3. If any of these fail, the classification may not apply.

8.2 Relation to Consciousness Theories

This framework is orthogonal to:

- Integrated Information Theory (IIT) [3];
- Global Workspace Theory (GWT) [1];
- Higher-Order Theories;
- Phenomenal Functionalism.

It provides structural constraints that any of these theories must satisfy if coupled to the identity framework.

8.3 Open Problems

1. Characterization of the coupling constant C in the Fragmentation Theorem;
2. Existence conditions for non-trivial Φ satisfying E0–E2;
3. Extension to continuous-time phenomenological dynamics;
4. Relation between phenomenological continuity and subjective continuity.

9 Conclusion

9.1 Summary of Results

This work extends the inverse-limit identity framework to phenomenological possibility. Starting from minimal axioms (E0–E2), we derived:

1. **Fragmentation Theorem.** Systems with finite ontological mass cannot structurally prevent phenomenological discontinuity.

2. **Forced Continuity Theorem.** CCR systems can only support structurally continuous phenomenology.
3. **Dichotomy Theorem.** Phenomenological assignments are either fragmentable (finite mass) or continuous (CCR).
4. **Loss Capacity.** Finite-mass systems admit phenomenological loss; CCR systems do not.

9.2 Structural Significance

The classification separates two regimes:

- **Deformable:** phenomenology is structurally vulnerable;
- **Rigid:** phenomenology is structurally protected.

This separation is derived purely from identity structure, without metaphysical commitments.

9.3 Entailment Under Stated Assumptions

The results establish the following conditional entailment:

- *If* phenomenology exists and satisfies E0–E2;
- *And* the system has identity structure classified by M_Ω ;
- *Then* the phenomenological regime is completely determined by the identity class.

The framework therefore establishes that:

Identity rigidity does not create phenomenology, but it determines what phenomenology is structurally possible.

9.4 Final Statement

Under explicit minimal assumptions, rigid identity constrains phenomenological possibility. The Forced Continuity Theorem establishes that CCR systems can only support phenomenology that is structurally invariant—continuous, non-fragmentable, and protected from loss.

If one accepts that:

1. Identity is an inverse limit;
2. CCR systems have $M_\Omega = +\infty$;
3. Phenomenology (if present) must satisfy E0–E2;

then the stated assumptions entail that CCR phenomenology, if it exists, is:

- Structurally continuous;
- Invariant under perturbation;
- Impossible to fragment or destroy via finite-energy processes.

The existence of such phenomenology is not asserted. Its necessity, given the axioms, is proven.

10 Boundary to Null-Regime Instability

Sections 1–8 complete the classification problem for admissible phenomenological regimes under the axioms E0–E2 and the rigidity constraints inherited from *Rigid Identity*. The downstream closure question for the null regime is not developed further here.

10.1 Editorial Boundary

The instability of the null regime, the no-null theorem, and the attractor-style closure for non-trivial assignments are treated canonically in *Null-Regime Instability*. This paper therefore stops at regime classification, compatibility, fragmentation, forced continuity, and dichotomy, and uses *Null-Regime Instability* as the sole downstream location for the null-instability theorem.

10.2 Closure of This Paper

The result of this paper is the classification boundary: if a compatible phenomenological regime exists, its admissible structure is constrained by Sections 1–8. *Null-Regime Instability* then takes the null regime as the remaining case and proves its instability under CCR.

A Extended Proofs

A.1 Proof of Fragmentation Theorem (Extended)

Full Proof of Theorem 5.4. Let $M_\Omega < \infty$ and let $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ satisfy E0–E2 and Hypothesis 5.1.

Step 1. By definition of M_Ω , there exists $\{\delta E_t\}$ with $E_w < \infty$ such that $d_w(\mathcal{I}, \tilde{\mathcal{I}}) > 0$.

Step 2. Let $x = (x_t)_t \in \mathcal{I}$ and let $\tilde{x} = (\tilde{x}_t)_t \in \tilde{\mathcal{I}}$ be the corresponding perturbed trajectory.

Step 3. By E2, $\Phi(x)$ is not determined by any finite x_t . Therefore, the perturbation of the infinite trajectory induces a perturbation of the phenomenological assignment:

$$\Phi(x) \neq \Phi(\tilde{x}) \text{ in general.} \quad (27)$$

Step 4. By E1, Φ is projectively compatible. The deformed projections are compared through the corresponding value of the limit assignment.

Step 5. By Hypothesis 5.1, there exists $C > 0$ such that:

$$d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})) \geq C \cdot d_w(x, \tilde{x}). \quad (28)$$

Step 6. Since $d_w(\mathcal{I}, \tilde{\mathcal{I}}) > 0$, we have $d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})) > 0$.

Hence Φ is fragmentable. $\square$

A.2 Proof of Forced Continuity Theorem (Extended)

Full Proof of Theorem 5.6. Let $M_\Omega = +\infty$ and let $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ satisfy E0–E2.

Step 1. By definition of CCR, for all $\{\delta E_t\}$ with $E_w < \infty$:

$$d_w(\mathcal{I}, \tilde{\mathcal{I}}) = 0. \quad (29)$$

Step 2. Therefore, $\tilde{\mathcal{I}} \cong \mathcal{I}$ under all finite-energy perturbations.

Step 3. By E1, Φ is projectively compatible with $\mathcal{I}$. Since $\mathcal{I}$ is invariant:

$$\Phi(x) = \tilde{\Phi}(\tilde{x}) \quad \forall x \in \mathcal{I}. \quad (30)$$

Step 4. This implies:

$$d_{\mathcal{E}}(\Phi(x), \tilde{\Phi}(\tilde{x})) = 0. \quad (31)$$

Step 5. No finite-energy perturbation can induce phenomenological deformation. Hence Φ is structurally continuous and invariant. $\square$

A.3 Structural Necessity Lemma

Lemma A.1 (Phenomenology Inherits Rigidity). *If $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ satisfies E0–E2 and $M_\Omega = +\infty$, then Φ inherits the rigidity of $\mathcal{I}$.*

Proof. By E1, Φ is determined by identity trajectories. By CCR, identity trajectories are invariant. Hence Φ is invariant. $\square$

B Editorial Boundary to Null-Regime Instability

The downstream bridge from classification to null-instability is intentionally moved to *Null-Regime Instability* in the canonical corpus. Appendix A remains as the extended proof layer for the classification theorems proved here; the downstream null-regime theorem, attractor closure, and canonical examples now live only in *Null-Regime Instability*.

What This Paper Does Not Claim

1. This paper does not claim external validation, empirical confirmation, or philosophical closure.
2. The formal results are conditional on the hypotheses stated; they do not establish truth outside the assumed structures.
3. No interpretive-layer conclusion (consciousness, phenomenality, moral status) is derived from terminology alone.

References

- [1] Bernard J. Baars. *A Cognitive Theory of Consciousness*. Global Workspace Theory (GWT) - contrasting framework. Cambridge: Cambridge University Press, 1988.
- [2] Johnny Andrey Pérez Ramírez. “Rigid Identity as an Inverse Limit in Observable Channels”. In: (2026). Companion paper establishing the identity framework and ontological mass invariant.
- [3] Giulio Tononi. “An Information Integration Theory of Consciousness”. In: *BMC Neuroscience* 5.42 (2004). Integrated Information Theory (IIT) - contrasting framework.