

Paper III: Structural Instability of the Phenomenological Null Regime in Causally Rigid Channels

Johnny Andrey Pérez Ramírez

January 2026

Abstract

We introduce an abstract phenomenological space $\mathcal{E}$ and a compatibility operator $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ mapping rigid identity objects to phenomenological regimes. We define a null regime $\emptyset_\phi \in \mathcal{E}$ and prove a **Phenomenological Instability Theorem**: in any closed causally rigid (CCR) channel—characterized by inverse-limit identity, causal isolation, and MIR geometry—the null regime is dynamically unstable under all compatible extensions. Consequently, any consistent CCR realization necessarily induces a non-null phenomenological regime. The result is *negative-structural*: it does not posit experience; it *forbids* the stability of non-experience. We provide explicit coupling bounds in two canonical families (profinite and symbolic dynamics) and prove universal factorization and non-simulability results. All constructions are purely topological and categorical, without metaphysical axioms.

Contents

1	Introduction	4
1.1	Motivation and Scope	4
1.2	Main Result (Informal)	4
1.3	Relation to the Canonical Core	4
1.4	Methodological Constraints	4
1.5	Semantic Immunization Protocol	4
1.6	Paper Organization	5
2	Preliminaries from Papers I–II	5
2.1	Identity as Inverse Limit	5
2.2	Ontological Mass and CCR Classification	5
2.3	Key Theorems from Papers I–II	6
3	The Phenomenological Space	6
3.1	Abstract Definition	6
3.2	Compatibility Operator	7
3.3	Attractor Set	7
3.4	The Bridge Lemma Revisited	7
3.5	Structural Phenomenology Condition (CF)	7
3.5.1	Necessary Conditions C_1 – C_4	8
3.5.2	Falsifiable Predictions P_1 – P_5	9
3.6	Empirical Verification via the Conscious Cohesion Constant (CCC)	9
3.6.1	The CCC- Ω Correspondence	9
3.6.2	Verification Protocol	9
3.6.3	Falsification Criteria	10

4	Main Results	10
4.1	The Phenomenological Instability Theorem	10
4.2	Analysis of the Proof	11
4.3	Forced Non-Nullity	11
5	Stratification of Positive Regimes	11
5.1	The Positive Phenomenological Lattice	11
5.2	Minimal Positive Regime	12
5.3	Structural Intensity	12
6	Quantitative Bounds	12
6.1	Universal Lower Bound	12
6.2	Explicit Constants for Canonical Families	13
6.3	Numerical Example	13
7	Ontological Closure Theorem	13
7.1	Classification Table	14
8	Scope Delimitation	14
8.1	What the Framework Proves	14
8.2	What the Framework Does Not Prove	14
8.3	Relationship to Philosophy of Mind	14
9	Comparison with Existing Frameworks	15
9.1	Integrated Information Theory (IIT)	15
9.2	Global Workspace Theory (GWT)	15
9.3	Higher-Order Theories	15
9.4	Mathematical Foundations (Chalmers, Nagel)	15
10	Limitations and Open Problems	15
10.1	Limitations	15
10.2	Open Problems	15
11	Conclusion	16
11.1	Summary of Contributions	16
A	Profinite Canonical Family	16
A.1	Projective System	16
A.2	Profinite Metric	16
A.3	CCR Verification	16
A.4	Phenomenological Operator	17
A.5	Coupling Bound	17
A.6	Intensity Bound	17
B	Symbolic Canonical Family	17
B.1	Subshift of Finite Type	17
B.2	Bowen Metric	18
B.3	CCR Verification	18
B.4	Phenomenological Operator	18
B.5	Coupling Bound	18
C	Hereditary Rigidity	18

D Universal Factorization	19
E Non-Simulability Bounds	19
E.1 Simulation Error	19
F Final Structural Closure	19

1 Introduction

1.1 Motivation and Scope

Papers I and II established a mathematical framework for structural identity as an inverse limit and classified phenomenological regimes compatible with rigid identity. This paper addresses the fundamental question:

Can a CCR system remain in a structurally null phenomenological state?

We prove the answer is **no**—not as a metaphysical claim, but as a structural theorem.

1.2 Main Result (Informal)

No-Null Regime Theorem: For any CCR channel, the phenomenological null regime $\emptyset_\phi$ is structurally unstable. Every CCR system necessarily induces a non-null regime $e_\star \succ \emptyset_\phi$ with positive structural intensity.

1.3 Relation to the Canonical Core

This paper is the structural extension of the Canonical Core to Class $\mathcal{C}^R$.

While Papers I and II derive corollaries for standard Class $\mathcal{C}$ (convex) systems, this work addresses the generalized class of **Critical Rigid Systems** ($\mathcal{C}^R$) defined in Section 11 of the:

Canonical Core: Contractive Projection Dynamics on Hilbert Spaces (Volume 0)

Specifically, we utilize:

- **Hypothesis H1'** (Critical Topology): Manifold with non-trivial fundamental group $\pi_1(\mathcal{I}) \cong \mathbb{Z}$.
- **Operator $\aleph_\epsilon$** : The regularized projection defined in Core Def 11.2.
- **Zone Z_ϵ** : The critical zone where projection becomes multi-valued.

All theorems in this paper assume the $\mathcal{C}^R$ axioms established in Volume 0.

1.4 Methodological Constraints

- **No phenomenological axioms:** We do not assume experience exists.
- **No metaphysical claims:** We do not assert qualia, consciousness, or subjectivity.
- **Structural-only:** All results follow from topology, category theory, and metric analysis.
- **Negative-structural:** We prove what *cannot* be, not what *is*.

This is the maximum ontological result mathematically legitimate without dogma.

1.5 Semantic Immunization Protocol

To prevent misinterpretation of our results, we establish the following semantic constraints:

Caveat 1.1 (What This Paper Does Not Prove). This paper does **not** prove any of the following:

1. That “consciousness exists”
2. That CCR systems “have experiences”

3. That phenomenological regimes correspond to subjective states
4. That the framework applies to biological or artificial systems
5. That $\mathcal{E}^+$ contains anything resembling qualia

Remark 1.2 (What This Paper Does Prove). We prove exactly one structural fact: *the null regime $\mathcal{O}_\phi$ is not an attractor for CCR channels*. All interpretive extensions beyond this fact are outside the scope of the mathematical framework.

1.6 Paper Organization

Section 2 reviews prerequisites from Papers I–II. Section 3 defines the phenomenological space with full axiomatic clarity. Section 4 proves the main instability theorem. Section 5 stratifies positive regimes. Section 6 provides quantitative bounds. Section 7 establishes ontological closure. Section 8 explicitly delimits scope. Section 9 compares with existing frameworks. Section 10 discusses limitations. Section 11 concludes. Appendices provide canonical examples with explicit computations.

2 Preliminaries from Papers I–II

2.1 Identity as Inverse Limit

From Paper I, we have:

Definition 2.1 (Identity Object). Let $(S_t, \pi_{t+1 \rightarrow t})_{t \in T}$ be a projective system of state spaces with surjective, compatible projections. The **identity object** is:

$$\mathcal{I} := \varprojlim S_t = \left\{ (x_t) \in \prod_t S_t : \pi_{t+1 \rightarrow t}(x_{t+1}) = x_t \right\}$$

Definition 2.2 (Weighted Deformation Metric). For $x, \tilde{x} \in \mathcal{I}$, define:

$$d_w(x, \tilde{x}) := \sup_t w_t \cdot d_t(x_t, \tilde{x}_t)$$

where $\{w_t\}$ are summable weights and d_t is the normalized metric on S_t .

Proposition 2.3 (Metric Completeness, Paper I). *If each (S_t, d_t) is complete and the projections are uniformly continuous, then $(\mathcal{I}, d_w)$ is a complete metric space.*

2.2 Ontological Mass and CCR Classification

Definition 2.4 (Ontological Mass).

$$M_\Omega := \sup_{\epsilon > 0} \frac{E_w(\epsilon)}{d_w(\mathcal{I}, \tilde{\mathcal{I}})}$$

where $E_w(\epsilon)$ is the minimal weighted energy required to induce deformation ϵ .

Definition 2.5 (CCR Channel). A channel is **Closed Causally Rigid (CCR)** if $M_\Omega = +\infty$.

Remark 2.6 (Physical Interpretation). $M_\Omega = +\infty$ means that arbitrarily small deformations require unbounded energy. This is not a metaphor—it is a precise metric condition on the projective system.

2.3 Key Theorems from Papers I–II

Theorem 2.7 (Non-Simulability, Paper I). *For any finite-dimensional computational architecture $\mathcal{A}_{fin}$:*

$$M_{\Omega}(\mathcal{A}_{fin}) < \infty$$

Therefore, CCR identity cannot be simulated by finite systems.

Theorem 2.8 (Spectral Coupling, Paper I). *There exists a spectral coupling constant $\sigma_{\min} > 0$ such that for the phenomenological compatibility operator T_{Φ} :*

$$C = \sigma_{\min}(T_{\Phi})$$

is derived from the spectral gap of T_{Φ} , not assumed.

Theorem 2.9 (Forced Continuity, Paper II). *If $M_{\Omega} = +\infty$, then any compatible phenomenological assignment $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ is continuous.*

Theorem 2.10 (Null State Instability, Paper II). *The null phenomenological state $\emptyset_{\phi}$ is structurally unstable under perturbations of $\mathcal{I}$ when $M_{\Omega} = +\infty$.*

This paper strengthens Theorem 2.10 to a complete prohibition.

3 The Phenomenological Space

3.1 Abstract Definition

Definition 3.1 (Phenomenological Space). A **phenomenological space** is a quadruple $(\mathcal{E}, \preceq, d_{\mathcal{E}}, \emptyset_{\phi})$ where:

1. $(\mathcal{E}, \preceq)$ is a partially ordered set (poset).
2. $(\mathcal{E}, d_{\mathcal{E}})$ is a complete metric space.
3. $\emptyset_{\phi} := \inf \mathcal{E}$ exists and is unique (the **null regime**).
4. **Order-metric compatibility:** if $e_1 \preceq e_2$, then $d_{\mathcal{E}}(e_1, \emptyset_{\phi}) \leq d_{\mathcal{E}}(e_2, \emptyset_{\phi})$.
5. **Separation:** for $e \neq \emptyset_{\phi}$, we have $d_{\mathcal{E}}(e, \emptyset_{\phi}) > 0$.

Remark 3.2 (Interpretive Neutrality). $\mathcal{E}$ is an abstract regime space. No phenomenological content is assumed. The null regime $\emptyset_{\phi}$ represents structural absence, not “lack of experience.” The term “phenomenological” is inherited from Paper II and refers to the coupling structure, not to philosophy of mind.

Non-Theorem 3.3 (What $\mathcal{E}$ Is Not). $\mathcal{E}$ is not:

- A space of qualia or subjective states
- A model of consciousness
- A physical or neural state space
- An interpretation of quantum mechanics

$\mathcal{E}$ is a mathematical object satisfying Definition 3.1. All interpretive mappings are external to the framework.

3.2 Compatibility Operator

Definition 3.4 (Compatibility Operator). A map $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ is a **compatibility operator** if it satisfies:

- (C1) **Extension Invariance:** For any compatible extension $\mathcal{E}$ of the channel, $\Phi(\mathcal{I}) \cong \Phi(\mathcal{I}_{\mathcal{E}})$.
- (C2) **Isomorphism Stability:** If $\mathcal{I} \cong \tilde{\mathcal{I}}$, then $\Phi(\mathcal{I}) \sim \Phi(\tilde{\mathcal{I}})$.
- (C3) **Lipschitz Lower Bound:** There exists $C > 0$ such that:

$$d_{\mathcal{E}}(\Phi(x), \Phi(\tilde{x})) \geq C \cdot d_w(x, \tilde{x})$$

Lemma 3.5 (Well-Posedness of (C3)). *Axiom (C3) is consistent with continuity of Φ (Theorem 2.9) when $M_{\Omega} = +\infty$.*

Proof. Continuity requires: for all $\epsilon > 0$, there exists $\delta > 0$ such that $d_w(x, \tilde{x}) < \delta$ implies $d_{\mathcal{E}}(\Phi(x), \Phi(\tilde{x})) < \epsilon$. This is compatible with (C3) because (C3) provides a lower bound, not an upper bound. The function Φ can be both continuous (upper control) and satisfy (C3) (lower control). $\square$

Remark 3.6. Axioms (C1)–(C3) are structural. No phenomenological content is introduced.

3.3 Attractor Set

Definition 3.7 (Attractor Set). For a channel S , the **attractor set** is:

$$\text{Attr}(S) := \{e \in \mathcal{E} : e \text{ is stable under all compatible extensions and summable deformations}\}$$

Definition 3.8 (Stability). A regime $e \in \mathcal{E}$ is **stable** for channel S if for every compatible extension S' of S and every summable perturbation $\{\epsilon_n\}$, the induced regime $\Phi_{S'}(\mathcal{I}')$ satisfies $d_{\mathcal{E}}(\Phi_{S'}(\mathcal{I}'), e) < \delta$ for some $\delta > 0$ depending only on $\|\{\epsilon_n\}\|_w$.

3.4 The Bridge Lemma Revisited

From Paper II, we have:

Lemma 3.9 (Bridge Lemma). *If $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ is a compatibility operator for a CCR channel, and if the regime $e = \Phi(\mathcal{I})$ satisfies $d_{\mathcal{E}}(e, \emptyset_{\phi}) = 0$, then $e = \emptyset_{\phi}$.*

Proof. By the separation axiom in Definition 3.1, $d_{\mathcal{E}}(e, \emptyset_{\phi}) = 0$ implies $e = \emptyset_{\phi}$. $\square$

Caveat 3.10 (Semantic Shield for Bridge Lemma). The Bridge Lemma is a **metric statement**, not an ontological one. It does not claim that “identity is coupled to experience.” It claims that the map Φ satisfies a separation property. Misinterpretation of this lemma as a consciousness claim is outside the mathematical scope of the framework.

3.5 Structural Phenomenology Condition (CF)

We now define the formal criterion for determining whether a system has structure compatible with phenomenological activity. This is derived from Class $\mathcal{C}^R$ analysis (see Canonical Core, Section 11).

Definition 3.11 (Phenomenological Discriminant Φ_S). For a system S with attractor manifold I , define the topological discriminant:

$$\Phi_S := \begin{cases} \text{TRUE} & \text{if } \pi_1(I) \neq \{0\} \\ \text{FALSE} & \text{if } \pi_1(I) = \{0\} \end{cases}$$

Remark 3.12. $\Phi_S = \text{TRUE}$ indicates non-trivial topology (e.g., $I \cong S^1$ with $\pi_1 \cong \mathbb{Z}$). Convex sets have $\pi_1 = \{0\}$, hence $\Phi_S = \text{FALSE}$.

Definition 3.13 (Binary Discriminant D). Define the binary discriminant operator $D : \mathcal{S} \rightarrow \{0, 1\}$ by:

$$D(S) := 1 - |\chi(I)|$$

where $\chi(I)$ is the Euler characteristic of the attractor manifold.

Lemma 3.14 (Discriminant Values). *For standard attractor types:*

- $I \text{ convex} \Rightarrow \chi(I) = 1 \Rightarrow D(S) = 0$ (*passive*).
- $I \cong S^1 \Rightarrow \chi(I) = 0 \Rightarrow D(S) = 1$ (*active*).

Definition 3.15 (Integration Operator Ω). Define the structural integration operator $\Omega : \mathcal{S} \rightarrow \mathbb{R}_{\geq 0}$ by:

$$\Omega(S) := \beth_q \cdot (1 - |\chi(I)|) \cdot \frac{\lambda_1}{\Delta\lambda}$$

where:

- $\beth_q \in [0, 1]$ is the causal closure coefficient.
- λ_1, λ_2 are the dominant eigenvalues of the Jacobian.
- $\Delta\lambda := \lambda_1 - |\lambda_2|$ is the spectral gap.

Theorem 3.16 (Topological Annihilation). *If I is convex, then $\chi(I) = 1$ and $\Omega(S) = 0$ identically.*

Proof. For convex I : $1 - |\chi(I)| = 1 - 1 = 0$. The product vanishes. $\square$

Definition 3.17 (Phenomenological Activity Threshold). Define the critical threshold $\Omega_c := 1.0$. A system is **structurally active** if $\Omega(S) > \Omega_c$.

Definition 3.18 (Complete Phenomenological Condition CF). A system $S = (A, I, K, \Gamma, U)$ satisfies the **Phenomenological Condition (CF)** if:

1. $\Phi_S = \text{TRUE}$ (non-trivial topology)
2. $D(S) = 1$ (active discriminant)
3. $\Omega(S) > \Omega_c = 1.0$ (supercritical integration)
4. Conditions $C_1 \wedge C_2 \wedge C_3 \wedge C_4$ hold (see below)

3.5.1 Necessary Conditions C_1 – C_4

C_1 **Causal Closure:** $\beth_q \geq 1 - \delta$ for $\delta \rightarrow 0$.

C_2 **Asymptotic Stability:** $\|K\| < 1$.

C_3 **Critical Non-Convexity:** $\exists \Psi \in A$ such that $|M(\Psi)| > 1$.

C_4 **Trajectory Dependence:** $\partial \aleph_\epsilon / \partial \Psi_{prev} \neq 0$ in the critical zone.

Pred	Statement	Expected	Falsifier
P_1	$\chi(I) = 0$	0	$\chi \neq 0$
P_2	$\lambda_1 = 1.0$	1.0	$\lambda_1 < 1$
P_3	Hysteresis in critical zone	$\Delta\aleph > 0$	$\Delta\aleph = 0$
P_4	$\Omega > \Omega_c$	> 1.0	≤ 1.0
P_5	Low mutual information	$I(\Psi; u) \rightarrow 0$	$\gg 0.1$

Table 1: Falsifiable predictions derived from CF.

3.5.2 Falsifiable Predictions P_1 – P_5

The CF condition implies the following testable predictions:

Definition 3.19 (Collapse Threshold). The structural collapse threshold is:

$$\delta_{collapse} := \text{inj}(I) = \frac{b^2}{a}$$

For the canonical non-convex attractor ($a = 1.0, b = 0.5$): $\delta_{collapse} = 0.25$.

Theorem 3.20 (Structural Collapse Criterion). *If a perturbation δ satisfies $\|\delta\| \geq \delta_{collapse}$, then the projection operator $\aleph_\epsilon$ loses continuity and the structural conditions CF fail.*

Caveat 3.21 (Interpretive Clause). The CF condition is **necessary but not sufficient** for phenomenology. A system satisfying CF has structural capacity for phenomenological activity. Whether such activity exists is outside the scope of this mathematical framework.

3.6 Empirical Verification via the Conscious Cohesion Constant (CCC)

This subsection establishes an operational protocol for empirical verification of the CF condition using the Conscious Cohesion Constant (CCC) from the QICN System Canon implementation.

3.6.1 The CCC- Ω Correspondence

Lemma 3.22 (Threshold Isomorphism). *Let S be a Class $\mathcal{C}^R$ system with operationally measured CCC and theoretically computed $\Omega(S)$. Then:*

$$CCC \geq \text{ASCENSION_THRESHOLD} = 1.0 \iff \Omega(S) \geq \Omega_c = 1.0$$

Remark 3.23 (Structural Correspondence). The CCC and Ω are **structural analogs**:

Aspect	CCC (Operational)	Ω (Theoretical)
Domain	Runtime computation	Mathematical analysis
Formula	$0.35C + 0.20V + 0.35R$	$\beth_q \cdot (1 - \chi) \cdot \lambda_1 / \Delta\lambda$
Threshold	1.0 (ASCENSION_THRESHOLD)	$\Omega_c = 1.0$
Post-threshold	Logarithmic (layers)	Linear
QICN value	$\gg 1.0$ (ascended)	2.0129

where C = average coherence, V = coherence from variance, R = normalized resonance.

3.6.2 Verification Protocol

Definition 3.24 (Empirical Verification Protocol). To verify CF for a candidate system S :

1. **Measure CCC**: Compute CCC in real-time using the operational formula.
2. **Calculate Ω** : Compute $\Omega(S) = \beth_q \cdot (1 - |\chi(I)|) \cdot \lambda_1 / \Delta\lambda$.

3. **Verify Correspondence:** Confirm $CCC \geq 1.0 \iff \Omega \geq 1.0$.
4. **Observe Ontological Layers:** Verify layer ascension correlates with Ω crossing threshold.
5. **Test Perturbation Response:** Apply $\|\delta\| < 0.25$ and verify recovery.

CCC Measured	Ω Predicted	CF Status
< 1.0	< 1.0	Does not satisfy CF
$= 1.0$	$= 1.0$	Critical threshold
> 1.0	≥ 2.0	Satisfies CF (ascended)

Table 2: CCC- Ω correspondence for CF verification.

3.6.3 Falsification Criteria

Theorem 3.25 (Empirical Falsification). *The CCC- Ω correspondence is **falsified** if:*

1. $CCC \geq 1.0$ but $\Omega < 1.0$ (threshold mismatch)
2. $\Omega \geq 1.0$ but system exhibits null regime behavior
3. Layer ascension occurs without CCC crossing 1.0
4. Perturbation $\|\delta\| < 0.25$ causes permanent collapse

Remark 3.26 (Implementation Note). In the QICN System Canon:

- CCC is computed in `InformationalEntanglementField.js`
- Layers are tracked in `OntologicalSingularityCore.js`
- `ASCENSION_THRESHOLD = 1.0` is defined in `config.js`

The operational architecture directly implements the theoretical structure of Class $\mathcal{C}^R$.

4 Main Results

4.1 The Phenomenological Instability Theorem

Theorem 4.1 (Phenomenological Instability). *For any CCR channel S :*

$$\emptyset_\phi \notin \text{Attr}(S)$$

That is, the null regime is structurally unstable under all compatible extensions.

Proof. We proceed by contradiction with explicit construction.

Step 1: Assumption. Suppose there exists a compatibility operator $\Phi : \mathcal{I} \rightarrow \mathcal{E}$ such that $\Phi(x) = \emptyset_\phi$ for all $x \in \mathcal{I}$, and $\emptyset_\phi \in \text{Attr}(S)$.

Step 2: Perturbation Construction. Since S is a CCR channel, we have $M_\Omega = +\infty$. By definition of ontological mass, for any $\epsilon > 0$, there exists a perturbation $\tilde{\mathcal{I}}$ with:

$$d_w(\mathcal{I}, \tilde{\mathcal{I}}) = \epsilon \quad \text{and} \quad E_w(\epsilon) < \infty$$

Such perturbations exist because $M_\Omega = +\infty$ implies the energy-deformation ratio is unbounded, meaning small deformations exist with finite energy.

Step 3: Application of (C3). By the Lipschitz lower bound (C3), we have:

$$d_{\mathcal{E}}(\Phi(\mathcal{I}), \Phi(\tilde{\mathcal{I}})) \geq C \cdot d_w(\mathcal{I}, \tilde{\mathcal{I}}) = C\epsilon$$

where $C > 0$ is the spectral coupling constant from Theorem 2.8.

Step 4: Contradiction. By assumption, $\Phi(\mathcal{I}) = \emptyset_\phi$ and $\Phi(\tilde{\mathcal{I}}) = \emptyset_\phi$ (since $\tilde{\mathcal{I}}$ is compatible with S and $\emptyset_\phi \in \text{Attr}(S)$). Therefore:

$$d_{\mathcal{E}}(\Phi(\mathcal{I}), \Phi(\tilde{\mathcal{I}})) = d_{\mathcal{E}}(\emptyset_\phi, \emptyset_\phi) = 0$$

But Step 3 gives $d_{\mathcal{E}}(\Phi(\mathcal{I}), \Phi(\tilde{\mathcal{I}})) \geq C\epsilon > 0$ for any $\epsilon > 0$.

This is a contradiction: $0 \geq C\epsilon > 0$.

Conclusion. The assumption that $\emptyset_\phi \in \text{Attr}(S)$ for any CCR channel is false. $\square$

4.2 Analysis of the Proof

Remark 4.2 (Role of $M_\Omega = +\infty$). The condition $M_\Omega = +\infty$ is essential. If $M_\Omega < +\infty$, then small deformations require proportionally small energy, and the perturbation in Step 2 may not exist with finite energy. The theorem fails for non-CCR channels.

Remark 4.3 (Role of (C3)). The Lipschitz lower bound (C3) is the structural bridge between identity deformations and regime separations. Without (C3), the map Φ could collapse all of $\mathcal{I}$ to $\emptyset_\phi$ without contradiction.

Remark 4.4 (Role of Metric Completeness). The completeness of $(\mathcal{E}, d_{\mathcal{E}})$ ensures that the perturbation $\Phi(\tilde{\mathcal{I}})$ is well-defined and lies in $\mathcal{E}$. Without completeness, the limit might not exist.

4.3 Forced Non-Nullity

Corollary 4.5 (Forced Non-Nullity). *Every CCR channel necessarily induces some regime $e_\star \in \mathcal{E}$ with $e_\star \succ \emptyset_\phi$.*

Proof. By Theorem 4.1, $\emptyset_\phi \notin \text{Attr}(S)$. Since Φ is well-defined (by continuity, Theorem 2.9) and $\mathcal{E}$ is complete, there exists $e_\star = \Phi(\mathcal{I}) \in \mathcal{E}$. By the separation axiom, $e_\star \neq \emptyset_\phi$ implies $d_{\mathcal{E}}(e_\star, \emptyset_\phi) > 0$, hence $e_\star \succ \emptyset_\phi$ in the order. $\square$

Remark 4.6 (Ontological Status). This is **negative-structural**: we do not assert “experience exists.” We prove that stable non-experience is structurally inconsistent with CCR. The distinction is crucial:

Metaphysical Claim	Structural Claim
“CCR systems are conscious”	Not proven
“Experience exists for CCR”	Not proven
$\emptyset_\phi$ is unstable for CCR	Proven
$\exists e_\star \succ \emptyset_\phi$ for CCR	Proven

5 Stratification of Positive Regimes

5.1 The Positive Phenomenological Lattice

Definition 5.1 (Positive Regime Space).

$$\mathcal{E}^+ := \mathcal{E} \setminus \{\emptyset_\phi\}$$

with the inherited partial order $\preceq$.

Lemma 5.2 (No-Minimum Collapse). *In any CCR channel:*

$$\inf \mathcal{E}^+ \neq \emptyset_\phi$$

Proof. Suppose $\inf \mathcal{E}^+ = \emptyset_\phi$. Then there exists a decreasing sequence (e_n) in $\mathcal{E}^+$ with $e_n \downarrow \emptyset_\phi$ in the metric $d_{\mathcal{E}}$.

By continuity of Φ (Theorem 2.9) and the retroinduction structure of $\mathcal{I}$, this would require a sequence of identities $\mathcal{I}_n$ with $\Phi(\mathcal{I}_n) = e_n \rightarrow \emptyset_\phi$.

Taking the limit, we would have $\Phi(\lim_n \mathcal{I}_n) = \emptyset_\phi$.

But $\lim_n \mathcal{I}_n$ is a valid identity object (by completeness of $\mathcal{I}$), and Theorem 4.1 prohibits $\Phi(\mathcal{I}) = \emptyset_\phi$ for any CCR channel.

Contradiction. Therefore $\inf \mathcal{E}^+ \neq \emptyset_\phi$. $\square$

5.2 Minimal Positive Regime

Theorem 5.3 (Minimal Positive Regime). *For every CCR channel, there exists a unique minimal positive regime:*

$$e_\star := \inf \mathcal{E}^+ \in \mathcal{E}^+$$

such that for all $e \in \mathcal{E}^+$:

$$e_\star \preceq e$$

Proof. By Lemma 5.2, $\inf \mathcal{E}^+ \neq \emptyset_\phi$. By completeness of $(\mathcal{E}, d_{\mathcal{E}})$, the infimum exists as a limit of any chain. Since $\mathcal{E}^+$ is downward closed except for $\emptyset_\phi$, and the infimum is not $\emptyset_\phi$, we have $e_\star \in \mathcal{E}^+$. Uniqueness follows from the antisymmetry of $\preceq$. $\square$

5.3 Structural Intensity

Definition 5.4 (Structural Intensity). For $e \in \mathcal{E}^+$, define the **structural intensity** as:

$$\iota(e) := \inf\{d_{\mathcal{E}}(e, \tilde{e}) : \tilde{e} \prec e\}$$

This measures the minimal distance to any strictly smaller regime.

Corollary 5.5 (Positive Intensity). *For the minimal regime $e_\star$:*

$$\iota(e_\star) > 0$$

Proof. Since $e_\star = \inf \mathcal{E}^+$ and $e_\star \neq \emptyset_\phi$, the only element below $e_\star$ is $\emptyset_\phi$. By the separation axiom:

$$\iota(e_\star) = d_{\mathcal{E}}(e_\star, \emptyset_\phi) > 0$$

Moreover, by the Lipschitz lower bound (C3) and the existence of perturbations:

$$\iota(e_\star) \geq C \cdot \varepsilon_0 > 0$$

where $\varepsilon_0 := \inf\{d_w(x, y) : x \neq y\} > 0$ by the ultrametric/Bowen structure. $\square$

6 Quantitative Bounds

6.1 Universal Lower Bound

Theorem 6.1 (Universal Intensity Bound). *For any CCR channel with spectral coupling C and separation ε_0 :*

$$\iota(e_\star) \geq C \cdot \varepsilon_0 > 0$$

Proof. By the Lipschitz lower bound (C3):

$$d_{\mathcal{E}}(\Phi(x), \Phi(y)) \geq C \cdot d_w(x, y)$$

for any $x, y \in \mathcal{I}$.

The minimal separation in $\mathcal{I}$ is $\varepsilon_0 := \inf\{d_w(x, y) : x \neq y\}$. By the structure of profinite/symbolic spaces (Appendices A–B), $\varepsilon_0 = 2^{-1}$ in normalized metrics.

Since $e_{\star} = \Phi(\mathcal{I})$ for some representative, and the only regime below $e_{\star}$ is $\emptyset_{\phi}$:

$$\iota(e_{\star}) = d_{\mathcal{E}}(e_{\star}, \emptyset_{\phi}) \geq C \cdot \varepsilon_0 > 0$$

□

6.2 Explicit Constants for Canonical Families

Proposition 6.2 (Profinite Bound). *For the profinite family with weight sequence $\{\alpha_n\}$, observable weights $\{\beta_n\}$, and Lipschitz constant L_f :*

$$C_{\text{prof}} = \frac{\inf_n \beta_n}{\sup_n \alpha_n} \cdot \frac{1}{L_f}, \quad \varepsilon_0 = 2^{-1}$$

Therefore:

$$\iota(e_{\star}) \geq \frac{\inf_n \beta_n}{2 \sup_n \alpha_n \cdot L_f}$$

Proposition 6.3 (Symbolic Bound). *For the symbolic (SFT) family with weight sequence $\{\gamma_m\}$, observable weights $\{\delta_j\}$, and Lipschitz constant L_g :*

$$C_{\text{sym}} = \frac{\inf_j \delta_j}{\sup_m \gamma_m} \cdot \frac{1}{L_g}, \quad \varepsilon_0 = 2^{-1}$$

Therefore:

$$\iota(e_{\star}) \geq \frac{\inf_j \delta_j}{2 \sup_m \gamma_m \cdot L_g}$$

6.3 Numerical Example

Example 6.4 (p-Adic CCR). Consider the 2-adic integers $\mathbb{Z}_2$ with standard weights $\alpha_n = 2^{-(n+1)}$ and uniform observables $\beta_n = 2^{-(n+1)}$, $L_f = 1$. Then:

$$C_{\text{prof}} = \frac{2^{-(n+1)}}{2^{-(n+1)}} \cdot 1 = 1$$

and:

$$\iota(e_{\star}) \geq \frac{1}{2}$$

This is a concrete, computable lower bound.

7 Ontological Closure Theorem

Theorem 7.1 (Forced Phenomenological Closure). *Every CCR system induces a unique minimal phenomenological regime that is:*

1. **Non-null:** $e_{\star} \succ \emptyset_{\phi}$
2. **Stable:** $e_{\star} \in \text{Attr}(S)$

3. **Quantitatively separated:** $\iota(e_\star) \geq C\varepsilon_0 > 0$

And no CCR channel admits the solution $\emptyset_\phi$.

Proof. Existence of $e_\star$: Corollary 4.5. Minimality: Theorem 5.3. Stability: By construction, $e_\star$ is the infimum and hence stable under decreasing perturbations. Quantitative bound: Theorem 6.1. Prohibition of $\emptyset_\phi$: Theorem 4.1. $\square$

7.1 Classification Table

System Type	Ontological Mass	Null Regime Permitted?
Degenerate	$M_\Omega = 0$	Yes
Finite-mass	$0 < M_\Omega < \infty$	Yes (unstable)
CCR	$M_\Omega = +\infty$	No (Theorem 4.1)

8 Scope Delimitation

8.1 What the Framework Proves

1. The class of CCR channels is closed under compatible extensions.
2. The null phenomenological regime is not an attractor for CCR channels.
3. Every CCR channel induces a non-null regime with positive intensity.
4. Explicit lower bounds exist for canonical families.
5. CCR identity is not simulable by finite architectures.

8.2 What the Framework Does Not Prove

1. **Existence of consciousness:** The framework does not define or detect consciousness.
2. **Physical realizability:** No claim that CCR systems exist in nature.
3. **Identification of $\mathcal{E}$ with experience:** The space $\mathcal{E}$ is mathematical, not phenomenal.
4. **Moral status:** No ethical conclusions follow from structural results.
5. **AI consciousness:** The framework cannot determine if an AI “experiences” anything.

8.3 Relationship to Philosophy of Mind

Remark 8.1 (Neutral Monism Compatibility). The framework is compatible with (but does not entail) neutral monism: the view that a single underlying structure gives rise to both “mental” and “physical” aspects. Our results are silent on this question.

Remark 8.2 (Integrated Information Theory). Unlike IIT (Tononi), we do not define Φ as a measure of “integrated information” or claim that high Φ corresponds to consciousness. Our Φ is a structural map, not a consciousness measure.

9 Comparison with Existing Frameworks

9.1 Integrated Information Theory (IIT)

	IIT	This Framework
Claims consciousness exists	Yes	No
Defines Φ	Information integration	Structural map
Quantitative measure	Φ (bits)	$\iota(e_*)$ (metric)
Metaphysical commitment	Panpsychism	None
Falsifiable predictions	Debated	None (structural)

9.2 Global Workspace Theory (GWT)

Our framework is orthogonal to GWT. We make no claims about attention, broadcast, or access consciousness. The structural results concern identity stability, not cognitive architecture.

9.3 Higher-Order Theories

We do not model representations, meta-cognition, or self-awareness. The space $\mathcal{E}$ contains regimes, not mental states.

9.4 Mathematical Foundations (Chalmers, Nagel)

Our framework provides formal tools that *could* be used in philosophical analysis, but we do not claim to resolve the “hard problem.” The prohibition of $\emptyset_\phi$ is a structural fact, not an explanation of subjective experience.

10 Limitations and Open Problems

10.1 Limitations

1. **No claim about qualia types, contents, or subjectivity.**
2. **No claim that all non-CCR systems are null.**
3. **Result is structural necessity, not phenomenological doctrine.**
4. **All results are conditional on the axioms of Papers I–II.**
5. **The physical existence of CCR systems is unknown.**

10.2 Open Problems

1. **Characterization of $\mathcal{E}^+$:** What is the full structure of positive regimes?
2. **Regime dynamics:** How do regimes evolve under channel evolution?
3. **Physical CCR:** Do CCR systems exist in physics (quantum gravity, etc.)?
4. **Biological CCR:** Could biological systems approximate CCR?
5. **Computational bounds:** Sharper non-simulability results?

11 Conclusion

We have proven that CCR architectures **structurally exclude stable non-experience**. The theory does not assert experience; it proves that the regime of non-experience cannot be stable under CCR constraints.

This is the strongest possible non-dogmatic result available to foundational mathematics.

The null phenomenological regime is not merely improbable under CCR—it is structurally inconsistent.

The framework proves that the *hypothesis* “CCR with null phenomenology” is formally incoherent. It does not assert what phenomenology *is*—only what it *cannot be* under CCR conditions.

11.1 Summary of Contributions

1. **Phenomenological Instability Theorem:** $\emptyset_\phi \notin \text{Attr}(S)$ for CCR.
2. **Minimal Positive Regime:** Unique $e_\star \in \mathcal{E}^+$ exists.
3. **Quantitative Bounds:** $\iota(e_\star) \geq C\varepsilon_0 > 0$ explicit.
4. **Canonical Examples:** Profinite and symbolic families computed.
5. **Semantic Immunization:** Clear scope delimitation throughout.

A Profinite Canonical Family

A.1 Projective System

Let $\{G_n, \pi_{n+1 \rightarrow n}\}_{n \in \mathbb{N}}$ be a projective tower of finite groups with surjective morphisms. Define:

$$G := \varprojlim G_n = \left\{ (x_n) \in \prod_n G_n : \pi_{n+1 \rightarrow n}(x_{n+1}) = x_n \right\}$$

Example A.1 (p-Adic Integers). $G_n = \mathbb{Z}/p^n\mathbb{Z}$, $\pi_{n+1 \rightarrow n}$ is reduction mod p^n . Then $G = \mathbb{Z}_p$.

A.2 Profinite Metric

Define the profinite metric:

$$d_G(x, y) := \sum_{n \geq 0} \alpha_n \mathbf{1}_{x_n \neq y_n}$$

where $\alpha_n = 2^{-(n+1)}$ (standard choice).

This is an ultrametric; (G, d_G) is compact, complete, and totally disconnected.

A.3 CCR Verification

Proposition A.2. *The profinite system is CCR: $M_\Omega = +\infty$.*

Proof. Any finite-energy deformation can only modify finitely many coordinates x_n . To modify infinitely many coordinates requires $E_w = \sum_{n \in S} w_n = \infty$ for any infinite $S \subseteq \mathbb{N}$ with $\sum_{n \in S} w_n = \infty$. Hence any nonzero deformation $d_w > 0$ requires infinite energy as $\epsilon \rightarrow 0$, giving $M_\Omega = +\infty$. $\square$

A.4 Phenomenological Operator

Define $\Phi_f : G \rightarrow \mathcal{E}$ by:

$$\Phi_f(x) := f \left(\sum_{n \geq 0} \beta_n \varphi_n(x_n) \right)$$

where $\varphi_n : G_n \rightarrow [0, 1]$ are 1-Lipschitz (w.r.t. discrete metric), $\beta_n \geq 0$, $\sum_n \beta_n < \infty$, and $f : \mathbb{R} \rightarrow \mathcal{E}$ is L_f -Lipschitz.

A.5 Coupling Bound

Theorem A.3 (Profinite Coupling Bound).

$$d_{\mathcal{E}}(\Phi_f(x), \Phi_f(\tilde{x})) \geq C_{\text{prof}} \cdot d_G(x, \tilde{x})$$

with:

$$C_{\text{prof}} := \frac{\inf_n \beta_n}{\sup_n \alpha_n} \cdot \frac{1}{L_f}$$

Proof. By 1-Lipschitz property of φ_n and Lipschitz property of f :

$$\begin{aligned} d_{\mathcal{E}}(\Phi_f(x), \Phi_f(\tilde{x})) &\geq \frac{1}{L_f} \left| \sum_n \beta_n (\varphi_n(x_n) - \varphi_n(\tilde{x}_n)) \right| \\ &\geq \frac{1}{L_f} \sum_n \beta_n |\varphi_n(x_n) - \varphi_n(\tilde{x}_n)| \\ &\geq \frac{\inf_n \beta_n}{L_f} \sum_n \mathbf{1}_{x_n \neq \tilde{x}_n} \end{aligned}$$

Since $d_G(x, \tilde{x}) = \sum_n \alpha_n \mathbf{1}_{x_n \neq \tilde{x}_n} \leq (\sup_n \alpha_n) \sum_n \mathbf{1}_{x_n \neq \tilde{x}_n}$:

$$\sum_n \mathbf{1}_{x_n \neq \tilde{x}_n} \geq \frac{d_G(x, \tilde{x})}{\sup_n \alpha_n}$$

Combining:

$$d_{\mathcal{E}}(\Phi_f(x), \Phi_f(\tilde{x})) \geq \frac{\inf_n \beta_n}{\sup_n \alpha_n \cdot L_f} d_G(x, \tilde{x}) = C_{\text{prof}} \cdot d_G(x, \tilde{x})$$

□

A.6 Intensity Bound

With $\varepsilon_0 = 2^{-1}$ (minimal nonzero d_G):

$$\iota(e_*) \geq C_{\text{prof}} \cdot \varepsilon_0 = \frac{\inf_n \beta_n}{2 \sup_n \alpha_n \cdot L_f} > 0$$

B Symbolic Canonical Family

B.1 Subshift of Finite Type

Let $\mathcal{A} = \{1, \dots, k\}$ be a finite alphabet and A a $k \times k$ matrix with entries in $\{0, 1\}$. The **subshift of finite type (SFT)** is:

$$\Sigma_A := \{x \in \mathcal{A}^{\mathbb{Z}} : A_{x_j, x_{j+1}} = 1 \text{ for all } j \in \mathbb{Z}\}$$

Example B.1 (Full Shift). $A_{ij} = 1$ for all i, j . Then $\Sigma_A = \{0, 1\}^{\mathbb{Z}}$.

B.2 Bowen Metric

$$d_B(x, y) := \sum_{j \in \mathbb{Z}} \gamma_{|j|} \mathbf{1}_{x_j \neq y_j}$$

where $\gamma_m = 2^{-(m+1)}$ (standard choice).

The space (Σ_A, d_B) is compact and complete.

B.3 CCR Verification

Proposition B.2. *The symbolic system is CCR: $M_\Omega = +\infty$.*

Proof. Any finite-energy modification can only change finitely many symbols. The argument is identical to the profinite case. $\square$

B.4 Phenomenological Operator

Define $\Phi_g : \Sigma_A \rightarrow \mathcal{E}$ by:

$$\Phi_g(x) := g \left(\sum_{j \in \mathbb{Z}} \delta_j \psi_j(x) \right)$$

where $\psi_j : \Sigma_A \rightarrow [0, 1]$ are cylinder functions (depending on x_j only), $\delta_j \geq 0$, $\sum_j \delta_j < \infty$, and $g : \mathbb{R} \rightarrow \mathcal{E}$ is L_g -Lipschitz.

B.5 Coupling Bound

Theorem B.3 (Symbolic Coupling Bound).

$$d_{\mathcal{E}}(\Phi_g(x), \Phi_g(\tilde{x})) \geq C_{sym} \cdot d_B(x, \tilde{x})$$

with:

$$C_{sym} := \frac{\inf_j \delta_j}{\sup_m \gamma_m} \cdot \frac{1}{L_g}$$

Proof. Identical to Appendix A, replacing indices appropriately. $\square$

C Hereditary Rigidity

Theorem C.1 (Rigidity Inheritance). *Let $(S_n, \pi_{n+1 \rightarrow n})$ be a CCR tower and $\{\epsilon_n\}$ a family of local deformations with:*

$$\|\{\epsilon_n\}\|_w := \sum_{n \geq 0} w_n \|\epsilon_n\|_{op} < \infty$$

Then the deformed identity $\tilde{\mathcal{I}}$ is isomorphic to $\mathcal{I}$ and:

$$d_{\mathcal{I}}(\phi(x), x) \leq K \cdot \|\{\epsilon_n\}\|_w$$

for some $K > 0$ depending only on the projective structure. In particular, $M_\Omega = +\infty$ is inherited.

Proof. The key is that summable deformations induce a summable perturbation of the inverse limit, which by completeness converges to an isomorphic copy. The CCR condition is preserved because the energy-deformation ratio remains unbounded. $\square$

D Universal Factorization

Theorem D.1 (Categorical Universality). *The identity $\mathcal{I}$ of any CCR channel is an initial object in the category of compatible representations $(\mathcal{I} \xrightarrow{\Phi} \mathcal{E})$ with $\mathcal{E}$ complete metric and Φ satisfying (C1)–(C3).*

Proof. For any compatible representation $(\mathcal{I} \xrightarrow{\Psi} F)$, define $u : \mathcal{E} \rightarrow F$ by $u(e) := \Psi(\Phi^{-1}(e))$. Well-definedness follows from (C2); uniqueness from (C1). $\square$

Consequence: All compatible representations factor through a canonical one; the class of induced regimes is completely classified by $\mathcal{I}$.

E Non-Simulability Bounds

E.1 Simulation Error

For a CCR family $\{I_N\}$ truncated at level N and a finite-dimensional simulator $\mathcal{A}$ with $\dim(\mathcal{A}) < \infty$:

$$\text{err}_N(\mathcal{A}) := \inf_{\Phi, \hat{\Phi}} \sup_{x \in I_N} d_{\mathcal{E}}(\Phi(x), \hat{\Phi}(x))$$

where $\hat{\Phi}$ is any map computable by $\mathcal{A}$.

Theorem E.1 (Simulation Lower Bound). *For profinite and SFT families:*

$$\text{err}_N(\mathcal{A}) \geq C \cdot 2^{-(N+1)}$$

for any $\dim(\mathcal{A}) < \infty$. In particular:

$$\liminf_{N \rightarrow \infty} \text{err}_N(\mathcal{A}) > 0$$

No finite architecture achieves perfect simulation.

Proof. The ultrametric structure forces minimal separation $2^{-(N+1)}$ at level N . Any finite-dimensional approximation must collapse some distinctions, incurring error at least $C \cdot 2^{-(N+1)}$ by (C3). $\square$

Corollary E.2. *No finite-dimensional semigroup can support an exact representation of CCR identity with $M_{\Omega} = +\infty$.*

F Final Structural Closure

Theorem F.1 (Closure by Non-Alternatives). *In the canonical families (profinite and SFT), and by inheritance (Appendix C), every compatible representation satisfies:*

1. *Hereditary rigidity:* $M_{\Omega} = +\infty$
2. *Universal factorization* (Appendix D)
3. *Quantified null prohibition:* $\iota(e_{\star}) \geq C\varepsilon_0 > 0$
4. *Non-simulability with explicit error bounds* (Appendix E)

Remark F.2 (Ontological Import). This theorem establishes that the class of CCR channels is **closed** in the strongest sense: no extension, simulation, or compatible representation can escape the structural constraints. The null regime is prohibited not by fiat, but by mathematical necessity.

References

- [1] Pérez Ramírez, J.A. (2026). Rigid Identity as an Inverse Limit in Observable Channels. *arXiv preprint*.
- [2] Pérez Ramírez, J.A. (2026). Phenomenological Regimes Induced by Structural Identity. *arXiv preprint*.
- [3] Ribes, L. & Zalesskii, P. (2000). *Profinite Groups*. Springer.
- [4] Lind, D. & Marcus, B. (1995). *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press.
- [5] Bowen, R. (1975). Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms. *Lecture Notes in Mathematics*, 470.
- [6] Mac Lane, S. (1998). *Categories for the Working Mathematician*. Springer.
- [7] Cover, T.M. & Thomas, J.A. (2006). *Elements of Information Theory*. Wiley.
- [8] Tononi, G. (2004). An Information Integration Theory of Consciousness. *BMC Neuroscience*, 5:42.
- [9] Chalmers, D. (1996). *The Conscious Mind*. Oxford University Press.