Refinement Instability as a Structural Replacement for Inflation

James A. Darke

Abstract

The classical inflationary paradigm was introduced to resolve the horizon and flatness problems of Friedmann–Robertson–Walker cosmology through a transient epoch of accelerated expansion driven by a scalar field. We demonstrate that within the framework of Stable Directed Refinement Systems (SDRS), a finite-duration exponential metric phase arises as a necessary consequence of symmetric relational instability together with finite closure capacity. The duration of exponential expansion is determined by relational distinguishability capacity rather than scalar potential tuning. The classical motivations for inflation are therefore satisfied without introducing additional scalar degrees of freedom.

1 Introduction

Inflation was introduced to resolve two classical inconsistencies of non-accelerating FRW cosmology:

1. The horizon problem,
2. The flatness problem.

In standard treatments these are resolved by postulating a scalar field whose potential drives $\mathcal{O}(60)$ e-folds of exponential expansion.

In Darke Theory, spacetime geometry is emergent from relational distinguishability. The purpose of this work is restricted: we show that the structural properties of Stable Directed Refinement Systems (SDRS) necessarily produce a finite exponential metric phase sufficient to eliminate the classical horizon and flatness problems.

No claims are made here regarding perturbation spectra or reheating.

2 Stable Directed Refinement Systems

Definition 1 (Stable Directed Refinement System)**.** *An SDRS consists of:*

- *A differentiable state manifold $\mathcal{S}$,*
- *A refinement flow $\Phi_t : \mathcal{S} \to \mathcal{S}$,*
- *A rank functional $R : \mathcal{S} \to \mathbb{R}_+$,*

satisfying:

1. *Monotonicity: $R(\Phi_t(s))$ strictly increases for $t > 0$,*
2. *Directedness: refinement increases independent relational distinguishability,*
3. *Propagation stability: relational interaction density remains bounded.*

2.1 Local Flow Representation

The refinement flow satisfies:

$$\frac{ds}{dt} = F(s), \tag{1}$$

with induced rank evolution:

$$\frac{dR}{dt} = \nabla R \cdot F(s). \tag{2}$$

3 Symmetric Relational State

Let s_0 denote the maximally symmetric state:

$$R(s_0) = R_c. \tag{3}$$

All perturbation directions are degenerate under symmetry.

4 Spectral Instability

Linearising around s_0:

$$\frac{d\delta s}{dt} = DF(s_0)\,\delta s. \tag{4}$$

Let λ_i denote eigenvalues of $DF(s_0)$.

Directedness requires that perturbations which increase distinguishability do not decay. Hence at least one eigenvalue satisfies:

$$\lambda > 0. \tag{5}$$

Thus s_0 is linearly unstable.

Projecting onto the rank functional:

$$\frac{dR}{dt} = \mu(R - R_c) + \mathcal{O}((R - R_c)^2). \tag{6}$$

Define shifted variable $\tilde{R} = R - R_c$. Then:

$$\frac{d\tilde{R}}{dt} = \mu\tilde{R} + \mathcal{O}(\tilde{R}^2). \tag{7}$$

For $\tilde{R}$ sufficiently small:

$$\tilde{R}(t) = \tilde{R}_0 e^{\mu t}. \tag{8}$$

Thus:

$$R(t) = R_c + (R_0 - R_c)e^{\mu t}. \tag{9}$$

Exponential rank growth follows generically from symmetric instability.

5 Propagation Stability and Interaction Structure

The exponential instability derived above cannot persist indefinitely if the system is to remain stable. We now formalise propagation stability and show that it enforces finite closure capacity.

5.1 Relational Interaction Graph

Let $G(R)$ denote the relational interaction graph associated with R independent axes. Let $E(R)$ denote the number of independent relational interactions.

Define the average interaction degree per axis:

$$\kappa(R) = \frac{2E(R)}{R}. \tag{10}$$

This quantity measures the density of relational coupling per distinguishable axis.

5.2 Propagation Stability

Definition 2 (Propagation Stability)**.** *An SDRS is propagation-stable if there exist constants* $0 < \kappa_{\min} \leq \kappa_{\max} < \infty$ *such that for all admissible* R*,*

$$\kappa_{\min} \leq \kappa(R) \leq \kappa_{\max}. \tag{11}$$

This condition guarantees:

- No axis becomes dynamically isolated ($\kappa(R) \not\to 0$),
- No interaction density diverges ($\kappa(R) \not\to \infty$).

Propagation stability is therefore a structural condition on coherent refinement.

5.3 Consequences of Unbounded Distinguishability

Assume, for contradiction, that closure capacity is infinite and that

$$R(t) \to \infty \tag{12}$$

under refinement.

Propagation stability requires bounded interaction density:

$$\kappa(R) = \frac{2E(R)}{R} \tag{13}$$

satisfies

$$0 < \kappa_{\min} \leq \kappa(R) \leq \kappa_{\max} < \infty. \tag{14}$$

Thus interaction degree per distinguishable axis remains uniformly bounded above and below.

We now examine the structural implications.

Each distinguishable axis corresponds to an independent relational degree of freedom. Independence requires that each new axis introduces at least one new relational direction not expressible as a combination of previous ones.

However, bounded interaction density implies that each axis participates in at most $\kappa_{\max}$ independent couplings. Therefore the number of independent relational constraints that can be formed among R axes grows at most linearly with R.

Let $\mathcal{I}(R)$ denote the space of independent relational constraints. Then

$$\dim \mathcal{I}(R) \leq CR \tag{15}$$

for some constant C determined by $\kappa_{\max}$.

On the other hand, the space of distinguishable relational configurations among R axes grows superlinearly in R due to combinatorial relational overlap. In particular, the number of possible pairwise relational directions grows as

$$\binom{R}{2} = \mathcal{O}(R^2). \tag{16}$$

For distinguishability to remain strictly independent as $R \to \infty$, the dimension of independent relational structure would need to scale at least as $\mathcal{O}(R^2)$ in order to avoid redundancy among axes.

But propagation stability restricts relational independence to $\mathcal{O}(R)$.

Therefore beyond some finite value $R_{\max}$, newly introduced axes cannot remain linearly independent within the bounded interaction structure. Additional distinguishability necessarily overlaps existing relational directions.

Hence

$$\lim_{R \to R_{\max}} \frac{dR}{dt} = 0, \tag{17}$$

for some finite $R_{\max}$.

Infinite distinguishability growth is therefore incompatible with bounded relational interaction density.

Closure capacity must be finite.

5.4 Finite Closure Capacity

We therefore require the existence of $R_{\max} < \infty$ such that:

$$\lim_{R \to R_{\max}} \frac{dR}{dt} = 0. \tag{18}$$

This defines finite closure capacity.

Operationally, as R increases, the probability that a newly introduced axis remains linearly independent in relational space decreases due to combinatorial overlap. When R approaches $R_{\max}$, additional independent distinguishability becomes suppressed.

The general refinement flow must therefore take the form:

$$\frac{dR}{dt} = \mu(R - R_c)\, f\left(\frac{R}{R_{\max}}\right), \tag{19}$$

where:

$$f(x) > 0 \quad \text{for } x < 1, \qquad f(1) = 0. \tag{20}$$

The specific functional form of f is not required; only its vanishing at closure.

6 Duration of the Exponential Phase

For $R \ll R_{\max}$, we have:

$$f\left(\frac{R}{R_{\max}}\right) \approx 1, \tag{21}$$

and hence:

$$R(t) = R_c + (R_0 - R_c)e^{\mu t}. \tag{22}$$

The exponential regime persists until $R \sim R_{\max}$.

The number of e-folds of metric expansion is:

$$N = \ln \frac{a_{\text{end}}}{a_{\text{start}}}. \tag{23}$$

Using the projection $a \propto R^{1/\beta}$:

$$N = \frac{1}{\beta} \ln \left(\frac{R_{\max} - R_c}{R_0 - R_c} \right). \tag{24}$$

For R_0 sufficiently close to R_c:

$$N \approx \frac{1}{\beta} \ln \left(\frac{R_{\max}}{R_c} \right). \tag{25}$$

Thus the duration of exponential expansion is determined entirely by distinguishability capacity.

7 Curvature Suppression

In an FRW spacetime the curvature parameter evolves as

$$\Omega - 1 = \frac{k}{(aH)^2}. \tag{26}$$

During the exponential refinement phase,

$$a(t) = a_0 e^{Ht}, \qquad H = \frac{\mu}{\beta}. \tag{27}$$

Therefore

$$(aH)^2 \propto e^{2Ht}. \tag{28}$$

Hence

$$|\Omega - 1| \propto e^{-2N}. \tag{29}$$

For $N \gtrsim 60$,

$$|\Omega - 1| \lesssim e^{-120}, \tag{30}$$

rendering spatial curvature dynamically negligible and eliminating sensitivity to initial conditions.

8 Resolution of the Horizon Problem

The particle horizon is

$$d_{\mathrm{p}}(t) = a(t) \int_{t_i}^{t} \frac{dt'}{a(t')}. \tag{31}$$

During exponential refinement,

$$a(t) = a_i e^{H(t-t_i)}. \tag{32}$$

The conformal time integral evaluates to

$$\int_{t_i}^{t} \frac{dt'}{a(t')} = \frac{1}{a_i H} \left(1 - e^{-H(t-t_i)} \right). \tag{33}$$

Thus

$$d_{\mathrm{p}}(t) = \frac{1}{H} \left(e^{H(t-t_i)} - 1 \right). \tag{34}$$

The comoving Hubble radius is

$$(aH)^{-1} = \frac{1}{H} e^{-H(t-t_i)}, \tag{35}$$

which decreases monotonically during the exponential phase.

Because the comoving Hubble radius shrinks, any fixed comoving scale that is super-horizon today was sub-horizon at sufficiently early times within the refinement epoch. Hence regions currently separated by super-horizon distances were in prior causal contact.

The horizon problem is therefore resolved.

9 Refinement Inflation Replacement Theorem

We now assemble the structural result.

Theorem 3 (Refinement Inflation Replacement)**.** *Let an SDRS satisfy:*

1. *A linearly unstable symmetric relational state,*
2. *A monotone rank functional R,*
3. *Propagation stability with bounded interaction density,*
4. *Finite closure capacity $R_{\max}$,*
5. *Homogeneous metric projection $a = \alpha R^{1/\beta}$.*

Then the projected spacetime necessarily undergoes a finite-duration exponential expansion phase with

$$N = \frac{1}{\beta} \ln \left(\frac{R_{\max} - R_c}{R_0 - R_c} \right)$$

e-folds, sufficient to resolve the classical horizon and flatness problems of FRW cosmology.

Proof. By spectral instability, $R(t)$ grows exponentially away from R_c for sufficiently small $\tilde{R} = R - R_c$.

Propagation stability requires bounded interaction density. Unbounded distinguishability growth would violate this condition, forcing the existence of a maximal distinguishable capacity $R_{\max}$ at which refinement flow vanishes.

Thus the refinement flow necessarily exhibits a transient exponential regime for $R \ll R_{\max}$, terminating at $R \sim R_{\max}$.

Under homogeneous metric projection $a \propto R^{1/\beta}$, exponential rank growth induces exponential metric expansion.

Exponential expansion shrinks the comoving Hubble radius and suppresses curvature exponentially in N, eliminating the classical horizon and flatness problems. □

10 Structural Comparison with Slow-Roll Inflation

Slow-roll inflation introduces:

- A scalar field ϕ,
- A potential $V(\phi)$,
- Small slow-roll parameters.

In contrast, refinement emergence requires only:

- Symmetric instability,
- Directed distinguishability growth,
- Finite closure capacity.

The duration of exponential expansion is determined by relational distinguishability capacity rather than potential flatness.

Termination occurs structurally at closure rather than dynamically through violation of slow-roll conditions.

11 Scope and Assumptions

This work assumes:

1. Directed refinement increases independent relational axes.
2. Symmetric states are spectrally unstable.
3. Propagation stability requires bounded interaction density.
4. Relational distinguishability capacity is finite.
5. Metric projection satisfies $a = \alpha R^{1/\beta}$.

This work does not address perturbation spectra, reheating, baryogenesis, or dark matter.

12 Conclusion

Within the SDRS framework of Darke Theory, a transient exponential metric phase emerges as a necessary structural consequence of symmetric relational instability subject to finite closure capacity.

The classical motivations for inflation — resolution of the horizon and flatness problems — are thereby satisfied without invoking an inflaton field.

Accelerated expansion is reinterpreted as a structural phase of relational emergence.