

TOP 200 DRUGS DECK

THE ESSENTIAL 200 CARDS

Double-Sided Study Cards · Color-Coded Categories

CLINICAL STUDY DECK SETUP

PRINTING SETTINGS

- **Paper:** Heavy cardstock, 8.5 x 11 in
- **Scale:** 100% / Actual Size
- **Sides:** Print double-sided
- **Flip:** On the long edge
- **Color:** Full color
- **Test first:** Print pages 3-4 on plain paper to check alignment

STUDY & ASSEMBLY

- **Cut:** Follow the dashed lines, using the scissor marks as guides.
- **Recall:** Look at the colored back first and recall the details before flipping.
- **Notes:** Add your own memory tricks in the dashed box on the front.

COLOR CODING SYSTEM

Every group below has its own color, shown on both sides of each card. Use the color to group related cards and quiz yourself by class before flipping.



Cardiovascular



Anti-Infectives



Vitamins & Supplements



Endocrine



Women's & Men's Health



Eye & Skin



Respiratory



Pain & Musculoskeletal



Immunology



Gastrointestinal



Renal & Electrolytes



Neuro & Psych





Allergy

ALIGNMENT TEST • FRONT

= 1 inch



ALIGNMENT TEST • BACK



ATORVASTATIN

CLASS: Statin

USE: High cholesterol, CV risk reduction

MOA: Inhibits HMG-CoA reductase, lowering LDL

SE: Muscle pain, rhabdomyolysis, raised LFTs

NOTES: Lipitor. Any time of day. Report unexplained muscle pain.

My Notes / Clues:

© 2020 Demo Study Shop

LEVOTHYROXINE

BLACK BOX WARNING

CLASS: Thyroid Hormone

USE: Hypothyroidism

MOA: Synthetic T4 replaces thyroid hormone

SE: Tachycardia, tremor, weight loss, insomnia (overdose)

NOTES: Synthroid. Empty stomach 30-60 min before breakfast. Separate from calcium/iron by 4 hr.

My Notes / Clues:

© 2020 Demo Study Shop

METFORMIN

BLACK BOX WARNING

CLASS: Biguanide

USE: Type 2 diabetes

MOA: Decreases liver glucose output; improves insulin sensitivity

SE: GI upset, diarrhea, B12 deficiency, rare lactic acidosis

NOTES: Glucophage. Take with meals. Hold for iodinated contrast; check kidney function.

My Notes / Clues:

© 2020 Demo Study Shop

AMLODIPINE

CLASS: Calcium Channel Blocker (DHP)

USE: Hypertension, angina

MOA: Blocks calcium entry into vascular smooth muscle

SE: Ankle edema, flushing, headache, dizziness

NOTES: Norvasc. Once daily. Edema is dose-related, not fluid overload.

My Notes / Clues:

© 2020 Demo Study Shop

LISINAPRIL

BLACK BOX WARNING

CLASS: ACE Inhibitor

USE: Hypertension, heart failure, post-MI

MOA: Blocks conversion of angiotensin I to II

SE: Dry cough, hyperkalemia, angioedema, dizziness

NOTES: Zestril / Prinivil. Monitor BP, K+, creatinine. Avoid in pregnancy.

My Notes / Clues:

© 2020 Demo Study Shop

ALBUTEROL

CLASS: Short-Acting Beta-2 Agonist

USE: Acute bronchospasm, asthma, COPD

MOA: Stimulates beta-2 receptors to relax airway muscle

SE: Tachycardia, tremor, nervousness, hypokalemia

NOTES: ProAir / Ventolin. Rescue inhaler. Needing it more than 2 days/week signals poor control.

My Notes / Clues:

© 2020 Demo Study Shop

LOSARTAN

BLACK BOX WARNING

CLASS: ARB

USE: Hypertension, diabetic nephropathy

MOA: Blocks angiotensin II at its receptor

SE: Dizziness, hyperkalemia, hypotension

NOTES: Cozaar. No ACE-inhibitor cough. Avoid in pregnancy.

My Notes / Clues:

© 2020 Demo Study Shop

METOPROLOL

BLACK BOX WARNING

CLASS: Beta Blocker (Beta-1 Selective)

USE: Hypertension, angina, heart failure, post-MI

MOA: Blocks beta-1 receptors, lowering HR and contractility

SE: Bradycardia, fatigue, hypotension, masks hypoglycemia

NOTES: Lopressor / Toprol-XL. Check apical pulse/BP. Never stop abruptly.

My Notes / Clues:

© 2020 Demo Study Shop

ROSUVASTATIN

CLASS: Statin

USE: High cholesterol, CV risk reduction

MOA: Inhibits HMG-CoA reductase

SE: Myalgia, raised LFTs, proteinuria

NOTES: Crestor. High-intensity statin. Any time of day. Lower doses in Asian patients.

My Notes / Clues:

© 2020 Demo Study Shop

BIGUANIDE

© 2026 Demo Study Shop

**THYROID
HORMONE**

© 2026 Demo Study Shop

STATIN

© 2026 Demo Study Shop

**SHORT-ACTING BETA-2
AGONIST**

© 2026 Demo Study Shop

ACE INHIBITOR

© 2026 Demo Study Shop

**CALCIUM CHANNEL
BLOCKER (DHP)**

© 2026 Demo Study Shop

STATIN

© 2026 Demo Study Shop

**BETA BLOCKER (BETA-1
SELECTIVE)**

© 2026 Demo Study Shop

ARB

© 2026 Demo Study Shop

OMEPRAZOLE

CLASS: Proton Pump Inhibitor

USE: GERD, ulcers, H. pylori (combo)

MOA: Irreversibly blocks the gastric proton pump

SE: Headache, diarrhea; long-term: C. diff, fractures, low Mg, low B12

NOTES: Prilosec. Take 30-60 min before breakfast. Use the lowest effective duration.

My Notes / Clues:

© 2026 Demo Study Shop

GABAPENTIN

CLASS: Gabapentinoid
Anticonvulsant

USE: Neuropathic pain, postherpetic neuralgia, seizures

MOA: Binds calcium channel alpha-2-delta subunit, reducing excitatory signals

SE: Drowsiness, dizziness, edema, ataxia

NOTES: Neurontin. Taper to stop. Respiratory depression risk with opioids. Adjust in kidney disease.

My Notes / Clues:

© 2026 Demo Study Shop

SERTRALINE

BLACK BOX WARNING

CLASS: SSRI

USE: Depression, anxiety, OCD, PTSD, panic

MOA: Blocks serotonin reuptake

SE: Nausea, diarrhea, insomnia, sexual dysfunction

NOTES: Zoloft. Takes 4-6 weeks. Watch for suicidal thoughts and serotonin syndrome.

My Notes / Clues:

© 2026 Demo Study Shop

ESCITALOPRAM

BLACK BOX WARNING

CLASS: SSRI

USE: Depression, generalized anxiety

MOA: Selectively blocks serotonin reuptake

SE: Nausea, insomnia, sexual dysfunction, QT prolongation

NOTES: Lexapro. Takes 4-6 weeks. Taper to stop.

My Notes / Clues:

© 2026 Demo Study Shop

SEMAGLUTIDE

BLACK BOX WARNING

CLASS: GLP-1 Receptor Agonist

USE: Type 2 diabetes, weight management, CV risk

MOA: Boosts glucose-dependent insulin, slows gastric emptying, reduces appetite

SE: Nausea, vomiting, diarrhea, pancreatitis, gallbladder disease

NOTES: Ozempic / Wegovy / Rybelsus. Weekly injection (oral daily). Contraindicated with MTC/MEN2 history.

My Notes / Clues:

© 2026 Demo Study Shop

AMPHETAMINE MIXED SALTS

BLACK BOX WARNING

CLASS: CNS Stimulant

USE: ADHD, narcolepsy

MOA: Releases dopamine and norepinephrine

SE: Insomnia, appetite loss, tachycardia, anxiety

NOTES: Adderall. Schedule II. Give in the morning. Monitor BP, HR, growth.

My Notes / Clues:

© 2026 Demo Study Shop

PANTOPRAZOLE

CLASS: Proton Pump Inhibitor

USE: GERD, erosive esophagitis, GI bleed prophylaxis

MOA: Irreversibly blocks the gastric proton pump

SE: Headache, diarrhea; long-term low Mg, C. diff

NOTES: Protonix. Before breakfast. IV form used in hospital for GI bleeds.

My Notes / Clues:

© 2026 Demo Study Shop

BUPROPION

BLACK BOX WARNING

CLASS: NDRI Antidepressant

USE: Depression, seasonal affective disorder, smoking cessation

MOA: Blocks dopamine and norepinephrine reuptake

SE: Insomnia, dry mouth, agitation, seizures (dose-related)

NOTES: Wellbutrin. No sexual side effects. Avoid with seizure or eating disorders.

My Notes / Clues:

© 2026 Demo Study Shop

HYDROCHLOROTHIAZIDE

CLASS: Thiazide Diuretic

USE: Hypertension, edema

MOA: Blocks sodium reabsorption in the distal tubule

SE: Hypokalemia, hyponatremia, hyperglycemia, hyperuricemia

NOTES: Microzide. Take in the morning. Monitor electrolytes. Sun sensitivity.

My Notes / Clues:

© 2026 Demo Study Shop

SSRI

© 2026 Demo Study Shop

**GABAPENTINOID
ANTICONVULSANT**

© 2026 Demo Study Shop

**PROTON PUMP
INHIBITOR**

© 2026 Demo Study Shop

CNS STIMULANT

© 2026 Demo Study Shop

**GLP-1 RECEPTOR
AGONIST**

© 2026 Demo Study Shop

SSRI

© 2026 Demo Study Shop

THIAZIDE DIURETIC

© 2026 Demo Study Shop

**NDRI
ANTIDEPRESSANT**

© 2026 Demo Study Shop

**PROTON PUMP
INHIBITOR**

© 2026 Demo Study Shop

FLUOXETINE

BLACK BOX WARNING

CLASS: SSRI

USE: Depression, OCD, bulimia, panic

MOA: Blocks serotonin reuptake

SE: Insomnia, anxiety, nausea, sexual dysfunction

NOTES: Prozac. Longest half-life SSRI; take in the morning.

My Notes / Clues:

© 2020 Demo Study Shop

TRAZODONE

BLACK BOX WARNING

CLASS: SARI Antidepressant

USE: Depression; off-label insomnia

MOA: Blocks serotonin 5-HT_{2A} receptors and reuptake

SE: Sedation, orthostatic hypotension, priapism

NOTES: Desyrel. Give at bedtime. Priapism is an emergency.

My Notes / Clues:

© 2020 Demo Study Shop

MONTELUKAST

BLACK BOX WARNING

CLASS: Leukotriene Receptor Antagonist

USE: Asthma maintenance, allergic rhinitis, exercise bronchospasm

MOA: Blocks leukotriene receptors

SE: Headache, mood changes, nightmares, suicidal thoughts

NOTES: Singulair. Evening dosing. Not for acute attacks. Report behavior changes.

My Notes / Clues:

© 2020 Demo Study Shop

AMOXICILLIN

CLASS: Aminopenicillin

USE: Ear, sinus, throat, skin infections; H. pylori

MOA: Inhibits bacterial cell wall synthesis

SE: Diarrhea, rash, allergic reactions

NOTES: Amoxil. Ask about penicillin allergy. Finish the full course.

My Notes / Clues:

© 2020 Demo Study Shop

FLUTICASONE (NASAL/INHALED)

CLASS: Corticosteroid

USE: Allergic rhinitis (nasal), asthma maintenance (inhaled)

MOA: Reduces local airway inflammation

SE: Nosebleeds, oral thrush, hoarseness

NOTES: Flonase / Flovent. Rinse mouth after inhaled use. Not a rescue drug.

My Notes / Clues:

© 2020 Demo Study Shop

TAMSULOSIN

CLASS: Alpha-1 Blocker

USE: Benign prostatic hyperplasia

MOA: Relaxes prostate and bladder-neck smooth muscle

SE: Dizziness, orthostatic hypotension, abnormal ejaculation

NOTES: Flomax. Take 30 min after the same meal daily. Tell eye surgeon (floppy iris).

My Notes / Clues:

© 2020 Demo Study Shop

APIXABAN

BLACK BOX WARNING

CLASS: Factor Xa Inhibitor (DOAC)

USE: AFib stroke prevention, DVT/PE

MOA: Directly inhibits factor Xa

SE: Bleeding, bruising

NOTES: Eliquis. No INR monitoring. Stopping early raises stroke risk. Reversal: andexanet alfa.

My Notes / Clues:

© 2020 Demo Study Shop

SIMVASTATIN

CLASS: Statin

USE: High cholesterol

MOA: Inhibits HMG-CoA reductase

SE: Myopathy, rhabdomyolysis, raised LFTs

NOTES: Zocor. Take in the evening. Many CYP3A4 interactions; avoid grapefruit.

My Notes / Clues:

© 2020 Demo Study Shop

INSULIN GLARGINE

HIGH-ALERT MEDICATION

CLASS: Long-Acting Insulin

USE: Type 1 and 2 diabetes (basal)

MOA: Replaces basal insulin; promotes glucose uptake

SE: Hypoglycemia, weight gain, injection-site lipohypertrophy

NOTES: Lantus / Basaglar / Toujeo. Once daily, same time. No peak. Do not mix with other insulins.

My Notes / Clues:

© 2020 Demo Study Shop

**LEUKOTRIENE
RECEPTOR
ANTAGONIST**

© 2026 Demo Study Shop

**SARI
ANTIDEPRESSANT**

© 2026 Demo Study Shop

SSRI

© 2026 Demo Study Shop

**ALPHA-1
BLOCKER**

© 2026 Demo Study Shop

CORTICOSTEROID

© 2026 Demo Study Shop

AMINOPENICILLIN

© 2026 Demo Study Shop

**LONG-ACTING
INSULIN**

© 2026 Demo Study Shop

STATIN

© 2026 Demo Study Shop

**FACTOR XA INHIBITOR
(DOAC)**

© 2026 Demo Study Shop

EMPAGLIFLOZIN

CLASS: SGLT2 Inhibitor

USE: Type 2 diabetes, heart failure, chronic kidney disease

MOA: Blocks renal glucose reabsorption

SE: UTIs, genital yeast infections, dehydration, ketoacidosis

NOTES: Jardiance. Hold before surgery. Good genital hygiene; stay hydrated.

My Notes / Clues:

© 2026 Demo Study Shop

FUROSEMIDE

BLACK BOX WARNING

CLASS: Loop Diuretic

USE: Edema, heart failure, hypertension

MOA: Blocks Na/K/2Cl in the loop of Henle

SE: Hypokalemia, dehydration, hypotension, ototoxicity

NOTES: Lasix. Monitor K⁺, weight, I&O. Give IV push slowly.

My Notes / Clues:

© 2026 Demo Study Shop

MELOXICAM

BLACK BOX WARNING

CLASS: NSAID (COX-2 Preferential)

USE: Osteoarthritis, rheumatoid arthritis

MOA: Inhibits prostaglandin synthesis

SE: GI bleeding, kidney injury, edema, raised BP

NOTES: Mobic. Take with food. Boxed warning: CV events and GI bleeding.

My Notes / Clues:

© 2026 Demo Study Shop

HYDROCODONE / ACETAMINOPHEN

BLACK BOX WARNING

CLASS: Opioid Combination

USE: Moderate to severe pain

MOA: Mu-opioid agonist plus central analgesic

SE: Constipation, sedation, respiratory depression, dependence

NOTES: Norco / Vicodin. Schedule II. Count all acetaminophen (max 4 g/day).

My Notes / Clues:

© 2026 Demo Study Shop

TIRZEPATIDE

BLACK BOX WARNING

CLASS: GIP / GLP-1 Receptor Agonist

USE: Type 2 diabetes, weight management

MOA: Activates GIP and GLP-1 receptors to raise insulin and reduce appetite

SE: Nausea, vomiting, diarrhea, pancreatitis

NOTES: Mounjaro / Zepbound. Weekly injection. May reduce oral contraceptive absorption.

My Notes / Clues:

© 2026 Demo Study Shop

METHYLPHENIDATE

BLACK BOX WARNING

CLASS: CNS Stimulant

USE: ADHD, narcolepsy

MOA: Blocks dopamine and norepinephrine reuptake

SE: Insomnia, appetite loss, tachycardia

NOTES: Ritalin / Concerta. Schedule II. Monitor growth in children. Morning dosing.

My Notes / Clues:

© 2026 Demo Study Shop

DULOXETINE

BLACK BOX WARNING

CLASS: SNRI

USE: Depression, anxiety, neuropathic pain, fibromyalgia

MOA: Blocks serotonin and norepinephrine reuptake

SE: Nausea, dry mouth, sweating, raised BP, liver injury

NOTES: Cymbalta. Taper to stop (discontinuation syndrome). Avoid heavy alcohol.

My Notes / Clues:

© 2026 Demo Study Shop

PREDNISONE

CLASS: Systemic Corticosteroid

USE: Inflammation, asthma flares, autoimmune disease

MOA: Suppresses inflammation and immune response

SE: Hyperglycemia, insomnia, mood changes, infection risk, bone loss

NOTES: Deltasone. Take in the morning with food. Taper long courses.

My Notes / Clues:

© 2026 Demo Study Shop

CARVEDILOL

CLASS: Beta Blocker with Alpha-1 Blockade

USE: Heart failure, hypertension, post-MI

MOA: Blocks beta-1, beta-2 and alpha-1 receptors

SE: Dizziness, orthostatic hypotension, bradycardia, fatigue

NOTES: Coreg. Take with food. Do not stop abruptly.

My Notes / Clues:

© 2026 Demo Study Shop

**NSAID (COX-2
PREFERENTIAL)**

© 2026 Demo Study Shop

LOOP DIURETIC

© 2026 Demo Study Shop

SGLT2 INHIBITOR

© 2026 Demo Study Shop

CNS STIMULANT

© 2026 Demo Study Shop

**GIP / GLP-1 RECEPTOR
AGONIST**

© 2026 Demo Study Shop

**OPIOID
COMBINATION**

© 2026 Demo Study Shop

**BETA BLOCKER WITH
ALPHA-1 BLOCKADE**

© 2026 Demo Study Shop

**SYSTEMIC
CORTICOSTEROID**

© 2026 Demo Study Shop

SNRI

© 2026 Demo Study Shop

FAMOTIDINE

CLASS: H2 Receptor Antagonist

USE: GERD, peptic ulcer, heartburn

MOA: Blocks histamine H2 receptors in the stomach

SE: Headache, dizziness, confusion in older adults

NOTES: Pepcid. Reduce dose in kidney disease.

My Notes / Clues:

© 2026 Demo Study Shop

IBUPROFEN

BLACK BOX WARNING

CLASS: NSAID

USE: Pain, fever, inflammation

MOA: Non-selectively inhibits COX-1 and COX-2

SE: GI upset/bleeding, kidney injury, raised BP

NOTES: Advil / Motrin. Take with food. Avoid in late pregnancy and kidney disease.

My Notes / Clues:

© 2026 Demo Study Shop

BUSPIRONE

CLASS: Anxiolytic (5-HT1A Partial Agonist)

USE: Generalized anxiety disorder

MOA: Partial agonist at serotonin 5-HT1A receptors

SE: Dizziness, headache, nausea

NOTES: Buspar. Not habit-forming. Takes 2-4 weeks. Not for acute panic.

My Notes / Clues:

© 2026 Demo Study Shop

VENLAFAXINE

BLACK BOX WARNING

CLASS: SNRI

USE: Depression, anxiety, panic

MOA: Blocks serotonin and norepinephrine reuptake

SE: Nausea, sweating, raised BP, insomnia

NOTES: Effexor XR. Severe discontinuation symptoms; taper slowly.

My Notes / Clues:

© 2026 Demo Study Shop

TRAMADOL

BLACK BOX WARNING

CLASS: Opioid Analgesic (Mu Agonist + SNRI)

USE: Moderate to severe pain

MOA: Weak mu agonist; blocks serotonin/NE reuptake

SE: Nausea, dizziness, seizures, serotonin syndrome

NOTES: Ultram. Schedule IV. Avoid with SSRIs/SNRIs and seizure history.

My Notes / Clues:

© 2026 Demo Study Shop

POTASSIUM CHLORIDE

CLASS: Electrolyte Supplement

USE: Hypokalemia prevention and treatment

MOA: Replaces potassium

SE: GI irritation, hyperkalemia

NOTES: K-Dur / Klor-Con. Take with food and a full glass of water; do not crush ER tabs.

My Notes / Clues:

© 2026 Demo Study Shop

HYDROXYZINE

CLASS: 1st-Gen Antihistamine

USE: Anxiety, itching, sedation

MOA: Blocks H1 receptors centrally and peripherally

SE: Drowsiness, dry mouth, QT prolongation

NOTES: Atarax / Vistaril. Fall risk in older adults (anticholinergic).

My Notes / Clues:

© 2026 Demo Study Shop

ALLOPURINOL

CLASS: Xanthine Oxidase Inhibitor

USE: Gout prevention, tumor lysis prevention

MOA: Blocks uric acid production

SE: Rash (stop immediately), SJS, GI upset

NOTES: Zyloprim. Not for acute attacks. Drink plenty of fluids.

My Notes / Clues:

© 2026 Demo Study Shop

CLOPIDOGREL

BLACK BOX WARNING

CLASS: P2Y12 Inhibitor (Antiplatelet)

USE: ACS, after stents, stroke/MI prevention

MOA: Irreversibly blocks platelet ADP receptors

SE: Bleeding, bruising, rare TTP

NOTES: Plavix. Poor CYP2C19 metabolizers get less effect. Hold 5 days before surgery.

My Notes / Clues:

© 2026 Demo Study Shop

**ANXIOLYTIC (5-HT1A
PARTIAL AGONIST)**

© 2026 Demo Study Shop

NSAID

© 2026 Demo Study Shop

**H2 RECEPTOR
ANTAGONIST**

© 2026 Demo Study Shop

**ELECTROLYTE
SUPPLEMENT**

© 2026 Demo Study Shop

**OPIOID ANALGESIC
(MU AGONIST + SNRI)**

© 2026 Demo Study Shop

SNRI

© 2026 Demo Study Shop

**P2Y12 INHIBITOR
(ANTIPLATELET)**

© 2026 Demo Study Shop

**XANTHINE OXIDASE
INHIBITOR**

© 2026 Demo Study Shop

**1ST-GEN
ANTIHISTAMINE**

© 2026 Demo Study Shop

ERGOCALCIFEROL (VITAMIN D2)

CLASS: Vitamin

USE: Vitamin D deficiency, hypoparathyroidism

MOA: Converted to active vitamin D; increases calcium absorption

SE: Hypercalcemia with excess (nausea, weakness)

NOTES: Drisdol. High-dose capsules are usually weekly.

My Notes / Clues:

© 2026 Demo Study Shop

CETIRIZINE

CLASS: 2nd-Gen Antihistamine

USE: Allergic rhinitis, hives

MOA: Blocks peripheral H1 receptors

SE: Mild drowsiness, dry mouth

NOTES: Zyrtec. Most sedating 2nd-gen antihistamine.

My Notes / Clues:

© 2026 Demo Study Shop

ONDANSETRON

CLASS: 5-HT3 Antagonist (Antiemetic)

USE: Nausea/vomiting (chemo, post-op)

MOA: Blocks serotonin 5-HT3 receptors

SE: Headache, constipation, QT prolongation

NOTES: Zofran. Check ECG risk factors. ODT dissolves on the tongue.

My Notes / Clues:

© 2026 Demo Study Shop

CYCLOBENZAPRINE

CLASS: Skeletal Muscle Relaxant

USE: Acute muscle spasm

MOA: Acts centrally at the brainstem to reduce muscle tone

SE: Drowsiness, dry mouth, dizziness

NOTES: Flexeril. Short-term use only. Avoid in older adults (Beers).

My Notes / Clues:

© 2026 Demo Study Shop

SPIRONOLACTONE

CLASS: Aldosterone Antagonist Diuretic

USE: Heart failure, hypertension, edema, acne

MOA: Blocks aldosterone in the distal tubule

SE: Hyperkalemia, gynecomastia, menstrual changes

NOTES: Aldactone. Avoid potassium supplements and salt substitutes.

My Notes / Clues:

© 2026 Demo Study Shop

OXYCODONE

BLACK BOX WARNING

CLASS: Opioid Analgesic

USE: Moderate to severe pain

MOA: Mu-opioid receptor agonist

SE: Constipation, sedation, respiratory depression, dependence

NOTES: Roxicodone / OxyContin. Schedule II. Never crush ER tablets. Keep naloxone available.

My Notes / Clues:

© 2026 Demo Study Shop

ESTRADIOL

BLACK BOX WARNING

CLASS: Estrogen

USE: Menopausal symptoms, hypoestrogenism, osteoporosis prevention

MOA: Replaces estrogen

SE: Breast tenderness, nausea, clots, stroke

NOTES: Estrace. Add progestin if uterus present. Lowest dose, shortest time.

My Notes / Clues:

© 2026 Demo Study Shop

ASPIRIN

CLASS: Antiplatelet / NSAID

USE: MI and stroke prevention, pain, fever

MOA: Irreversibly inhibits COX, blocking thromboxane

SE: GI bleeding, tinnitus (toxicity)

NOTES: Bayer / Ecotrin. Avoid in children with viral illness (Reye's syndrome).

My Notes / Clues:

© 2026 Demo Study Shop

GLIPIZIDE

HIGH-ALERT MEDICATION

CLASS: Sulfonylurea

USE: Type 2 diabetes

MOA: Stimulates insulin release from beta cells

SE: Hypoglycemia, weight gain

NOTES: Glucotrol. Take 30 min before breakfast. Do not skip meals.

My Notes / Clues:

© 2026 Demo Study Shop

**5-HT₃ ANTAGONIST
(ANTIEMETIC)**

© 2026 Demo Study Shop

**2ND-GEN
ANTIHISTAMINE**

© 2026 Demo Study Shop

VITAMIN

© 2026 Demo Study Shop

OPIOID ANALGESIC

© 2026 Demo Study Shop

**ALDOSTERONE
ANTAGONIST DIURETIC**

© 2026 Demo Study Shop

**SKELETAL MUSCLE
RELAXANT**

© 2026 Demo Study Shop

SULFONYLUREA

© 2026 Demo Study Shop

**ANTIPLATELET /
NSAID**

© 2026 Demo Study Shop

ESTROGEN

© 2026 Demo Study Shop

ZOLPIDEM

BLACK BOX WARNING

CLASS: Nonbenzodiazepine Hypnotic

USE: Insomnia

MOA: Enhances GABA-A activity

SE: Next-day drowsiness, complex sleep behaviors

NOTES: Ambien. Take right before bed with 7-8 hours to sleep. Lower dose in women.

My Notes / Clues:

© 2026 Demo Study Shop

LAMOTRIGINE

BLACK BOX WARNING

CLASS: Anticonvulsant / Mood Stabilizer

USE: Seizures, bipolar maintenance

MOA: Blocks voltage-gated sodium channels

SE: Dizziness, rash, Stevens-Johnson syndrome

NOTES: Lamictal. Slow titration. Stop and report any rash.

My Notes / Clues:

© 2026 Demo Study Shop

ALPRAZOLAM

BLACK BOX WARNING

CLASS: Benzodiazepine

USE: Anxiety, panic disorder

MOA: Enhances GABA-A activity

SE: Sedation, dependence, respiratory depression with opioids

NOTES: Xanax. Schedule IV. Taper to stop; withdrawal can cause seizures.

My Notes / Clues:

© 2026 Demo Study Shop

CITALOPRAM

BLACK BOX WARNING

CLASS: SSRI

USE: Depression

MOA: Blocks serotonin reuptake

SE: Nausea, sexual dysfunction, QT prolongation

NOTES: Celexa. Max 20 mg in patients over 60.

My Notes / Clues:

© 2026 Demo Study Shop

PREGABALIN

CLASS: Gabapentinoid Anticonvulsant

USE: Neuropathic pain, fibromyalgia, seizures

MOA: Binds calcium channel alpha-2-delta subunit

SE: Dizziness, drowsiness, edema, weight gain

NOTES: Lyrica. Schedule V. Taper to stop.

My Notes / Clues:

© 2026 Demo Study Shop

CHOLECALCIFEROL (VITAMIN D3)

CLASS: Vitamin

USE: Vitamin D deficiency, bone health

MOA: Converted to active vitamin D; increases calcium absorption

SE: Hypercalcemia with excess

NOTES: Various. Fat-soluble; take with a meal containing fat.

My Notes / Clues:

© 2026 Demo Study Shop

CLONAZEPAM

BLACK BOX WARNING

CLASS: Benzodiazepine

USE: Panic disorder, seizures

MOA: Enhances GABA-A activity

SE: Sedation, ataxia, dependence

NOTES: Klonopin. Schedule IV. Long-acting; taper slowly.

My Notes / Clues:

© 2026 Demo Study Shop

AZITHROMYCIN

CLASS: Macrolide Antibiotic

USE: Respiratory infections, chlamydia

MOA: Inhibits bacterial protein synthesis (50S)

SE: Diarrhea, nausea, QT prolongation

NOTES: Zithromax. Short course (Z-Pak). Separate from Al/Mg antacids.

My Notes / Clues:

© 2026 Demo Study Shop

PRAVASTATIN

CLASS: Statin

USE: High cholesterol

MOA: Inhibits HMG-CoA reductase

SE: Myalgia, raised LFTs

NOTES: Pravachol. Few CYP interactions compared with other statins.

My Notes / Clues:

© 2026 Demo Study Shop

BENZODIAZEPINE

© 2026 Demo Study Shop

**ANTICONVULSANT /
MOOD STABILIZER**

© 2026 Demo Study Shop

**NONBENZODIAZEPINE
HYPNOTIC**

© 2026 Demo Study Shop

VITAMIN

© 2026 Demo Study Shop

**GABAPENTINOID
ANTICONVULSANT**

© 2026 Demo Study Shop

SSRI

© 2026 Demo Study Shop

STATIN

© 2026 Demo Study Shop

**MACROLIDE
ANTIBIOTIC**

© 2026 Demo Study Shop

BENZODIAZEPINE

© 2026 Demo Study Shop

VALSARTAN

BLACK BOX WARNING

CLASS: ARB

USE: Hypertension, heart failure

MOA: Blocks angiotensin II receptors

SE: Dizziness, hyperkalemia, hypotension

NOTES: Diovan. Avoid in pregnancy. Monitor K⁺ and creatinine.

My Notes / Clues:

© 2026 Demo Study Shop

EZETIMIBE

CLASS: Cholesterol Absorption Inhibitor

USE: High cholesterol (with or without statin)

MOA: Blocks intestinal cholesterol absorption

SE: Diarrhea, joint pain, raised LFTs with statins

NOTES: Zetia. Take 2 hr before or 4 hr after bile acid sequestrants.

My Notes / Clues:

© 2026 Demo Study Shop

DICLOFENAC

BLACK BOX WARNING

CLASS: NSAID

USE: Arthritis pain, inflammation (oral and topical gel)

MOA: Inhibits COX-1 and COX-2

SE: GI bleeding, liver injury, raised BP

NOTES: Voltaren. Topical gel has less systemic exposure.

My Notes / Clues:

© 2026 Demo Study Shop

INSULIN LISPRO

HIGH-ALERT MEDICATION

CLASS: Rapid-Acting Insulin

USE: Type 1 and 2 diabetes (mealtime)

MOA: Promotes glucose uptake into cells

SE: Hypoglycemia, hypokalemia

NOTES: Humalog / Admelog. Onset ~15 min. Give within 15 min of eating.

My Notes / Clues:

© 2026 Demo Study Shop

NORETHINDRONE / ETHINYL ESTRADIOL

BLACK BOX WARNING

CLASS: Combined Oral Contraceptive

USE: Pregnancy prevention

MOA: Suppresses ovulation; thickens cervical mucus

SE: Spotting, nausea, clots, raised BP

NOTES: Loestrin / Junel. Smoking over age 35 greatly raises CV risk.

My Notes / Clues:

© 2026 Demo Study Shop

PROPRANOLOL

BLACK BOX WARNING

CLASS: Beta Blocker (Non-Selective)

USE: Hypertension, migraine prevention, tremor, anxiety

MOA: Blocks beta-1 and beta-2 receptors

SE: Bradycardia, bronchospasm, fatigue, masks hypoglycemia

NOTES: Inderal. Avoid in asthma. Do not stop abruptly.

My Notes / Clues:

© 2026 Demo Study Shop

LATANOPROST

CLASS: Prostaglandin Analog (Eye)

USE: Glaucoma, ocular hypertension

MOA: Increases aqueous humor outflow

SE: Iris darkening, eyelash growth, eye redness

NOTES: Xalatan. One drop at bedtime. Remove contacts; wait 15 min.

My Notes / Clues:

© 2026 Demo Study Shop

ATENOLOL

BLACK BOX WARNING

CLASS: Beta Blocker (Beta-1 Selective)

USE: Hypertension, angina, post-MI

MOA: Blocks beta-1 receptors

SE: Bradycardia, fatigue, dizziness

NOTES: Tenormin. Renally cleared; adjust in kidney disease. Taper to stop.

My Notes / Clues:

© 2026 Demo Study Shop

LISDEXAMFETAMINE

BLACK BOX WARNING

CLASS: CNS Stimulant (Prodrug)

USE: ADHD, binge-eating disorder

MOA: Converted to dextroamphetamine; releases dopamine/NE

SE: Insomnia, appetite loss, tachycardia, dry mouth

NOTES: Vyvanse. Schedule II. Capsule can be opened into water or yogurt.

My Notes / Clues:

© 2026 Demo Study Shop

NSAID

© 2026 Demo Study Shop

**CHOLESTEROL
ABSORPTION
INHIBITOR**

© 2026 Demo Study Shop

ARB

© 2026 Demo Study Shop

**BETA BLOCKER
(NON-SELECTIVE)**

© 2026 Demo Study Shop

**COMBINED ORAL
CONTRACEPTIVE**

© 2026 Demo Study Shop

**RAPID-ACTING
INSULIN**

© 2026 Demo Study Shop

**CNS STIMULANT
(PRODRUG)**

© 2026 Demo Study Shop

**BETA BLOCKER (BETA-1
SELECTIVE)**

© 2026 Demo Study Shop

**PROSTAGLANDIN
ANALOG (EYE)**

© 2026 Demo Study Shop

DOXYCYCLINE

CLASS: Tetracycline Antibiotic

USE: Acne, Lyme disease, chlamydia, pneumonia

MOA: Inhibits bacterial protein synthesis (30S)

SE: Photosensitivity, esophagitis, GI upset

NOTES: Vibramycin. Full glass of water; stay upright 30 min. Avoid in pregnancy and young children.

My Notes / Clues:

© 2026 Demo Study Shop

AMOXICILLIN / CLAVULANATE

CLASS: Penicillin + Beta-Lactamase Inhibitor

USE: Sinusitis, otitis, bites, skin and respiratory infections

MOA: Cell wall inhibitor; clavulanate blocks beta-lactamase

SE: Diarrhea (more than amoxicillin), rash

NOTES: Augmentin. Take with food to reduce GI upset.

My Notes / Clues:

© 2026 Demo Study Shop

DULAGLUTIDE

BLACK BOX WARNING

CLASS: GLP-1 Receptor Agonist

USE: Type 2 diabetes, CV risk reduction

MOA: Increases insulin, slows gastric emptying

SE: Nausea, vomiting, diarrhea, pancreatitis

NOTES: Trulicity. Once-weekly injection.

My Notes / Clues:

© 2026 Demo Study Shop

LISINAPRIL / HYDROCHLOROTHIAZIDE

BLACK BOX WARNING

CLASS: ACE Inhibitor + Thiazide

USE: Hypertension

MOA: Blocks angiotensin II formation plus thiazide diuresis

SE: Cough, dizziness, electrolyte changes

NOTES: Zestoretic. Monitor K⁺, Na⁺, creatinine. Avoid in pregnancy.

My Notes / Clues:

© 2026 Demo Study Shop

LORAZEPAM

BLACK BOX WARNING

CLASS: Benzodiazepine

USE: Anxiety, status epilepticus, sedation

MOA: Enhances GABA-A activity

SE: Sedation, respiratory depression, dependence

NOTES: Ativan. Schedule IV. IV drug of choice for status epilepticus.

My Notes / Clues:

© 2026 Demo Study Shop

FLUTICASONE / SALMETEROL

CLASS: ICS + LABA

USE: Asthma and COPD maintenance

MOA: Reduces airway inflammation plus long-acting bronchodilation

SE: Thrush, hoarseness, headache

NOTES: Advair. Rinse mouth after use. Not for acute attacks.

My Notes / Clues:

© 2026 Demo Study Shop

INSULIN ASPART

HIGH-ALERT MEDICATION

CLASS: Rapid-Acting Insulin

USE: Type 1 and 2 diabetes (mealtime)

MOA: Promotes glucose uptake into cells

SE: Hypoglycemia, hypokalemia

NOTES: Novolog. Give 5-10 min before meals.

My Notes / Clues:

© 2026 Demo Study Shop

CELECOXIB

BLACK BOX WARNING

CLASS: NSAID (COX-2 Selective)

USE: Osteoarthritis, rheumatoid arthritis, acute pain

MOA: Selectively inhibits COX-2

SE: CV events, GI bleeding (less), edema

NOTES: Celebrex. Caution with sulfonamide allergy.

My Notes / Clues:

© 2026 Demo Study Shop

FINASTERIDE

CLASS: 5-Alpha Reductase Inhibitor

USE: BPH, male-pattern hair loss

MOA: Blocks conversion of testosterone to DHT

SE: Decreased libido, erectile dysfunction

NOTES: Proscar / Propecia. Pregnant people should not handle crushed tablets. Lowers PSA.

My Notes / Clues:

© 2026 Demo Study Shop

**GLP-1 RECEPTOR
AGONIST**

© 2026 Demo Study Shop

**PENICILLIN +
BETA-LACTAMASE
INHIBITOR**

© 2026 Demo Study Shop

**TETRACYCLINE
ANTIBIOTIC**

© 2026 Demo Study Shop

ICS + LABA

© 2026 Demo Study Shop

BENZODIAZEPINE

© 2026 Demo Study Shop

**ACE INHIBITOR +
THIAZIDE**

© 2026 Demo Study Shop

**5-ALPHA REDUCTASE
INHIBITOR**

© 2026 Demo Study Shop

**NSAID (COX-2
SELECTIVE)**

© 2026 Demo Study Shop

**RAPID-ACTING
INSULIN**

© 2026 Demo Study Shop

QUETIAPINE

BLACK BOX WARNING

CLASS: Atypical Antipsychotic

USE: Schizophrenia, bipolar disorder, adjunct for depression

MOA: Blocks dopamine D2 and serotonin 5-HT_{2A}

SE: Sedation, weight gain, hyperglycemia, orthostasis

NOTES: Seroquel. Monitor weight, glucose, lipids.

My Notes / Clues:

© 2026 Demo Study Shop

CLONIDINE

CLASS: Central Alpha-2 Agonist

USE: Hypertension, ADHD, withdrawal symptoms

MOA: Reduces sympathetic outflow from the brain

SE: Drowsiness, dry mouth, bradycardia

NOTES: Catapres. Stopping abruptly causes rebound hypertension.

My Notes / Clues:

© 2026 Demo Study Shop

ARIPIPRAZOLE

BLACK BOX WARNING

CLASS: Atypical Antipsychotic

USE: Schizophrenia, bipolar disorder, adjunct for depression

MOA: Partial agonist at D2 and 5-HT_{1A}

SE: Akathisia, insomnia, impulse-control problems

NOTES: Abilify. Lower metabolic risk than other atypicals.

My Notes / Clues:

© 2026 Demo Study Shop

CEPHALEXIN

CLASS: 1st-Gen Cephalosporin

USE: Skin infections, UTI, strep throat

MOA: Inhibits bacterial cell wall synthesis

SE: Diarrhea, rash

NOTES: Keflex. Low cross-reactivity with penicillin allergy; ask about the reaction.

My Notes / Clues:

© 2026 Demo Study Shop

ALENDRONATE

CLASS: Bisphosphonate

USE: Osteoporosis

MOA: Inhibits bone breakdown by osteoclasts

SE: Esophagitis, GI upset, rare jaw osteonecrosis

NOTES: Fosamax. Empty stomach, full glass of water, stay upright 30 min.

My Notes / Clues:

© 2026 Demo Study Shop

TOPIRAMATE

CLASS: Anticonvulsant

USE: Seizures, migraine prevention

MOA: Blocks sodium channels, enhances GABA

SE: Cognitive slowing, paresthesia, kidney stones, weight loss

NOTES: Topamax. Drink plenty of water. Teratogenic; reduces OC effectiveness at high doses.

My Notes / Clues:

© 2026 Demo Study Shop

TIZANIDINE

CLASS: Central Alpha-2 Agonist (Muscle Relaxant)

USE: Spasticity

MOA: Reduces spinal motor neuron activity

SE: Drowsiness, hypotension, dry mouth, liver injury

NOTES: Zanaflex. Avoid with ciprofloxacin or fluvoxamine. Taper to stop.

My Notes / Clues:

© 2026 Demo Study Shop

DAPAGLIFLOZIN

CLASS: SGLT2 Inhibitor

USE: Type 2 diabetes, heart failure, CKD

MOA: Blocks renal glucose reabsorption

SE: Genital infections, UTIs, dehydration, ketoacidosis

NOTES: Farxiga. Hold before surgery. Stay hydrated.

My Notes / Clues:

© 2026 Demo Study Shop

OXYCODONE / ACETAMINOPHEN

BLACK BOX WARNING

CLASS: Opioid Combination

USE: Moderate to severe pain

MOA: Mu-opioid agonist plus central analgesic

SE: Constipation, sedation, respiratory depression

NOTES: Percocet. Schedule II. Count total daily acetaminophen.

My Notes / Clues:

© 2026 Demo Study Shop

**ATYPICAL
ANTIPSYCHOTIC**

© 2026 Demo Study Shop

**CENTRAL ALPHA-2
AGONIST**

© 2026 Demo Study Shop

**ATYPICAL
ANTIPSYCHOTIC**

© 2026 Demo Study Shop

ANTICONVULSANT

© 2026 Demo Study Shop

BISPHOSPHONATE

© 2026 Demo Study Shop

**1ST-GEN
CEPHALOSPORIN**

© 2026 Demo Study Shop

**OPIOID
COMBINATION**

© 2026 Demo Study Shop

SGLT2 INHIBITOR

© 2026 Demo Study Shop

**CENTRAL ALPHA-2
AGONIST (MUSCLE
RELAXANT)**

© 2026 Demo Study Shop

LOSARTAN / HYDROCHLOROTHIAZIDE

BLACK BOX WARNING

CLASS: ARB + Thiazide

USE: Hypertension

MOA: Blocks angiotensin II receptors plus thiazide diuresis

SE: Dizziness, electrolyte changes

NOTES: Hyzaar. Avoid in pregnancy. Monitor K+.

My Notes / Clues:

© 2026 Demo Study Shop

OLMESARTAN

BLACK BOX WARNING

CLASS: ARB

USE: Hypertension

MOA: Blocks angiotensin II receptors

SE: Dizziness, hyperkalemia, sprue-like diarrhea

NOTES: Benicar. Report severe chronic diarrhea. Avoid in pregnancy.

My Notes / Clues:

© 2026 Demo Study Shop

TESTOSTERONE

BLACK BOX WARNING

CLASS: Androgen

USE: Low testosterone (hypogonadism)

MOA: Replaces testosterone

SE: Acne, polycythemia, raised BP, edema

NOTES: AndroGel / Depo-Testosterone. Gel: cover site and wash hands; avoid skin contact with women/children.

My Notes / Clues:

© 2026 Demo Study Shop

AMITRIPTYLINE

BLACK BOX WARNING

CLASS: Tricyclic Antidepressant

USE: Depression, neuropathic pain, migraine prevention

MOA: Blocks serotonin and norepinephrine reuptake

SE: Sedation, dry mouth, constipation, orthostasis

NOTES: Elavil. Give at bedtime. Overdose is highly cardiotoxic.

My Notes / Clues:

© 2026 Demo Study Shop

FOLIC ACID

CLASS: Vitamin (B9)

USE: Folate deficiency, pregnancy (neural tube prevention), methotrexate

MOA: Needed for DNA synthesis and red cell production

SE: Rarely: GI upset

NOTES: Folvite. Can mask B12 deficiency anemia.

My Notes / Clues:

© 2026 Demo Study Shop

RIVAROXABAN

BLACK BOX WARNING

CLASS: Factor Xa Inhibitor (DOAC)

USE: AFib stroke prevention, DVT/PE

MOA: Directly inhibits factor Xa

SE: Bleeding, bruising

NOTES: Xarelto. 15 and 20 mg doses with the evening meal. Stopping early raises stroke risk.

My Notes / Clues:

© 2026 Demo Study Shop

FENOFIBRATE

CLASS: Fibrate

USE: High triglycerides

MOA: Activates PPAR-alpha, increasing fat breakdown

SE: Myopathy (with statins), gallstones, raised LFTs

NOTES: Tricor. Monitor kidney function.

My Notes / Clues:

© 2026 Demo Study Shop

TRIAMCINOLONE

CLASS: Corticosteroid

USE: Skin inflammation (topical), joint injections

MOA: Reduces inflammation and immune response

SE: Skin thinning (topical), raised glucose (injection)

NOTES: Kenalog. Avoid long use on face and skin folds.

My Notes / Clues:

© 2026 Demo Study Shop

PAROXETINE

BLACK BOX WARNING

CLASS: SSRI

USE: Depression, anxiety, PTSD, OCD

MOA: Blocks serotonin reuptake

SE: Sedation, weight gain, sexual dysfunction, anticholinergic effects

NOTES: Paxil. Worst SSRI discontinuation syndrome. Avoid in pregnancy.

My Notes / Clues:

© 2026 Demo Study Shop

ANDROGEN

© 2026 Demo Study Shop

ARB

© 2026 Demo Study Shop

ARB + THIAZIDE

© 2026 Demo Study Shop

**FACTOR XA INHIBITOR
(DOAC)**

© 2026 Demo Study Shop

VITAMIN (B9)

© 2026 Demo Study Shop

**TRICYCLIC
ANTIDEPRESSANT**

© 2026 Demo Study Shop

SSRI

© 2026 Demo Study Shop

CORTICOSTEROID

© 2026 Demo Study Shop

FIBRATE

© 2026 Demo Study Shop

FERROUS SULFATE

CLASS: Iron Supplement

USE: Iron-deficiency anemia

MOA: Replaces iron for hemoglobin production

SE: Constipation, dark stools, GI upset

NOTES: Feosol. Best on empty stomach with vitamin C. Keep away from children.

My Notes / Clues:

© 2026 Demo Study Shop

LORATADINE

CLASS: 2nd-Gen Antihistamine

USE: Allergic rhinitis, hives

MOA: Blocks peripheral H1 receptors

SE: Headache, minimal drowsiness

NOTES: Claritin. Once daily; non-sedating for most people.

My Notes / Clues:

© 2026 Demo Study Shop

WARFARIN

BLACK BOX WARNING

CLASS: Vitamin K Antagonist

USE: AFib, DVT/PE, mechanical valves

MOA: Inhibits vitamin K-dependent clotting factors (II, VII, IX, X)

SE: Bleeding, bruising, skin necrosis

NOTES: Coumadin / Jantoven. Monitor INR (usually 2-3). Consistent vitamin K diet. Antidote: vitamin K.

My Notes / Clues:

© 2026 Demo Study Shop

VALACYCLOVIR

CLASS: Antiviral (Nucleoside Analog)

USE: Herpes zoster, genital herpes, cold sores

MOA: Prodrug of acyclovir; inhibits viral DNA polymerase

SE: Headache, nausea, kidney injury

NOTES: Valtrex. Start early (within 72 hr for shingles). Stay hydrated.

My Notes / Clues:

© 2026 Demo Study Shop

BUDESONIDE / FORMOTEROL

CLASS: ICS + LABA

USE: Asthma and COPD maintenance

MOA: Reduces inflammation plus long-acting bronchodilation

SE: Thrush, hoarseness, tremor

NOTES: Symbicort. Rinse mouth after use. Some asthma plans also use it as reliever.

My Notes / Clues:

© 2026 Demo Study Shop

NORGESTIMATE / ETHINYL ESTRADIOL

BLACK BOX WARNING

CLASS: Combined Oral Contraceptive

USE: Pregnancy prevention, acne

MOA: Suppresses ovulation

SE: Spotting, nausea, clots

NOTES: Ortho Tri-Cyclen. Smoking over age 35 greatly raises CV risk.

My Notes / Clues:

© 2026 Demo Study Shop

FLUTICASONE / UMECLIDINIUM / VILANTEROL

CLASS: ICS + LAMA + LABA

USE: COPD and asthma maintenance

MOA: Anti-inflammatory plus two long-acting bronchodilators

SE: Thrush, headache, upper respiratory infection

NOTES: Trelegy Ellipta. One inhalation daily. Rinse mouth after use.

My Notes / Clues:

© 2026 Demo Study Shop

METHOCARBAMOL

CLASS: Skeletal Muscle Relaxant

USE: Acute muscle spasm

MOA: General CNS depression

SE: Drowsiness, dizziness, dark urine

NOTES: Robaxin. Harmless urine color change (brown, black, green).

My Notes / Clues:

© 2026 Demo Study Shop

ACETAMINOPHEN

CLASS: Analgesic / Antipyretic

USE: Pain, fever

MOA: Inhibits prostaglandin synthesis in the CNS

SE: Liver injury in overdose

NOTES: Tylenol. Max 4 g/day (lower with liver disease or alcohol use). Antidote: acetylcysteine.

My Notes / Clues:

© 2026 Demo Study Shop

**VITAMIN K
ANTAGONIST**

© 2026 Demo Study Shop

**2ND-GEN
ANTIHISTAMINE**

© 2026 Demo Study Shop

**IRON
SUPPLEMENT**

© 2026 Demo Study Shop

**COMBINED ORAL
CONTRACEPTIVE**

© 2026 Demo Study Shop

ICS + LABA

© 2026 Demo Study Shop

**ANTIVIRAL
(NUCLEOSIDE
ANALOG)**

© 2026 Demo Study Shop

**ANALGESIC /
ANTIPYRETIC**

© 2026 Demo Study Shop

**SKELETAL MUSCLE
RELAXANT**

© 2026 Demo Study Shop

ICS + LAMA + LABA

© 2026 Demo Study Shop

DILTIAZEM

CLASS: Calcium Channel Blocker (Non-DHP)

USE: AFib rate control, angina, hypertension

MOA: Slows AV-node conduction; relaxes vessels

SE: Bradycardia, hypotension, edema, constipation

NOTES: Cardizem. Hold for low HR/BP. Many CYP3A4 interactions.

My Notes / Clues:

© 2026 Demo Study Shop

THYROID (DESICCATED)

BLACK BOX WARNING

CLASS: Thyroid Hormone

USE: Hypothyroidism

MOA: Natural T3 and T4 from porcine thyroid

SE: Palpitations, tremor, insomnia (excess)

NOTES: Armour Thyroid. Potency can vary more than levothyroxine.

My Notes / Clues:

© 2026 Demo Study Shop

BACLOFEN

CLASS: GABA-B Agonist (Muscle Relaxant)

USE: Spasticity (MS, spinal cord injury)

MOA: Activates GABA-B receptors in the spinal cord

SE: Drowsiness, weakness, dizziness

NOTES: Lioresal. Never stop abruptly (hallucinations, seizures).

My Notes / Clues:

© 2026 Demo Study Shop

NAPROXEN

BLACK BOX WARNING

CLASS: NSAID

USE: Pain, arthritis, gout, dysmenorrhea

MOA: Non-selectively inhibits COX

SE: GI bleeding, kidney injury, raised BP

NOTES: Aleve / Naprosyn. Take with food. Twice-daily dosing.

My Notes / Clues:

© 2026 Demo Study Shop

SUMATRIPTAN

CLASS: Triptan (5-HT1B/1D Agonist)

USE: Acute migraine

MOA: Constricts cranial blood vessels

SE: Chest tightness, tingling, dizziness

NOTES: Imitrex. Avoid with CAD or uncontrolled hypertension. Not for prevention.

My Notes / Clues:

© 2026 Demo Study Shop

PREDNISOLONE

CLASS: Systemic Corticosteroid

USE: Asthma flares, inflammation (liquid for children)

MOA: Suppresses inflammation and immune response

SE: Hyperglycemia, mood changes, insomnia

NOTES: Millipred / Orapred. Take with food in the morning.

My Notes / Clues:

© 2026 Demo Study Shop

ESOMEPRAZOLE

CLASS: Proton Pump Inhibitor

USE: GERD, erosive esophagitis, H. pylori

MOA: Irreversibly blocks the gastric proton pump

SE: Headache, diarrhea; long-term low Mg, fractures

NOTES: Nexium. Take at least 1 hr before a meal.

My Notes / Clues:

© 2026 Demo Study Shop

MIRTAZAPINE

BLACK BOX WARNING

CLASS: Tetracyclic Antidepressant

USE: Depression

MOA: Blocks alpha-2 receptors, increasing serotonin and NE

SE: Sedation, increased appetite, weight gain

NOTES: Remeron. Bedtime dosing. Useful when sleep and appetite are poor.

My Notes / Clues:

© 2026 Demo Study Shop

LEVETIRACETAM

CLASS: Anticonvulsant

USE: Focal and generalized seizures

MOA: Binds synaptic vesicle protein SV2A

SE: Drowsiness, irritability, mood changes

NOTES: Keppra. Few drug interactions. Adjust dose in kidney disease.

My Notes / Clues:

© 2026 Demo Study Shop

**GABA-B AGONIST
(MUSCLE RELAXANT)**

© 2026 Demo Study Shop

**THYROID
HORMONE**

© 2026 Demo Study Shop

**CALCIUM CHANNEL
BLOCKER (NON-DHP)**

© 2026 Demo Study Shop

**SYSTEMIC
CORTICOSTEROID**

© 2026 Demo Study Shop

**TRIPTAN (5-HT_{1B/1D}
AGONIST)**

© 2026 Demo Study Shop

NSAID

© 2026 Demo Study Shop

ANTICONVULSANT

© 2026 Demo Study Shop

**TETRACYCLIC
ANTIDEPRESSANT**

© 2026 Demo Study Shop

**PROTON PUMP
INHIBITOR**

© 2026 Demo Study Shop

NORETHINDRONE

CLASS: Progestin-Only Contraceptive

USE: Pregnancy prevention (incl. breastfeeding)

MOA: Thickens cervical mucus

SE: Irregular bleeding, headache

NOTES: Camila / Errin. Take at the same time daily; a 3-hr delay needs backup.

My Notes / Clues:

© 2026 Demo Study Shop

TADALAFIL

CLASS: PDE-5 Inhibitor

USE: Erectile dysfunction, BPH, pulmonary hypertension

MOA: Increases cGMP, relaxing smooth muscle

SE: Headache, flushing, back pain

NOTES: Cialis. Never with nitrates (severe hypotension). Lasts up to 36 hr.

My Notes / Clues:

© 2026 Demo Study Shop

NIFEDIPINE

CLASS: Calcium Channel Blocker (DHP)

USE: Hypertension, angina, Raynaud's

MOA: Relaxes vascular smooth muscle

SE: Edema, flushing, headache, reflex tachycardia

NOTES: Procardia / Adalat CC. Use extended-release; do not crush.

My Notes / Clues:

© 2026 Demo Study Shop

PROGESTERONE

BLACK BOX WARNING

CLASS: Progestin

USE: Endometrial protection with estrogen, amenorrhea

MOA: Replaces progesterone

SE: Drowsiness, dizziness, breast tenderness

NOTES: Prometrium. Take at bedtime. Contains peanut oil.

My Notes / Clues:

© 2026 Demo Study Shop

INSULIN DEGLUDEC

HIGH-ALERT MEDICATION

CLASS: Ultra-Long-Acting Insulin

USE: Type 1 and 2 diabetes (basal)

MOA: Replaces basal insulin; lasts over 42 hr

SE: Hypoglycemia, weight gain

NOTES: Tresiba. Once daily, flexible timing (8 hr apart minimum).

My Notes / Clues:

© 2026 Demo Study Shop

CYANOCOBALAMIN (VITAMIN B12)

CLASS: Vitamin

USE: B12 deficiency, pernicious anemia

MOA: Needed for red cell production and nerve function

SE: Rare: injection-site pain, hypokalemia early in treatment

NOTES: Nascobal. Pernicious anemia needs injection or nasal form.

My Notes / Clues:

© 2026 Demo Study Shop

HYDRALAZINE

CLASS: Direct Arterial Vasodilator

USE: Hypertension, heart failure (with nitrates)

MOA: Relaxes arteriolar smooth muscle

SE: Reflex tachycardia, headache, lupus-like syndrome

NOTES: Apresoline. Often given with a beta blocker.

My Notes / Clues:

© 2026 Demo Study Shop

ADALIMUMAB

BLACK BOX WARNING

CLASS: TNF-Alpha Inhibitor

USE: RA, psoriasis, Crohn's, ulcerative colitis

MOA: Monoclonal antibody that neutralizes TNF-alpha

SE: Serious infections, injection-site reactions, lymphoma

NOTES: Humira. Screen for TB and hepatitis B first. No live vaccines.

My Notes / Clues:

© 2026 Demo Study Shop

AZELASTINE

CLASS: Intranasal Antihistamine

USE: Allergic and nonallergic rhinitis

MOA: Blocks H1 receptors in the nose

SE: Bitter taste, drowsiness, nasal irritation

NOTES: Astelin / Astepro. Tilt head down while spraying to reduce bitter taste.

My Notes / Clues:

© 2026 Demo Study Shop

**CALCIUM CHANNEL
BLOCKER (DHP)**

© 2026 Demo Study Shop

PDE-5 INHIBITOR

© 2026 Demo Study Shop

**PROGESTIN-ONLY
CONTRACEPTIVE**

© 2026 Demo Study Shop

VITAMIN

© 2026 Demo Study Shop

**ULTRA-LONG-ACTING
INSULIN**

© 2026 Demo Study Shop

PROGESTIN

© 2026 Demo Study Shop

**INTRANASAL
ANTIHISTAMINE**

© 2026 Demo Study Shop

**TNF-ALPHA
INHIBITOR**

© 2026 Demo Study Shop

**DIRECT ARTERIAL
VASODILATOR**

© 2026 Demo Study Shop

SITAGLIPTIN

CLASS: DPP-4 Inhibitor

USE: Type 2 diabetes

MOA: Increases incretin levels (GLP-1)

SE: Headache, nasopharyngitis, pancreatitis

NOTES: Januvia. Weight neutral. Adjust dose in kidney disease.

My Notes / Clues:

© 2026 Demo Study Shop

GLIMEPIRIDE

HIGH-ALERT MEDICATION

CLASS: Sulfonylurea

USE: Type 2 diabetes

MOA: Stimulates insulin release from beta cells

SE: Hypoglycemia, weight gain

NOTES: Amaryl. Take with breakfast. Caution in older adults.

My Notes / Clues:

© 2026 Demo Study Shop

CHLORTHALIDONE

CLASS: Thiazide-Like Diuretic

USE: Hypertension, edema

MOA: Blocks sodium reabsorption in the distal tubule

SE: Hypokalemia, hyponatremia, hyperuricemia

NOTES: Thalitone. Longer-acting than HCTZ. Take in the morning.

My Notes / Clues:

© 2026 Demo Study Shop

OXYBUTYNIN

CLASS: Antimuscarinic (Bladder)

USE: Overactive bladder

MOA: Blocks muscarinic receptors in the bladder

SE: Dry mouth, constipation, confusion in older adults

NOTES: Ditropan. Anticholinergic; avoid in older adults when possible (Beers).

My Notes / Clues:

© 2026 Demo Study Shop

PIOGLITAZONE

BLACK BOX WARNING

CLASS: Thiazolidinedione

USE: Type 2 diabetes

MOA: Improves insulin sensitivity via PPAR-gamma

SE: Edema, weight gain, heart failure, fractures

NOTES: Actos. Avoid in heart failure. Monitor weight and LFTs.

My Notes / Clues:

© 2026 Demo Study Shop

KETOCONAZOLE

BLACK BOX WARNING

CLASS: Azole Antifungal

USE: Fungal skin infections, dandruff (topical)

MOA: Inhibits ergosterol synthesis

SE: Topical: irritation. Oral: liver toxicity, QT prolongation

NOTES: Nizoral. Oral use is limited due to hepatotoxicity; many interactions.

My Notes / Clues:

© 2026 Demo Study Shop

IRBESARTAN

BLACK BOX WARNING

CLASS: ARB

USE: Hypertension, diabetic nephropathy

MOA: Blocks angiotensin II receptors

SE: Dizziness, hyperkalemia

NOTES: Avapro. Avoid in pregnancy.

My Notes / Clues:

© 2026 Demo Study Shop

HYDROXYCHLOROQUINE

CLASS: Antimalarial / DMARD

USE: Lupus, rheumatoid arthritis, malaria

MOA: Modulates immune response

SE: Retinal toxicity, GI upset, QT prolongation

NOTES: Plaquenil. Baseline and yearly eye exams.

My Notes / Clues:

© 2026 Demo Study Shop

SULFAMETHOXAZOLE / TRIMETHOPRIM

CLASS: Sulfonamide Antibiotic

USE: UTI, MRSA skin infections, PCP

MOA: Blocks bacterial folate synthesis

SE: Rash, SJS, hyperkalemia, photosensitivity

NOTES: Bactrim. Drink plenty of water. Check sulfa allergy. Raises warfarin effect.

My Notes / Clues:

© 2026 Demo Study Shop

**THIAZIDE-LIKE
DIURETIC**

© 2026 Demo Study Shop

SULFONYLUREA

© 2026 Demo Study Shop

DPP-4 INHIBITOR

© 2026 Demo Study Shop

AZOLE ANTIFUNGAL

© 2026 Demo Study Shop

THIAZOLIDINEDIONE

© 2026 Demo Study Shop

**ANTIMUSCARINIC
(BLADDER)**

© 2026 Demo Study Shop

**SULFONAMIDE
ANTIBIOTIC**

© 2026 Demo Study Shop

**ANTIMALARIAL /
DMARD**

© 2026 Demo Study Shop

ARB

© 2026 Demo Study Shop

BENZONATATE

CLASS: Antitussive

USE: Cough

MOA: Numbs stretch receptors in the lungs

SE: Drowsiness, dizziness; choking if chewed

NOTES: Tessalon. Swallow whole. Fatal overdose in young children.

My Notes / Clues:

© 2026 Demo Study Shop

NITROFURANTOIN

CLASS: Urinary Antibacterial

USE: Uncomplicated UTI

MOA: Damages bacterial DNA in the urine

SE: Nausea, brown urine, rare lung or liver toxicity

NOTES: Macrobid. Take with food. Avoid with poor kidney function.

My Notes / Clues:

© 2026 Demo Study Shop

ISOSORBIDE MONONITRATE

CLASS: Nitrate Vasodilator

USE: Angina prevention

MOA: Releases nitric oxide, dilating veins

SE: Headache, dizziness, hypotension

NOTES: Imdur. Needs a nitrate-free period daily. Never with PDE-5 inhibitors.

My Notes / Clues:

© 2026 Demo Study Shop

METHYLPREDNISOLONE

CLASS: Systemic Corticosteroid

USE: Inflammation, allergic reactions, flares

MOA: Suppresses inflammation and immune response

SE: Hyperglycemia, insomnia, mood changes

NOTES: Medrol. Dose pack tapers over 6 days. Take with food.

My Notes / Clues:

© 2026 Demo Study Shop

TRIAMTERENE / HYDROCHLOROTHIAZIDE

BLACK BOX WARNING

CLASS: K-Sparing + Thiazide Diuretic

USE: Hypertension, edema

MOA: Blocks sodium channels plus thiazide diuresis

SE: Hyperkalemia, dizziness

NOTES: Maxzide / Dyazide. Avoid potassium supplements and salt substitutes.

My Notes / Clues:

© 2026 Demo Study Shop

LEVOCETIRIZINE

CLASS: 2nd-Gen Antihistamine

USE: Allergic rhinitis, hives

MOA: Blocks peripheral H1 receptors

SE: Drowsiness, dry mouth

NOTES: Xyzal. Take in the evening.

My Notes / Clues:

© 2026 Demo Study Shop

CLINDAMYCIN

BLACK BOX WARNING

CLASS: Lincosamide Antibiotic

USE: Skin, dental, anaerobic infections

MOA: Inhibits bacterial protein synthesis (50S)

SE: Diarrhea, C. difficile colitis, rash

NOTES: Cleocin. Report severe or persistent diarrhea.

My Notes / Clues:

© 2026 Demo Study Shop

ATOMOXETINE

BLACK BOX WARNING

CLASS: Non-Stimulant ADHD Medication

USE: ADHD

MOA: Selectively blocks norepinephrine reuptake

SE: Nausea, decreased appetite, suicidal thoughts, liver injury

NOTES: Strattera. Not controlled. Takes several weeks to work.

My Notes / Clues:

© 2026 Demo Study Shop

DEXMETHYLPHENIDATE

BLACK BOX WARNING

CLASS: CNS Stimulant

USE: ADHD

MOA: Blocks dopamine and norepinephrine reuptake

SE: Insomnia, appetite loss, tachycardia

NOTES: Focalin. Schedule II. Morning dosing.

My Notes / Clues:

© 2026 Demo Study Shop

**NITRATE
VASODILATOR**

© 2026 Demo Study Shop

**URINARY
ANTIBACTERIAL**

© 2026 Demo Study Shop

ANTITUSSIVE

© 2026 Demo Study Shop

**2ND-GEN
ANTIHISTAMINE**

© 2026 Demo Study Shop

**K-SPARING + THIAZIDE
DIURETIC**

© 2026 Demo Study Shop

**SYSTEMIC
CORTICOSTEROID**

© 2026 Demo Study Shop

CNS STIMULANT

© 2026 Demo Study Shop

**NON-STIMULANT
ADHD MEDICATION**

© 2026 Demo Study Shop

**LINCOSAMIDE
ANTIBIOTIC**

© 2026 Demo Study Shop

METHOTREXATE

BLACK BOX WARNING

CLASS: Antimetabolite / DMARD

USE: RA, psoriasis, cancers

MOA: Folic acid antagonist

SE: Bone marrow suppression, liver toxicity, mouth sores

NOTES: Trexall / Otrexup. Weekly (not daily) for RA. Folic acid supplement. Avoid pregnancy.

My Notes / Clues:

© 2026 Demo Study Shop

DONEPEZIL

CLASS: Cholinesterase Inhibitor

USE: Alzheimer's dementia

MOA: Increases acetylcholine in the brain

SE: Nausea, diarrhea, bradycardia, vivid dreams

NOTES: Aricept. Slows decline; does not cure.

My Notes / Clues:

© 2026 Demo Study Shop

DESVENLAFAXINE

BLACK BOX WARNING

CLASS: SNRI

USE: Depression

MOA: Blocks serotonin and norepinephrine reuptake

SE: Nausea, sweating, raised BP

NOTES: Pristiq. Taper to stop.

My Notes / Clues:

© 2026 Demo Study Shop

ROPINIROLE

CLASS: Dopamine Agonist

USE: Parkinson's disease, restless legs syndrome

MOA: Stimulates dopamine receptors

SE: Nausea, sleep attacks, impulse-control disorders

NOTES: Requip. Warn about sudden sleepiness while driving.

My Notes / Clues:

© 2026 Demo Study Shop

CLOBETASOL

CLASS: Super-High-Potency Topical Steroid

USE: Severe psoriasis, eczema

MOA: Reduces skin inflammation

SE: Skin thinning, striae, HPA suppression

NOTES: Temovate. Limit to 2 weeks. Avoid face, groin, armpits.

My Notes / Clues:

© 2026 Demo Study Shop

FLUTICASONE / VILANTEROL

CLASS: ICS + LABA

USE: Asthma and COPD maintenance

MOA: Reduces inflammation plus long-acting bronchodilation

SE: Thrush, headache, upper respiratory infection

NOTES: Breo Ellipta. Once daily. Rinse mouth after use.

My Notes / Clues:

© 2026 Demo Study Shop

ALBUTEROL / IPRATROPIUM

CLASS: SABA + SAMA

USE: COPD bronchospasm

MOA: Beta-2 agonist plus anticholinergic bronchodilation

SE: Dry mouth, tremor, tachycardia

NOTES: Combivent / DuoNeb. Avoid spraying into eyes (glaucoma risk).

My Notes / Clues:

© 2026 Demo Study Shop

PHENTERMINE

CLASS: Sympathomimetic Anorectic

USE: Short-term weight loss

MOA: Releases norepinephrine, suppressing appetite

SE: Insomnia, raised BP and HR, dry mouth

NOTES: Adipex-P. Schedule IV. Short-term use. Avoid with heart disease.

My Notes / Clues:

© 2026 Demo Study Shop

OFLOXACIN

CLASS: Fluoroquinolone Antibiotic

USE: Ear and eye infections (drops)

MOA: Inhibits bacterial DNA gyrase

SE: Local irritation, taste changes (drops)

NOTES: Floxin. Systemic fluoroquinolones carry tendon rupture warnings.

My Notes / Clues:

© 2026 Demo Study Shop

SNRI

© 2026 Demo Study Shop

**CHOLINESTERASE
INHIBITOR**

© 2026 Demo Study Shop

**ANTIMETABOLITE /
DMARD**

© 2026 Demo Study Shop

ICS + LABA

© 2026 Demo Study Shop

**SUPER-HIGH-POTENCY
TOPICAL STEROID**

© 2026 Demo Study Shop

DOPAMINE AGONIST

© 2026 Demo Study Shop

**FLUOROQUINOLONE
ANTIBIOTIC**

© 2026 Demo Study Shop

**SYMPATHOMIMETIC
ANORECTIC**

© 2026 Demo Study Shop

SABA + SAMA

© 2026 Demo Study Shop

MUIPIROCIN

CLASS: Topical Antibiotic

USE: Impetigo, MRSA nasal decolonization

MOA: Inhibits bacterial protein synthesis

SE: Burning, itching at site

NOTES: Bactroban. Apply thin layer; wash hands afterward.

My Notes / Clues:

© 2026 Demo Study Shop

VERAPAMIL

CLASS: Calcium Channel Blocker (Non-DHP)

USE: Hypertension, angina, SVT, migraine prevention

MOA: Slows SA/AV node conduction

SE: Constipation, bradycardia, hypotension

NOTES: Calan. Avoid with beta blockers in heart failure. Increase fiber.

My Notes / Clues:

© 2026 Demo Study Shop

OLANZAPINE

BLACK BOX WARNING

CLASS: Atypical Antipsychotic

USE: Schizophrenia, bipolar disorder

MOA: Blocks D2 and 5-HT_{2A} receptors

SE: Major weight gain, hyperglycemia, sedation

NOTES: Zyprexa. Monitor weight, glucose, lipids.

My Notes / Clues:

© 2026 Demo Study Shop

METRONIDAZOLE

BLACK BOX WARNING

CLASS: Nitroimidazole Antimicrobial

USE: C. diff, bacterial vaginosis, trichomoniasis

MOA: Disrupts DNA in anaerobes and protozoa

SE: Metallic taste, nausea, neuropathy

NOTES: Flagyl. No alcohol during and for 3 days after.

My Notes / Clues:

© 2026 Demo Study Shop

PRAZOSIN

CLASS: Alpha-1 Blocker

USE: Hypertension, PTSD nightmares (off-label)

MOA: Blocks alpha-1 receptors, dilating vessels

SE: First-dose syncope, dizziness

NOTES: Minipress. Give the first dose at bedtime.

My Notes / Clues:

© 2026 Demo Study Shop

CIPROFLOXACIN

BLACK BOX WARNING

CLASS: Fluoroquinolone Antibiotic

USE: UTI, GI infections, bone infections

MOA: Inhibits bacterial DNA gyrase

SE: Tendon rupture, neuropathy, QT prolongation, C. diff

NOTES: Cipro. Separate from dairy, antacids, iron by 2 hr.

My Notes / Clues:

© 2026 Demo Study Shop

SILDENAFIL

CLASS: PDE-5 Inhibitor

USE: Erectile dysfunction, pulmonary hypertension

MOA: Increases cGMP, relaxing smooth muscle

SE: Headache, flushing, blue-tinted vision

NOTES: Viagra / Revatio. Never with nitrates. Seek care for erections over 4 hr.

My Notes / Clues:

© 2026 Demo Study Shop

LEVONORGESTREL / ETHINYL ESTRADIOL

BLACK BOX WARNING

CLASS: Combined Oral Contraceptive

USE: Pregnancy prevention

MOA: Suppresses ovulation

SE: Spotting, nausea, clots

NOTES: Aviane / Seasonique. Smoking over age 35 greatly raises CV risk.

My Notes / Clues:

© 2026 Demo Study Shop

EVOLOCUMAB

CLASS: PCSK9 Inhibitor

USE: High LDL, clinical ASCVD

MOA: Antibody that keeps LDL receptors active

SE: Injection-site reactions, nasopharyngitis

NOTES: Repatha. Subcut every 2 weeks or monthly.

My Notes / Clues:

© 2026 Demo Study Shop

**ATYPICAL
ANTIPSYCHOTIC**

© 2026 Demo Study Shop

**CALCIUM CHANNEL
BLOCKER (NON-DHP)**

© 2026 Demo Study Shop

TOPICAL ANTIBIOTIC

© 2026 Demo Study Shop

**FLUOROQUINOLONE
ANTIBIOTIC**

© 2026 Demo Study Shop

**ALPHA-1
BLOCKER**

© 2026 Demo Study Shop

**NITROIMIDAZOLE
ANTIMICROBIAL**

© 2026 Demo Study Shop

PCSK9 INHIBITOR

© 2026 Demo Study Shop

**COMBINED ORAL
CONTRACEPTIVE**

© 2026 Demo Study Shop

PDE-5 INHIBITOR

© 2026 Demo Study Shop

MESALAMINE

CLASS: Aminosalicylate (5-ASA)

USE: Ulcerative colitis

MOA: Reduces inflammation in the colon lining

SE: Headache, abdominal pain, kidney injury

NOTES: Lialda / Asacol. Swallow whole. Monitor kidney function.

My Notes / Clues:

© 2026 Demo Study Shop

MEMANTINE

CLASS: NMDA Receptor Antagonist

USE: Moderate to severe Alzheimer's dementia

MOA: Blocks excess glutamate activity

SE: Dizziness, headache, constipation

NOTES: Namenda. Often combined with donepezil.

My Notes / Clues:

© 2026 Demo Study Shop

ETONOGESTREL / ETHINYL ESTRADIOL

BLACK BOX WARNING

CLASS: Combined Hormonal Contraceptive Ring

USE: Pregnancy prevention

MOA: Suppresses ovulation

SE: Vaginal irritation, headache, clots

NOTES: NuvaRing. Leave in 3 weeks, out 1 week. Smoking over 35 raises CV risk.

My Notes / Clues:

© 2026 Demo Study Shop

DIAZEPAM

BLACK BOX WARNING

CLASS: Benzodiazepine

USE: Anxiety, muscle spasm, alcohol withdrawal, seizures

MOA: Enhances GABA-A activity

SE: Sedation, ataxia, respiratory depression

NOTES: Valium. Schedule IV. Very long half-life; fall risk in older adults.

My Notes / Clues:

© 2026 Demo Study Shop

LITHIUM

BLACK BOX WARNING

CLASS: Mood Stabilizer

USE: Bipolar disorder

MOA: Alters sodium transport and neurotransmission

SE: Tremor, polyuria, thirst, hypothyroidism

NOTES: Lithobid. Level 0.6-1.2 mEq/L. Keep salt and fluid intake steady.

My Notes / Clues:

© 2026 Demo Study Shop

NORTRIPTYLINE

BLACK BOX WARNING

CLASS: Tricyclic Antidepressant

USE: Depression, neuropathic pain

MOA: Blocks norepinephrine and serotonin reuptake

SE: Dry mouth, constipation, orthostasis

NOTES: Pamelor. Fewer anticholinergic effects than amitriptyline.

My Notes / Clues:

© 2026 Demo Study Shop

TIMOLOL

CLASS: Beta Blocker (Eye)

USE: Glaucoma

MOA: Reduces aqueous humor production

SE: Eye stinging, bradycardia, bronchospasm

NOTES: Timoptic. Systemic absorption possible; press on tear duct after drops.

My Notes / Clues:

© 2026 Demo Study Shop

CEFDINIR

CLASS: 3rd-Gen Cephalosporin

USE: Otitis media, sinusitis, pneumonia, skin infections

MOA: Inhibits bacterial cell wall synthesis

SE: Diarrhea, red stools (with iron)

NOTES: Omnicef. Separate from iron and antacids by 2 hr.

My Notes / Clues:

© 2026 Demo Study Shop

MAGNESIUM SALTS

CLASS: Electrolyte Supplement

USE: Low magnesium, constipation, heartburn

MOA: Replaces magnesium

SE: Diarrhea; toxicity in kidney failure

NOTES: Mag-Ox. Separate from many oral antibiotics.

My Notes / Clues:

© 2026 Demo Study Shop

**COMBINED
HORMONAL
CONTRACEPTIVE RING**

© 2026 Demo Study Shop

**NMDA RECEPTOR
ANTAGONIST**

© 2026 Demo Study Shop

**AMINOSALICYLATE
(5-ASA)**

© 2026 Demo Study Shop

**TRICYCLIC
ANTIDEPRESSANT**

© 2026 Demo Study Shop

**MOOD
STABILIZER**

© 2026 Demo Study Shop

BENZODIAZEPINE

© 2026 Demo Study Shop

**ELECTROLYTE
SUPPLEMENT**

© 2026 Demo Study Shop

**3RD-GEN
CEPHALOSPORIN**

© 2026 Demo Study Shop

BETA BLOCKER (EYE)

© 2026 Demo Study Shop

MECLIZINE

CLASS: 1st-Gen Antihistamine (Vertigo)

USE: Vertigo, motion sickness

MOA: Blocks H1 receptors in the vestibular system

SE: Drowsiness, dry mouth

NOTES: Antivert. Take 1 hr before travel. Fall risk in older adults.

My Notes / Clues:

© 2026 Demo Study Shop

CYCLOSPORINE

BLACK BOX WARNING

CLASS: Calcineurin Inhibitor

USE: Transplant rejection prevention, RA, psoriasis

MOA: Inhibits calcineurin, blocking T-cell activation

SE: Nephrotoxicity, hypertension, gum overgrowth

NOTES: Neoral / Sandimmune. Monitor levels. Avoid grapefruit.

My Notes / Clues:

© 2026 Demo Study Shop

FLUCONAZOLE

CLASS: Azole Antifungal

USE: Candida infections, cryptococcal meningitis

MOA: Inhibits fungal ergosterol synthesis

SE: Headache, nausea, liver injury, QT prolongation

NOTES: Diflucan. Many interactions (raises warfarin effect).

My Notes / Clues:

© 2026 Demo Study Shop

RIMEGEPANT

CLASS: CGRP Receptor Antagonist

USE: Acute migraine and prevention

MOA: Blocks CGRP receptors

SE: Nausea

NOTES: Nurtec ODT. Dissolve on the tongue. No vasoconstriction (unlike triptans).

My Notes / Clues:

© 2026 Demo Study Shop

TRETINOIN

CLASS: Topical Retinoid

USE: Acne, photoaging

MOA: Increases skin cell turnover

SE: Dryness, peeling, sun sensitivity

NOTES: Retin-A. Apply at night; use sunscreen. Avoid in pregnancy.

My Notes / Clues:

© 2026 Demo Study Shop

LOVASTATIN

CLASS: Statin

USE: High cholesterol

MOA: Inhibits HMG-CoA reductase

SE: Myalgia, raised LFTs

NOTES: Mevacor. Take with the evening meal. Avoid grapefruit.

My Notes / Clues:

© 2026 Demo Study Shop

DROSPIRENONE / ETHINYL ESTRADIOL

BLACK BOX WARNING

CLASS: Combined Oral Contraceptive

USE: Pregnancy prevention, PMDD, acne

MOA: Suppresses ovulation; drospirenone is anti-mineralocorticoid

SE: Hyperkalemia, clots, spotting

NOTES: Yaz / Yasmin. Check K+ with ACE inhibitors or ARBs. Smoking over 35 raises risk.

My Notes / Clues:

© 2026 Demo Study Shop

TIOTROPIUM

CLASS: LAMA

USE: COPD and asthma maintenance

MOA: Blocks muscarinic receptors in the airways

SE: Dry mouth, urinary retention, constipation

NOTES: Spiriva. Do not swallow capsules; use with the inhaler device.

My Notes / Clues:

© 2026 Demo Study Shop

DOCUSATE

CLASS: Stool Softener

USE: Constipation prevention

MOA: Lets water into the stool

SE: Mild cramping, diarrhea

NOTES: Colace. Does not stimulate the bowel; drink plenty of water.

My Notes / Clues:

© 2026 Demo Study Shop

AZOLE ANTIFUNGAL

© 2026 Demo Study Shop

**CALCINEURIN
INHIBITOR**

© 2026 Demo Study Shop

**1ST-GEN
ANTIHISTAMINE
(VERTIGO)**

© 2026 Demo Study Shop

STATIN

© 2026 Demo Study Shop

TOPICAL RETINOID

© 2026 Demo Study Shop

**CGRP RECEPTOR
ANTAGONIST**

© 2026 Demo Study Shop

STOOL SOFTENER

© 2026 Demo Study Shop

LAMA

© 2026 Demo Study Shop

**COMBINED ORAL
CONTRACEPTIVE**

© 2026 Demo Study Shop

TEMAZEPAM

BLACK BOX WARNING

CLASS: Benzodiazepine (Hypnotic)

USE: Short-term insomnia

MOA: Enhances GABA-A activity

SE: Drowsiness, dizziness, dependence

NOTES: Restoril. Schedule IV. Take 30 min before bed.

My Notes / Clues:

© 2020 Demo Study Shop

BENAZEPRIL

BLACK BOX WARNING

CLASS: ACE Inhibitor

USE: Hypertension

MOA: Blocks conversion of angiotensin I to II

SE: Dry cough, hyperkalemia, angioedema

NOTES: Lotensin. Avoid in pregnancy.

My Notes / Clues:

© 2020 Demo Study Shop

BRIMONIDINE

CLASS: Alpha-2 Agonist (Eye)

USE: Glaucoma, ocular hypertension

MOA: Reduces aqueous humor production

SE: Eye redness, dry mouth, fatigue

NOTES: Alphagan. Not for infants (CNS depression).

My Notes / Clues:

© 2020 Demo Study Shop

SACUBITRIL / VALSARTAN

BLACK BOX WARNING

CLASS: ARNI

USE: Heart failure (HFrEF)

MOA: Neprilysin inhibitor plus ARB

SE: Hypotension, hyperkalemia, angioedema

NOTES: Entresto. 36-hr washout after stopping an ACE inhibitor. Avoid in pregnancy.

My Notes / Clues:

© 2020 Demo Study Shop

VALPROATE / DIVALPROEX

BLACK BOX WARNING

CLASS: Anticonvulsant / Mood Stabilizer

USE: Seizures, bipolar mania, migraine prevention

MOA: Increases GABA; blocks sodium and calcium channels

SE: Liver toxicity, pancreatitis, tremor, weight gain

NOTES: Depakote. Monitor LFTs and levels (50-100 mcg/mL). Highly teratogenic.

My Notes / Clues:

© 2020 Demo Study Shop

NEBIVOLOL

CLASS: Beta-1 Selective Blocker (Vasodilating)

USE: Hypertension

MOA: Blocks beta-1; releases nitric oxide

SE: Headache, fatigue, bradycardia

NOTES: Bystolic. Do not stop abruptly.

My Notes / Clues:

© 2020 Demo Study Shop

ACYCLOVIR

CLASS: Antiviral (Nucleoside Analog)

USE: Herpes simplex, shingles, chickenpox

MOA: Inhibits viral DNA polymerase

SE: Nausea, kidney injury (IV)

NOTES: Zovirax. Hydrate well, especially with IV use.

My Notes / Clues:

© 2020 Demo Study Shop

MORPHINE

BLACK BOX WARNING

CLASS: Opioid Analgesic

USE: Severe pain

MOA: Mu-opioid receptor agonist

SE: Respiratory depression, constipation, hypotension

NOTES: MS Contin. Schedule II. Never crush ER tablets. Antidote: naloxone.

My Notes / Clues:

© 2020 Demo Study Shop

DOXAZOSIN

CLASS: Alpha-1 Blocker

USE: BPH, hypertension

MOA: Blocks alpha-1 receptors

SE: First-dose syncope, dizziness

NOTES: Cardura. First dose at bedtime.

My Notes / Clues:

© 2020 Demo Study Shop

**ALPHA-2 AGONIST
(EYE)**

© 2026 Demo Study Shop

ACE INHIBITOR

© 2026 Demo Study Shop

**BENZODIAZEPINE
(HYPNOTIC)**

© 2026 Demo Study Shop

**BETA-1 SELECTIVE
BLOCKER
(VASODILATING)**

© 2026 Demo Study Shop

**ANTICONVULSANT /
MOOD STABILIZER**

© 2026 Demo Study Shop

ARNI

© 2026 Demo Study Shop

**ALPHA-1
BLOCKER**

© 2026 Demo Study Shop

OPIOID ANALGESIC

© 2026 Demo Study Shop

**ANTIVIRAL
(NUCLEOSIDE
ANALOG)**

© 2026 Demo Study Shop

AMLODIPINE / BENAZEPRIL

BLACK BOX WARNING

CLASS: CCB + ACE Inhibitor

USE: Hypertension

MOA: Vasodilation plus angiotensin blockade

SE: Edema, cough, dizziness

NOTES: Lotrel. Avoid in pregnancy.

My Notes / Clues:

© 2020 Demo Study Shop

NIRMATRELVIR / RITONAVIR

BLACK BOX WARNING

CLASS: SARS-CoV-2 Protease Inhibitor + Booster

USE: Mild-moderate COVID-19 in high-risk patients

MOA: Blocks viral protease; ritonavir boosts levels

SE: Altered taste, diarrhea

NOTES: Paxlovid. Start within 5 days of symptoms. Check all drug interactions.

My Notes / Clues:

© 2020 Demo Study Shop

EMTRICITABINE / TENOFOVIR

BLACK BOX WARNING

CLASS: NRTI Combination

USE: HIV treatment, HIV PrEP

MOA: Blocks HIV reverse transcriptase

SE: Nausea, kidney injury, bone loss

NOTES: Truvada / Descovy. Confirm HIV-negative before PrEP. HBV flare on stopping.

My Notes / Clues:

© 2020 Demo Study Shop

BUDESONIDE

CLASS: Corticosteroid

USE: Asthma (inhaled), Crohn's disease (oral)

MOA: Local anti-inflammatory effect

SE: Thrush (inhaled), headache

NOTES: Pulmicort / Entocort. Rinse mouth after inhaled use.

My Notes / Clues:

© 2020 Demo Study Shop

TACROLIMUS

BLACK BOX WARNING

CLASS: Calcineurin Inhibitor

USE: Organ transplant rejection prevention

MOA: Inhibits calcineurin, blocking T-cell activation

SE: Nephrotoxicity, tremor, hyperglycemia, hyperkalemia

NOTES: Prograf. Monitor trough levels. Avoid grapefruit.

My Notes / Clues:

© 2020 Demo Study Shop

DUPILUMAB

CLASS: IL-4 / IL-13 Blocker (Monoclonal Antibody)

USE: Eczema, asthma, chronic sinusitis with polyps

MOA: Blocks IL-4 and IL-13 signaling

SE: Conjunctivitis, injection-site reactions

NOTES: Dupixent. Subcut every 2 weeks. Avoid live vaccines.

My Notes / Clues:

© 2020 Demo Study Shop

ENALAPRIL

BLACK BOX WARNING

CLASS: ACE Inhibitor

USE: Hypertension, heart failure

MOA: Blocks conversion of angiotensin I to II

SE: Dry cough, hyperkalemia, angioedema

NOTES: Vasotec. IV form: enalaprilat. Avoid in pregnancy.

My Notes / Clues:

© 2020 Demo Study Shop

ANASTROZOLE

CLASS: Aromatase Inhibitor

USE: Hormone-receptor-positive breast cancer (postmenopausal)

MOA: Blocks estrogen production

SE: Hot flashes, joint pain, bone loss

NOTES: Arimidex. Monitor bone density.

My Notes / Clues:

© 2020 Demo Study Shop

MELATONIN

CLASS: Sleep Aid / Supplement

USE: Insomnia, jet lag

MOA: Mimics the natural sleep hormone

SE: Drowsiness, headache, vivid dreams

NOTES: Various. Take 30-60 min before bed. Supplements are not FDA-regulated for dose accuracy.

My Notes / Clues:

© 2020 Demo Study Shop

NRTI COMBINATION

© 2026 Demo Study Shop

**SARS-COV-2 PROTEASE
INHIBITOR + BOOSTER**

© 2026 Demo Study Shop

CCB + ACE INHIBITOR

© 2026 Demo Study Shop

**IL-4 / IL-13 BLOCKER
(MONOCLONAL
ANTIBODY)**

© 2026 Demo Study Shop

**CALCINEURIN
INHIBITOR**

© 2026 Demo Study Shop

CORTICOSTEROID

© 2026 Demo Study Shop

**SLEEP AID /
SUPPLEMENT**

© 2026 Demo Study Shop

**AROMATASE
INHIBITOR**

© 2026 Demo Study Shop

ACE INHIBITOR

© 2026 Demo Study Shop

TORSEMIDE

CLASS: Loop Diuretic

USE: Edema, heart failure, hypertension

MOA: Blocks Na/K/2Cl in the loop of Henle

SE: Hypokalemia, dehydration, dizziness

NOTES: Demadex. Longer-acting than furosemide. Monitor K⁺ and weight.

My Notes / Clues:

© 2020 Demo Study Shop

POLYETHYLENE GLYCOL 3350

CLASS: Osmotic Laxative

USE: Constipation, bowel prep

MOA: Draws water into the colon

SE: Bloating, cramping, diarrhea

NOTES: MiraLAX. Mix in 4-8 oz of fluid. Works in 1-3 days.

My Notes / Clues:

© 2020 Demo Study Shop

OSMOTIC LAXATIVE

© 2026 Demo Study Shop

LOOP DIURETIC

© 2026 Demo Study Shop

EDUCATIONAL USE ONLY

NOT A CLINICAL REFERENCE

This study aid is a learning and exam-review tool for nursing and pharmacy students. It must not be used to make patient-care decisions, including dosing, administration, or monitoring.

ALWAYS VERIFY

Drug information, boxed (black box) warnings, and high-alert designations change over time. Before applying anything in practice, check the current FDA-approved labeling, your facility's policies, and an up-to-date drug reference. Normal lab ranges vary between laboratories; use the ranges your program or facility provides.

RECOMMENDED REFERENCES FOR VERIFICATION

- FDA drug labels: DailyMed (dailymed.nlm.nih.gov)
- ISMP List of High-Alert Medications ([ismp.org](https://www.ismp.org))
- A current nursing drug handbook or your program's required drug reference
- Your course textbook and instructor guidance

ACCURACY

Content was reviewed for accuracy at the time of publication, but errors or omissions are possible. If you spot one, please let us know so it can be corrected in the next update.

TERMS OF USE

For personal study use by the purchaser only. You may print as many copies as you need for your own studying. Do not resell, redistribute, upload, or share this file.