



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Applied stochastic processes


Jie Xiong

Department of Mathematics
Southern University of Science and Technology
Shenzhen, China, 518055

Math 208, Spring 2023



群聊：应随 2023



该二维码7天内(2月20日前)有效，重新进入将更新



Outline for this semester

- 1 Revisiting of probability



Outline for this semester

- 1 Revisiting of probability
- 2 Conditional Expectation



Outline for this semester

- ① Revisiting of probability
- ② Conditional Expectation
- ③ Discrete Time Markov Chains (I)



Outline for this semester

- ① Revisiting of probability
- ② Conditional Expectation
- ③ Discrete Time Markov Chains (I)
- ④ Discrete Time Markov Chains (II)



Outline for this semester

- ① Revisiting of probability
- ② Conditional Expectation
- ③ Discrete Time Markov Chains (I)
- ④ Discrete Time Markov Chains (II)
- ⑤ Poisson Processes

Outline for this semester

- ① Revisiting of probability
- ② Conditional Expectation
- ③ Discrete Time Markov Chains (I)
- ④ Discrete Time Markov Chains (II)
- ⑤ Poisson Processes
- ⑥ Continuous Time Markov Chains



Outline for this semester

- ① Revisiting of probability
- ② Conditional Expectation
- ③ Discrete Time Markov Chains (I)
- ④ Discrete Time Markov Chains (II)
- ⑤ Poisson Processes
- ⑥ Continuous Time Markov Chains
- ⑦ Brownian Motion Processes



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Chapter 1. Revisiting of probability



Chapter 1. Revisiting of probability

1.1. Basic probabilities



Chapter 1. Revisiting of probability

1.1. Basic probabilities

S =sample space= all possible outcomes of an experiment.



Chapter 1. Revisiting of probability

1.1. Basic probabilities

S =sample space= all possible outcomes of an experiment.

P : probability measure which assign a number to each event

$E \subset S$.



Chapter 1. Revisiting of probability

1.1. Basic probabilities

S =sample space= all possible outcomes of an experiment.

P : probability measure which assign a number to each event

$E \subset S$.

① $0 \leq P(E) \leq 1$



Chapter 1. Revisiting of probability

1.1. Basic probabilities

S =sample space= all possible outcomes of an experiment.

P : probability measure which assign a number to each event

$E \subset S$.

① $0 \leq P(E) \leq 1$

② $P(S) = 1$



Chapter 1. Revisiting of probability

1.1. Basic probabilities

S =sample space= all possible outcomes of an experiment.

P : probability measure which assign a number to each event

$E \subset S$.

- ① $0 \leq P(E) \leq 1$
- ② $P(S) = 1$
- ③ If E_1, E_2, \dots are disjoint, then

$$P(\cup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} P(E_n).$$



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Random variable (r.v.): $X : S \rightarrow \mathbb{R}$.



Random variable (r.v.): $X : S \rightarrow \mathbb{R}$.

Random vector: $X = (X_1, \dots, X_d) : S \rightarrow \mathbb{R}^d$.



Random variable (r.v.): $X : S \rightarrow \mathbb{R}$.

Random vector: $X = (X_1, \dots, X_d) : S \rightarrow \mathbb{R}^d$.

Distribution function (d.f.) of X :

$$F(x) = P(X \leq x), \quad x \in \mathbb{R}.$$



Random variable (r.v.): $X : S \rightarrow \mathbb{R}$.

Random vector: $X = (X_1, \dots, X_d) : S \rightarrow \mathbb{R}^d$.

Distribution function (d.f.) of X :

$$F(x) = P(X \leq x), \quad x \in \mathbb{R}.$$

Joint distribution function of (X_1, \dots, X_d) :

$$F(x_1, \dots, x_d) = P(X_1 \leq x_1, \dots, X_d \leq x_d).$$



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Two types of r.v.'s: Discrete and continuous.



Two types of r.v.'s: Discrete and continuous.

If X takes only countable many possible values x_1, x_2, \dots , then X is a discrete r.v.



Two types of r.v.'s: Discrete and continuous.

If X takes only countable many possible values x_1, x_2, \dots , then X is a discrete r.v.

Probability mass function (pmf):

$$p(x) = P(X = x), \quad x = x_1, x_2, \dots$$



Two types of r.v.'s: Discrete and continuous.

If X takes only countable many possible values x_1, x_2, \dots , then X is a discrete r.v.

Probability mass function (pmf):

$$p(x) = P(X = x), \quad x = x_1, x_2, \dots$$

- $0 \leq p(x) \leq 1$



Two types of r.v.'s: Discrete and continuous.

If X takes only countable many possible values x_1, x_2, \dots , then X is a discrete r.v.

Probability mass function (pmf):

$$p(x) = P(X = x), \quad x = x_1, x_2, \dots$$

- $0 \leq p(x) \leq 1$
- $\sum_x p(x) = 1$



For any $A \subset \mathbb{R}$,

$$P(X \in A) = \sum_{x_i \in A} p(x_i).$$



For any $A \subset \mathbb{R}$,

$$P(X \in A) = \sum_{x_i \in A} p(x_i).$$

Expectation (what do you expect) of X is

$$\mu = \mathbb{E}(X) = \sum_x xp(x).$$



Variance (average square distance to μ) of X is

$$\sigma^2 = V(X) = \mathbb{E}((X - \mu)^2) = \sum_x (x - \mu)^2 p(x).$$



Variance (average square distance to μ) of X is

$$\sigma^2 = V(X) = \mathbb{E}((X - \mu)^2) = \sum_x (x - \mu)^2 p(x).$$

Another formula

$$\sigma^2 = \mathbb{E}(X^2) - \mu^2.$$



X is a continuous r.v. if

$$P(X \leq x) = F(x) = \int_{-\infty}^x f(y)dy$$

for a suitable function f , which is called the probability density function (p.d.f.) of X .



X is a continuous r.v. if

$$P(X \leq x) = F(x) = \int_{-\infty}^x f(y)dy$$

for a suitable function f , which is called the probability density function (p.d.f.) of X .

- $f(x) \geq 0$

X is a continuous r.v. if

$$P(X \leq x) = F(x) = \int_{-\infty}^x f(y)dy$$

for a suitable function f , which is called the probability density function (p.d.f.) of X .

- $f(x) \geq 0$



$$\int_{-\infty}^{\infty} f(x)dx = 1.$$



Expectation of X is

$$\mu = \mathbb{E}(X) = \int_{-\infty}^{\infty} x f(x) dx.$$



Expectation of X is

$$\mu = \mathbb{E}(X) = \int_{-\infty}^{\infty} x f(x) dx.$$

Variance of X is

$$\sigma^2 = V(X) = \mathbb{E}((X - \mu)^2) = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx.$$



If X has pdf f_X and g is 1-to-1, then $Y = g(X)$ has pdf



If X has pdf f_X and g is 1-to-1, then $Y = g(X)$ has pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|.$$



If X has pdf f_X and g is 1-to-1, then $Y = g(X)$ has pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|.$$

If $X = (X_1, X_2)$ has joint pdf $f_X(x)$ and $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is 1-to-1, then $(Y_1, Y_2) = Y = g(X)$ has joint pdf f_Y given by

If X has pdf f_X and g is 1-to-1, then $Y = g(X)$ has pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|.$$

If $X = (X_1, X_2)$ has joint pdf $f_X(x)$ and $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is 1-to-1, then $(Y_1, Y_2) = Y = g(X)$ has joint pdf f_Y given by

$$f_Y(y) = f_X(g^{-1}(y)) |J|^{-1}$$

If X has pdf f_X and g is 1-to-1, then $Y = g(X)$ has pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|.$$

If $X = (X_1, X_2)$ has joint pdf $f_X(x)$ and $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is 1-to-1, then $(Y_1, Y_2) = Y = g(X)$ has joint pdf f_Y given by

$$f_Y(y) = f_X(g^{-1}(y)) |J|^{-1}$$

where J is the Jacobian determinant

If X has pdf f_X and g is 1-to-1, then $Y = g(X)$ has pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|.$$

If $X = (X_1, X_2)$ has joint pdf $f_X(x)$ and $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is 1-to-1, then $(Y_1, Y_2) = Y = g(X)$ has joint pdf f_Y given by

$$f_Y(y) = f_X(g^{-1}(y)) |J|^{-1}$$

where J is the Jacobian determinant

$$J = \begin{vmatrix} \frac{\partial g_1}{\partial x_1} & \frac{\partial g_1}{\partial x_2} \\ \frac{\partial g_2}{\partial x_1} & \frac{\partial g_2}{\partial x_2} \end{vmatrix}$$



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

1.2. Independence



1.2. Independence

Two events A , B are independent if

$$P(AB) = P(A)P(B).$$



1.2. Independence

Two events A , B are independent if

$$P(AB) = P(A)P(B).$$

n events A_1, \dots, A_n are independent if

1.2. Independence

Two events A , B are independent if

$$P(AB) = P(A)P(B).$$

n events A_1, \dots, A_n are independent if for any $2 \leq k \leq n$,
 $\{i_1, \dots, i_k\} \subset \{1, \dots, n\}$,

$$P(A_{i_1} \cdots A_{i_k}) = P(A_{i_1}) \cdots P(A_{i_k}).$$



Example

If $n = 3$, there are 4 equations need to be verified:



Example

If $n = 3$, there are 4 equations need to be verified:

$$P(AB) = P(A)P(B),$$



Example

If $n = 3$, there are 4 equations need to be verified:

$$P(AB) = P(A)P(B),$$

$$P(AC) = P(A)P(C),$$



Example

If $n = 3$, there are 4 equations need to be verified:

$$P(AB) = P(A)P(B),$$

$$P(AC) = P(A)P(C),$$

$$P(BC) = P(B)P(C),$$



Example

If $n = 3$, there are 4 equations need to be verified:

$$P(AB) = P(A)P(B),$$

$$P(AC) = P(A)P(C),$$

$$P(BC) = P(B)P(C),$$

$$P(ABC) = P(A)P(B)P(C).$$



Two r.v.'s are independent if

$$P(X \leq x, Y \leq y) = P(X \leq x)P(Y \leq y),$$



Two r.v.'s are independent if

$$P(X \leq x, Y \leq y) = P(X \leq x)P(Y \leq y),$$

namely,

$$F(x, y) = F_X(x)F_Y(y).$$

Two r.v.'s are independent if

$$P(X \leq x, Y \leq y) = P(X \leq x)P(Y \leq y),$$

namely,

$$F(x, y) = F_X(x)F_Y(y).$$

- Discrete r.v.: $p(x, y) = p_X(x)p_Y(y)$.

Two r.v.'s are independent if

$$P(X \leq x, Y \leq y) = P(X \leq x)P(Y \leq y),$$

namely,

$$F(x, y) = F_X(x)F_Y(y).$$

- Discrete r.v.: $p(x, y) = p_X(x)p_Y(y)$.
- Continuous r.v.: $f(x, y) = f_X(x)f_Y(y)$.



Conditional probability: If A , B are two events and $P(B) > 0$, then



Conditional probability: If A , B are two events and $P(B) > 0$, then

$$P(A|B) = \frac{P(AB)}{P(B)}.$$



Conditional probability: If A , B are two events and $P(B) > 0$, then

$$P(A|B) = \frac{P(AB)}{P(B)}.$$

If A and B are independent, then



Conditional probability: If A , B are two events and $P(B) > 0$, then

$$P(A|B) = \frac{P(AB)}{P(B)}.$$

If A and B are independent, then

$$P(A|B) = P(A).$$



Conditional probability: If A , B are two events and $P(B) > 0$, then

$$P(A|B) = \frac{P(AB)}{P(B)}.$$

If A and B are independent, then

$$P(A|B) = P(A).$$

Equivalently,



Conditional probability: If A , B are two events and $P(B) > 0$, then

$$P(A|B) = \frac{P(AB)}{P(B)}.$$

If A and B are independent, then

$$P(A|B) = P(A).$$

Equivalently,

$$P(AB) = P(A)P(B).$$



1.3. Moment generating function



1.3. Moment generating function

The moment generating function (m.g.f.) of X is



1.3. Moment generating function

The moment generating function (m.g.f.) of X is

$$\phi(t) = \mathbb{E}(e^{tX})$$



1.3. Moment generating function

The moment generating function (m.g.f.) of X is

$$\phi(t) = \mathbb{E}(e^{tX}) = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ discrete} \end{cases}$$



1.3. Moment generating function

The moment generating function (m.g.f.) of X is

$$\phi(t) = \mathbb{E}(e^{tX}) = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ discrete} \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ continuous.} \end{cases}$$



ϕ is the mgf since we can obtain all moments from it.



ϕ is the mgf since we can obtain all moments from it. In fact, by Taylor's formula, we have

$$\phi(t) = \sum_{n=0}^{\infty} \frac{\phi^{(n)}(0)}{n!} t^n,$$



ϕ is the mgf since we can obtain all moments from it. In fact, by Taylor's formula, we have

$$\phi(t) = \sum_{n=0}^{\infty} \frac{\phi^{(n)}(0)}{n!} t^n,$$

and

$$\phi(t) = E(e^{tX})$$

ϕ is the mgf since we can obtain all moments from it. In fact, by Taylor's formula, we have

$$\phi(t) = \sum_{n=0}^{\infty} \frac{\phi^{(n)}(0)}{n!} t^n,$$

and

$$\phi(t) = E(e^{tX}) = \mathbb{E} \sum_{n=0}^{\infty} \frac{(tX)^n}{n!}$$

ϕ is the mgf since we can obtain all moments from it. In fact, by Taylor's formula, we have

$$\phi(t) = \sum_{n=0}^{\infty} \frac{\phi^{(n)}(0)}{n!} t^n,$$

and

$$\phi(t) = E(e^{tX}) = \mathbb{E} \sum_{n=0}^{\infty} \frac{(tX)^n}{n!} = \sum_{n=0}^{\infty} \frac{\mathbb{E}(X^n)}{n!} t^n.$$

ϕ is the mgf since we can obtain all moments from it. In fact, by Taylor's formula, we have

$$\phi(t) = \sum_{n=0}^{\infty} \frac{\phi^{(n)}(0)}{n!} t^n,$$

and

$$\phi(t) = E(e^{tX}) = \mathbb{E} \sum_{n=0}^{\infty} \frac{(tX)^n}{n!} = \sum_{n=0}^{\infty} \frac{\mathbb{E}(X^n)}{n!} t^n.$$

Thus,

$$\mathbb{E}(X^n) = \phi^{(n)}(0).$$



Example

If $X \sim N(\mu, \sigma^2)$, find its mgf.



Example

If $X \sim N(\mu, \sigma^2)$, find its mgf.

Solution: Do it on board!



Example

If $X \sim P(\lambda)$, find its mgf.



Example

If $X \sim P(\lambda)$, find its mgf.

Solution: Do it on board!



Proposition

If X and Y are independent and a, b are constants, then



Proposition

If X and Y are independent and a, b are constants, then

$$\phi_{aX+bY}(t) = \phi_X(at)\phi_Y(bt).$$

Proof: Do it on board!



Proposition

If X and Y are independent and a, b are constants, then

$$\phi_{aX+bY}(t) = \phi_X(at)\phi_Y(bt).$$

Proof: Do it on board!



Example

If X and Y are independent standard normal and a, b are constants, show that $aX + bY$ is still normal.



Example

If X and Y are independent standard normal and a, b are constants, show that $aX + bY$ is still normal.

Proof: Do it on board!



1.4. Stochastic processes



1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?



1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?

It is a r.v. depending on time.



1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?

It is a r.v. depending on time. Or, it is a family of r.v.'s.



1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?

It is a r.v. depending on time. Or, it is a family of r.v.'s. For example, $X_1 = \$153$, $X_2 = \$120$, ...



1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?

It is a r.v. depending on time. Or, it is a family of r.v.'s. For example, $X_1 = \$153$, $X_2 = \$120$, ...

Time set: $\mathbb{T} = \{0, 1, 2, \dots, N\}$ or $\{0, 1, 2, \dots\}$, or $[0, T]$ or $[0, \infty)$.



1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?

It is a r.v. depending on time. Or, it is a family of r.v.'s. For example, $X_1 = \$153$, $X_2 = \$120$, ...

Time set: $\mathbb{T} = \{0, 1, 2, \dots, N\}$ or $\{0, 1, 2, \dots\}$, or $[0, T]$ or $[0, \infty)$.

State space S :



1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?

It is a r.v. depending on time. Or, it is a family of r.v.'s. For example, $X_1 = \$153$, $X_2 = \$120$, ...

Time set: $\mathbb{T} = \{0, 1, 2, \dots, N\}$ or $\{0, 1, 2, \dots\}$, or $[0, T]$ or $[0, \infty)$.

State space S : The set that the process takes its values,

1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?

It is a r.v. depending on time. Or, it is a family of r.v.'s. For example, $X_1 = \$153$, $X_2 = \$120$, ...

Time set: $\mathbb{T} = \{0, 1, 2, \dots, N\}$ or $\{0, 1, 2, \dots\}$, or $[0, T]$ or $[0, \infty)$.

State space S : The set that the process takes its values, e.g., $S = [0, \infty)$ in this example.

1.4. Stochastic processes

Example

What is the stock price of Apple (AAPL)?

It is a r.v. depending on time. Or, it is a family of r.v.'s. For example, $X_1 = \$153$, $X_2 = \$120$, ...

Time set: $\mathbb{T} = \{0, 1, 2, \dots, N\}$ or $\{0, 1, 2, \dots\}$, or $[0, T]$ or $[0, \infty)$.

State space S : The set that the process takes its values, e.g., $S = [0, \infty)$ in this example.

Definition

A stochastic process is a family of random variables $\{X_t : t \in \mathbb{T}\}$ with time set \mathbb{T} and state space S .



Example

Suppose that you will bid \$1 in each play. You will win \$1 with probability p and lose \$1 with probability $q = 1 - p$. You will stop if you have nothing to bid or you played 3 bids. If you start with \$1, what is the probability that you will end up bankrupted?



Example

Suppose that you will bid \$1 in each play. You will win \$1 with probability p and lose \$1 with probability $q = 1 - p$. You will stop if you have nothing to bid or you played 3 bids. If you start with \$1, what is the probability that you will end up bankrupted?

Solution: Do it on board!



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

HW: Ch1, 8, 10, 13, 36. Ch2, 1, 4, 5, 30, 33, 43, 46, 67, 69.

Note: Ch2 means Chapter 2 in the textbook.