

Step-1

Consider the function,

$$f(x) = \sin 2x.$$

The objective is to find the closest function $a \cos x + b \sin x$ to the function f on the interval from $-\pi$ to π .

Step-2

The closest function $a \cos x + b \sin x$ to the function $f(x)$ is $b_1 \sin 2x$.

Here, the coefficient b_1 is the least squares solution to the inconsistent equation $b_1 \sin 2x = f(x)$ and $b_1 = \frac{(f(x), \sin x)}{(\sin x, \sin x)}$.

The formula $b_1 = \frac{(f(x), \sin x)}{(\sin x, \sin x)}$ brings the $b_1 \sin 2x$ as close as possible to $f(x)$.

Now find the coefficient $b_1 = \frac{(f(x), \sin x)}{(\sin x, \sin x)}$ as follows:

$$\begin{aligned}(f(x), \sin x) &= \int_{-\pi}^{\pi} f(x) \sin x \, dx \\&= \int_{-\pi}^{\pi} \sin 2x \sin x \, dx \quad \text{Use } f(x) = \sin 2x \\&= 2 \int_0^{\pi} \sin 2x \sin x \, dx \quad \left(\begin{array}{l} \text{For any even function } f, \text{ i.e. } f(-x) = f(x), \\ \int_{-a}^a f(x) \, dx = 2 \int_0^a f(x) \, dx \\ \text{Here, } \sin 2x \cdot \sin x \text{ is an even function.} \end{array} \right) \\&= \int_0^{\pi} 2 \sin 2x \sin x \, dx\end{aligned}$$

Step-3

Further simplification is as follows:

$$\begin{aligned}
&= \int_0^{\pi} (\cos(2x+x) - \cos(2x-x)) \, dx \\
&\quad \text{(Use } 2 \sin x \sin y = \cos(x+y) - \cos(x-y) \text{)} \\
&= \int_0^{\pi} (\cos 3x - \cos x) \, dx \\
&= \left[\frac{\sin 3x}{3} - \sin x \right]_0^{\pi} \\
&= \frac{\sin 3\pi}{3} - \sin \pi - \frac{\sin(0)}{3} + \sin(0) \\
&= 0 \quad \text{Use } \sin n\pi = 0, \text{ for all } n = 1, 2, 3, \dots
\end{aligned}$$

Therefore, $(f(x), \sin x) = 0$.

Step-4

Now find the inner product $(\sin x, \sin x)$.

$$\begin{aligned}
(\sin x, \sin x) &= \int_{-\pi}^{\pi} (\sin x)(\sin x) \, dx \\
&= \int_{-\pi}^{\pi} \sin^2 x \, dx \\
&= 2 \int_0^{\pi} \sin^2 x \, dx \quad \text{Here, } \sin^2 x \text{ is an even function.} \\
&= 2 \int_0^{\pi} \left(\frac{1 - \cos 2x}{2} \right) \, dx \quad \text{Use } \cos 2x = 1 - 2 \sin^2 x
\end{aligned}$$

$$\begin{aligned}
&= \int_0^{\pi} (1 - \cos 2x) \, dx \\
&= \left[x - \frac{\sin 2x}{2} \right]_0^{\pi} \\
&= \pi - \frac{\sin 2\pi}{2} - 0 + \frac{\sin 2(0)}{2} \\
&= \pi - 0 \\
&= \pi
\end{aligned}$$

Therefore, $(\sin x, \sin x) = \pi$.

Step-5

Substitute the known values in the formula $b_1 = \frac{(f(x), \sin x)}{(\sin x, \sin x)}.$

$$\begin{aligned} b_1 &= \frac{(f(x), \sin x)}{(\sin x, \sin x)} \\ &= \frac{0}{\pi} \\ &= 0 \end{aligned}$$

$$b_1 \sin 2x = 0 \cdot \sin 2x$$

Therefore, the closest function is $= 0.$

Hence, the closest function to the function $f(x)$ is $\boxed{0 \cdot \sin 2x = 0}.$

Step-6

Now the objective is to find the closest straight line $c + dx$ to the function $f(x) = \sin 2x.$

The closest straight line $c + dx$ to the function f is $a_1 + b_1x.$

Here, the coefficients a_1, b_1 are given by the formulas,

$$a_1 = \frac{(f(x), \cos x)}{(\cos x, \cos x)} \text{ and } b_1 = \frac{(f(x), \sin x)}{(\sin x, \sin x)}.$$

From the above calculations, obtained that $b_1 = \frac{(f(x), \sin x)}{(\sin x, \sin x)} = 0.$

Step-7

Now find the value of the coefficient $a_1.$

First compute the inner product $(f(x), \cos x).$

$$\begin{aligned}
 (f(x), \cos x) &= \int_{-\pi}^{\pi} f(x) \cos x \, dx \\
 &= \int_{-\pi}^{\pi} \sin 2x \cos x \, dx \quad \text{Use } f(x) = \sin 2x \\
 &= 0 \quad \left(\begin{array}{l} \text{For any odd function } f, \text{ i.e. } f(-x) = -f(x), \\ \int_{-a}^a f(x) \, dx = 0. \\ \text{Here, } \sin 2x \cdot \cos x \text{ is an odd function.} \end{array} \right)
 \end{aligned}$$

Therefore, $(f(x), \cos x) = 0$.

Step-8

Now compute the inner product $(\cos x, \cos x)$.

$$\begin{aligned}
 (\cos x, \cos x) &= \int_{-\pi}^{\pi} (\cos x)(\cos x) \, dx \\
 &= \int_{-\pi}^{\pi} \cos^2 x \, dx \\
 &= 2 \int_0^{\pi} \cos^2 x \, dx \quad \text{Here, } \cos^2 x \text{ is an even function.} \\
 &= 2 \int_0^{\pi} \left(\frac{1 + \cos 2x}{2} \right) \, dx \quad \text{Use } \cos 2x = 2 \cos^2 x - 1
 \end{aligned}$$

$$\begin{aligned}
 &= \int_0^{\pi} (1 + \cos 2x) \, dx \\
 &= \left[x + \frac{\sin 2x}{2} \right]_0^{\pi} \\
 &= \pi + \frac{\sin 2\pi}{2} - 0 - \frac{\sin 2(0)}{2} \\
 &= \pi - 0 \\
 &= \pi
 \end{aligned}$$

Therefore, $(\cos x, \cos x) = \pi$.

Step-9

Substitute the known values in the formula $a_1 = \frac{(f(x), \cos x)}{(\cos x, \cos x)}$.

$$\begin{aligned} a_1 &= \frac{(f(x), \cos x)}{(\cos x, \cos x)} \\ &= \frac{0}{\pi} \\ &= 0 \end{aligned}$$

Substitute the values of a_1 and b_1 in the function $a_1 + b_1x$.

$$\begin{aligned} a_1 + b_1x &= 0 + 0x \\ &= 0 \end{aligned}$$

Hence, the closest straight line to the function f is $\boxed{0 + 0x = 0}$.