



南方科技大学  
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Please write carefully and clearly in complete sentences. Your explanations are your only representative when your work is being graded.

Unless otherwise noted, vector spaces are over  $\mathbb{F}$ , where  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{F} = \mathbb{C}$ .

1. Label the following statements as **True** or **False**. **Along with your answer, provide an informal proof, counterexample, or other explanation.**

- (1) If  $T : V \rightarrow W$  is a linear map and  $v_1, v_2, \dots, v_n$  are vectors in  $V$ , and  $Tv_1, Tv_2, \dots, Tv_n$  is linearly independent, then  $v_1, v_2, \dots, v_n$  is linearly independent.
- (2) Suppose that  $p_0, p_1, \dots, p_m$  are polynomials in  $\mathcal{P}_m(\mathbb{F})$  such that  $p_j(-1) = 0$  for all  $j$ , then  $p_0, p_1, \dots, p_m$  is not linearly independent in  $\mathcal{P}_m(\mathbb{F})$ .
- (3) Let  $T$  be a linear operator defined on a 3 dimensional real vector space, then  $T$  always has an eigenvalue.
- (4) If  $v_1, v_2$ , and  $v_3$  are eigenvectors of  $T$  such that  $v_3 = v_1 + v_2$ , then all three vectors have the same eigenvalue.
- (5) Let  $V$  be a complex vector space and  $U_1, U_2, U_3$  be its subspaces with intersection  $U_1 \cap U_2 \cap U_3 = \{\mathbf{0}\}$ . Then  $U_1 + U_2 + U_3$  is a direct sum.

2. If  $U_1$  and  $U_2$  are subspaces of a finite-dimensional vector space. Prove that

$$\dim(U_1 + U_2) = \dim U_1 + \dim U_2 - \dim(U_1 \cap U_2).$$

3. Define  $T \in \mathcal{L}(\mathcal{P}_3(\mathbb{R}))$  by

$$(Tp)(x) = xp'(x)$$

for all  $x \in \mathbb{R}$ . Find all eigenvalues and eigenvectors of  $T$ .

4. Define  $V^4$  by

$$V^4 = V \times V \times V \times V.$$

Prove that  $V^4$  and  $\mathcal{L}(\mathbb{F}^4, V)$  are isomorphic vector spaces.

5. Suppose  $V$  is finite-dimensional and  $S, T \in \mathcal{L}(V)$ . Prove that  $ST = I$  if and only if  $TS = I$ .
6. Consider the standard basis  $1, x, x^2$  of  $V = \mathcal{P}_2(\mathbb{R})$ . Let  $\varphi_1, \varphi_2, \varphi_3$  be corresponding the dual basis of  $\mathcal{P}_2(\mathbb{R})'$ . Let  $\varphi : V \rightarrow \mathbb{R}$  be the linear function

$$f(x) \mapsto f(2) + \int_0^1 f(x)dx.$$

Find the coefficients  $a, b, c$  for which  $\varphi = a\varphi_1 + b\varphi_2 + c\varphi_3$ .

7. Let  $V$  be a vector space that is generated by  $v_1, v_2, \dots, v_n$ , and let  $u_1, u_2, \dots, u_m$  be a linearly independent list of vectors in  $V$ . Show that  $m \leq n$  and there exists a subset  $H$  of  $v_1, v_2, \dots, v_n$  containing exactly  $n - m$  vectors such that  $H \cup \{u_1, u_2, \dots, u_m\}$  generates  $V$ .
8. Let  $V$  be a finite-dimensional complex vector space and let  $S$  and  $T$  be linear operators on  $V$  such that  $ST = TS$ . Prove that if  $S$  and  $T$  can each be diagonalized, then there is a basis for  $V$  which simultaneously diagonalizes  $S$  and  $T$ .